

Croat Med J. 2015;56:257-62
doi: 10.3325/cmj.2015.56.257

Identification of human remains from the Second World War mass graves uncovered in Bosnia and Herzegovina

Damir Marjanović^{1,2,3}, Negra Hadžić Metjahić⁴, Jasmina Čakar⁴, Mirela Džehverović⁴, Serkan Dogan¹, Elma Ferić¹, Snježana Džijan⁵, Vedrana Škaro^{5,6}, Petar Projić^{5,6}, Tomislav Madžar⁷, Eduard Rod⁸, Dragan Primorac^{6,8,9,10,11}

Aim To present the results obtained in the identification of human remains from World War II found in two mass graves in Ljubuški, Bosnia and Herzegovina.

Methods Samples from 10 skeletal remains were collected. Teeth and femoral fragments were collected from 9 skeletons and only a femoral fragment from 1 skeleton. DNA was isolated from bone and teeth samples using an optimized phenol/chloroform DNA extraction procedure. All samples required a pre-extraction decalcification with EDTA and additional post-extraction DNA purification using filter columns. Additionally, DNA from 12 reference samples (buccal swabs from potential living relatives) was extracted using the Qiagen DNA extraction method. Quantifiler™ Human DNA Quantification Kit was used for DNA quantification. PowerPlex ESI kit was used to simultaneously amplify 15 autosomal short tandem repeat (STR) loci, and PowerPlex Y23 was used to amplify 23 Y chromosomal STR loci. Matching probabilities were estimated using a standard statistical approach.

Results A total of 10 samples were processed, 9 teeth and 1 femoral fragment. Nine of 10 samples were profiled using autosomal STR loci, which resulted in useful DNA profiles for 9 skeletal remains. A comparison of established victims' profiles against a reference sample database yielded 6 positive identifications.

Conclusion DNA analysis may efficiently contribute to the identification of remains even seven decades after the end of the World War II. The significant percentage of positively identified remains (60%), even when the number of the examined possible living relatives was relatively small (only 12), proved the importance of cooperation with the members of the local community, who helped to identify the closest missing persons' relatives and collect referent samples from them.

¹International Burch University, Sarajevo, Bosnia and Herzegovina

²Institute for Anthropological Researches, Zagreb, Croatia

³Faculty of Science, University of Sarajevo, Sarajevo, Bosnia and Herzegovina

⁴Institute for Genetic Engineering and Biotechnology, University of Sarajevo, Sarajevo, Bosnia and Herzegovina

⁵Genos DNA Laboratory, Zagreb, Croatia

⁶University of Osijek, Medical School, Osijek, Croatia

⁷University of Zagreb, Medical School, Zagreb, Croatia

⁸St. Catherine Hospital, Zabok and Zagreb, Croatia

⁹Eberly College of Science, The Pennsylvania State University, University Park, PA, USA,

¹⁰University of Split, Medical School, Split, Croatia

¹¹Children's Hospital Srebrnjak, Zagreb, Croatia

Received: May 15, 2015

Accepted: June 9, 2015

Correspondence to:

Damir Marjanović
International Burch University,
Institute for Anthropological
Research
Francuske revolucije bb
71 000 Sarajevo, Bosnia and
Herzegovina
damir.marjanovic@ibu.edu.ba

DNA analysis plays a key role in the identification of missing persons and victims of mass fatality incidents, and DNA profiling could be regarded as a core method for such identification (1). The primary value of this procedure significantly increased over the last twenty years due to the introduction of short tandem repeat (STR) loci in routine forensic testing, especially their optimized versions, miniSTRs or STR markers located at the Y and X chromosomes. Data obtained by DNA typing are highly reliable and can be used as a powerful tool that produces reliable results (2).

Various procedures could be used to identify human remains (3) and the choice of an appropriate procedure and its usefulness primarily depend on the condition of the remains. When it comes to skeletal remains from World War II (WWII), DNA analysis has already proved to be efficient, and in most cases, the only applicable approach to human identification (4,5). Furthermore, a seven-decade long soil deposition is a significant adverse factor, which is why most of the laboratories use optimized procedures in the analysis of these samples. Over the last 20 years, DNA identification of victims of the wars in Bosnia and Herzegovina (B&H) (6) and Croatia in the 1990s (3), the analysis of WWII skeletal remains in Slovenia (4,5), and the analysis of several centuries old archeological samples (7) successfully identified a number of human skeletal remains and overcame a number of challenges, such as optimization of DNA extraction protocols, specific laboratory procedures, communication with the relatives, etc. Here we once again utilized the experience gained from these projects to identify skeletal remains from the WWII mass graves in B&H.

There are no precise official data about the number of killed and missing persons from the WWII period in B&H, although this number is roughly estimated to about 180 000 (8). This estimate is even higher if we include the victims from immediate post-war incidents and mass executions by communist authorities. Most of those crimes were hidden for almost 70 years. For example, more than 2300 persons from the Ljubuški municipality alone were killed during WWII, and immediately after it, that is, in 1945. Almost 50% of the victims are still considered missing and their burial sites are not known (9).

As a consequence of continuous pressure from living relatives, the regional authorities and local communities recently put in significant efforts to identify individuals discovered in several mass graves scattered throughout the region. They commissioned a DNA analysis of skeletal remains and reference samples. Here, we report the results of the identification of the WWII victims from the city of Ljubuški, south Herzegovina.

MATERIALS AND METHODS

Handling of skeletal remains

More than 60 skeletal remains from different locations in and around Ljubuški city (south Bosnia and Herzegovina) were exhumed during 2010 and 2011 (Figure 1. Exhumations were initiated by family members as well as the local Commission for Marking and Maintenance of Graves from World War II and the Post-War Period in the municipality of Ljubuški. The work was performed by forensic experts




FIGURE 1. Location of (A) Bosnia and Herzegovina and (B) Ljubuški area.

from the University Clinical Center Split, Croatia. Three years later mortal remains of 10 persons, which were exhumed from the uncovered graves closest to the center of the city Ljubuški, were selected for DNA analysis. Testimonies from living relatives indicated that these remains belonged to the inhabitants of Ljubuški that were shot by communist forces in the spring of 1945. According to these testimonies, local people knew about this burial site, but it could not be marked until the end of the communist rule in B&H.

The remains were exhumed and samples for DNA analysis (femoral fragments and teeth) were collected and labeled. The samples (teeth and bone fragments from nine bodies and bone fragment from one body) were subsequently transported to Genos DNA laboratory in Zagreb and the Laboratory of Forensic Genetics at the Institute for Genetic Engineering and Biotechnology (INGEB) University of Sarajevo, where they were stored at -80°C until further processing.

Sample preparation

For the bone sample, the entire exterior was sanded clean to remove potential contaminants. Bone surface was cleaned from remnant soft tissue and soil traces using a grinding stone attached to a Dremel[®] rotary tool. The sample was then successively washed in mild detergent, 5% bleach, sterile distilled water, and 100% ethanol and subsequently air-dried. The thoroughly dried sample was pulverized using a sterilized Waring[®] blender and powder and then transferred to sterile 15-mL conical polypropylene tubes. With the exception of the sanding phase, the same procedure was applied to teeth. An EDTA decalcification step was used for each of the samples. Also, the same extractions were performed for each sample following a previously described phenol-chloroform DNA extraction protocol (10). DNA analysis included teeth from 9 skeletons (laboratory code S1 to S9/03-14) and a femoral fragment from one skeleton (laboratory code S10/03-14) because of absence of teeth. Filter units were used for DNA purification and concentration. The concentrates were subsequently transferred to 1.5-mL microcentrifuge tubes and diluted with DNA-free ddH_2O to a final volume of maximum 100 μL .

DNA analysis

The DNA concentration was determined using Quantifiler Human DNA Quantification Kit (Applied Biosystems Foster

City, CA, USA) (11). The reaction was carried out in the AB 7300 Real-Time PCR System (Applied Biosystems) according to manufacturer's recommendations.

The PowerPlex[®]ESI kit (Promega Corp., Madison, WI, USA) was used to simultaneously amplify 15 STR loci: D3S1358, D8S1179, D18S51, D21S11, FGA, TH01, vWA, D2S441, D10S1248, D22S1045, D1S1656, D12S391, D2S1338, D16S539, D19S433 as well as the gender determination locus, Amelogenin. Amplification was carried out as described previously (12). The total volume of each reaction was 25 μL . Additionally, the PowerPlex[®]Y23 kit (Promega Corp.) was used to simultaneously amplify 23 Y-STR loci according to manufacturer's recommendations (13). PCR amplification was carried out using the Applied Biosystems[®] GeneAmp[®] PCR System 9700 (Life Technologies) according to manufacturer's recommendations. Electrophoresis of the amplification products was performed on an ABI PRISM 310 Genetic Analyzer (Applied Biosystems). The raw data were compiled and analyzed using 310 Data Collection Software and GeneMapper[™] 3.2.

Handling of referent samples

Referent samples (12 buccal swabs) from potential living relatives were collected, recorded, and preliminarily labeled by local DNA experts. Living relatives were directly contacted by a local person (Mr Jure Lauc), who was present during the collection of each sample. Collection was performed with an Internal GENOS collection kit, and dried and labeled samples were transported to the laboratory. Upon arrival, the samples were relabeled and the relevant information entered into the Chain of Custody forms. The samples were stored at -80°C until DNA extraction was performed using the QIAmp DNA Mini Kit (14).

PowerPlex[®] ESI kit (Promega Corp.) was used for further analysis. Similar amounts of DNA were used in all PCR reactions. Amplification was carried out as described previously (12). The total reaction volume was 10 μL . Additionally, PowerPlex[®]Y23 kit (Promega Corp.) was used to simultaneously amplify 23 Y-STR loci according to manufacturer's recommendations (13). PCR amplification was carried out by the Applied Biosystems[®] GeneAmp[®] PCR System 9700 (Life Technologies) according to manufacturer's recommendations. Electrophoresis of the amplification products was performed on an ABI PRISM 310 Genetic Analyzer (Applied Biosystems). The raw data were compiled and analyzed using 310 Data Collection Software and GeneMapper[™] 3.2.

Statistical analysis

Comparative analysis of DNA profiles obtained from skeletal remains and referent samples, estimation of potential familial relationships, calculation of paternity and sibling indexes, as well as calculation of matching probability were performed according to the previously used statistical approach (15).

RESULTS

Ten skeletal remains samples (9 teeth and 1 bone) were processed. The quantification results illustrated that the mean concentration of isolated DNA varied between 4.00×10^{-3} ng/ μ L and 7.00×10^{-1} ng/ μ L, for samples from which DNA profiles were obtained. Based on the quantification results, an extended PCR procedure (up to 32 cycles with an extended elongation time) was recommended for 3 samples.

A total of 9 DNA profiles were obtained, which means that profiling success rate was 90% (9/10 remains; Table 1). The number of successfully analyzed loci per established STR profile varied from 12 to 16. Also, 12 referent buccal swabs collected from the missing persons' relatives were 100% successfully profiled.

Comparison of victims' profiles against the referent database resulted in 6 positive identifications. All identified persons were men and first 5 were identified by autosomal DNA analysis, while 1 was identified by a combination of autosomal and Y-STR analysis. Five of the 6 identified persons were identified using the living children's DNA profiles and 1 was identified using the living nephew's DNA profile in combination with the established DNA profile of

his deceased brother who was buried in the same grave and was one of the 5 previously mentioned identified persons (Table 1).

DISCUSSION

This study demonstrated that the experience gathered over the last two decades through identification of missing persons in this region could be almost routinely applied without significant modifications in the analysis of WWII skeletal remains. DNA analysis successfully identified remains even seven decades after the end of the World War II and proved to be the essential approach for the identification of remains from that period.

Genetic typing by STR loci analysis has over the last two decades (3-5) become a primary method of choice for the identification of human remains. The latest studies conducted on the WWII samples (4,5,16-19) were done using STR techniques instead of mtDNA, which used to be the mostly used identification method for the samples from that period (20).

Nowadays, application of improved old or completely new procedures increases the possibility for nuclear DNA profiling of degraded skeletal remains and gives us an opportunity to increase success of the skeletal remains DNA profiling routine procedures. Additionally, these results emphasize the importance of targeted, prompt, and efficient sample collection from living relatives. This approach yielded a 60% success identification rate, which is considerably higher than rate from our previous studies (4,5). The crucial reason for this success was the involvement of individuals from the local community who searched for victims' close relatives. This is why we had so many samples from the di-

TABLE 1. Summary of DNA analysis of skeletal remains

Body code	Sample code	Collected samples	EDTA decalc. step	Polymerase chain reaction protocol	Successfully DNA profiled	Number of loci with detected alleles	Type of analysis	Positive identification	Identified by DNA profile from
G2K1	S1/03-14	Tooth/bone	+	Standard	Yes	14/16	Autosomal	No	-
G1K2	S2/03-14	Tooth/bone	+	Extended	Yes	15/16	Autosomal	Yes	Daughter
G1K6	S3/03-14	Tooth/bone	+	Standard	Yes	14/16	Autosomal/ Y chromosomal	Yes	Living nephew/ deceased brother
G1K4	S4/03-14	Tooth/bone	+	Standard	Yes	15/16	Autosomal	Yes	Son
G1K5	S5/03-14	Tooth/bone	+	Standard	Yes	14/16	Autosomal	Yes	Son
G2K3	S6/03-14	Tooth/bone	+	Standard	Yes	12/16	Autosomal	No	-
G1K1	S7/03-14	Tooth/bone	+	Standard	Yes	16/16	Autosomal	Yes	Daughter
G1K3	S8/03-14	Tooth/bone	+	Standard	Yes	15/16	Autosomal	Yes	Son
G2K6	S9/03-14	Tooth/bone	+	Extended	Yes	13/16	Autosomal	No	-
G2K2	S10/03-14	Bone	+	Extended	No	3/16	Autosomal	No	-

rect offspring (daughters and sons) of the victims, which led to statistically satisfying and powerful results of the identification although the number of processed referent samples was small. This was something that our previous studies (4,5) lacked and that had a significant influence on the final success rate of identification.

The biggest limitation of this type of studies is the lack of clear strategy for identification of WWII victims. Therefore, we suggest that our strategy should be used as a model for further activities in this field, as well for additional DNA identification of human remains exhumed from the WWII mass graves in B&H and other neighboring countries. Once again it was proven that forensic science might bring closure to families who had been searching for their loved ones for decades. It is important to highlight that many of the closest relatives of the missing are aged and it is urgent to collect samples from them. It is safe to assume that without their DNA profile the entire identification process would be much more complicated if not impossible.

Acknowledgments We thank the Lauc family, especially Mr Jure Lauc. Without his enormous effort and true commitment to this mission, skeletal remains of the victims would be still hidden nameless in the mass graves.

Funding DNA analysis was financially supported by the living relatives.

Ethical approval obtained from the Ethical Committee of the International Burch University, Sarajevo.

Declaration of authorship DM, leader of the project, prepared and approved the final version of the manuscript. NHM took part in lab-work and sub-coordination of the project. MDŽ, JČ, VŠ, SDŽ, PP took part in laboratory work with samples and generation of DNA profiles. EF, SD performed statistical analysis. TM took part in manuscript preparation and collection of the historical data from the area. ER took part in optimization of the handling procedures with bone samples. DP took part in supervision of the project and finalization of the manuscript.

Competing interests All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- Butcher BA, Bieher FR, Budimilija ZM, Dennis SM, Desire MA. Identification of missing persons and mass disaster victim identification by DNA. In: Primorac D, Schanfield MS, editors. *Forensic DNA application: An interdisciplinary perspective*. London, UK: CRC Press; 2014. p. 277-92.
- Butler JM. *Fundamentals of forensic DNA typing*. London, UK: Elsevier Academic Press; 2010.
- Anđelinović S, Sutilović D, Erceg Ivkošić I, Škaro V, Ivkošić A, Paić F, et al. Twelve-year experience in identification of skeletal remains from mass graves. *Croat Med J*. 2005;46:530-9. [Medline:16100755](#)
- Marjanović D, Durmić-Pašić A, Bakal N, Haverić S, Kalamujić B, Kovačević L, et al. DNA Identification of Skeletal Remains from the Second World War Mass Graves Uncovered in Slovenia. *Croat Med J*. 2007;48:513-9. [Medline:17696306](#)
- Marjanović D, Durmić-Pašić A, Kovačević L, Avdić J, Džehverović M, Haverić S, et al. DNA Identification of skeletal remains of communist armed forces victims during and after World War II: combined Y-chromosome short tandem repeat (STR) and miniSTR approach. *Croat Med J*. 2009;50:296-304. [Medline:19480024](#) [doi:10.3325/cmj.2009.50.296](#)
- Definis-Gojanović M, Ivanović J, Drmić I, Galić M, Anđelinović Š. Identification of fifty-nine victims of the war from the Kupres battlefield, Bosnia and Herzegovina. *Croat Med J*. 1995;36:61-4.
- Marjanović D, Kovačević L, Čakar J, Džehverović M, Musemić Dž, Bajrović K. DNA analysis of ancient skeletal remains from old Bosnian graves. 7th ISABS Conference in Forensic Genetic, Anthropologic and Medical genetics and Mayo Clinic Lectures in Translational Medicine, Bol, Croatia, June 20-24, 2011 (Book of Abstracts:60).
- Krušelj Ž, Zagorac Đ. *Sporna knjiga mrtvih. Danas.1989* [in Serbian]. Novembar 12; p24.
- Lučić I. Pronalaženje posmrtnih ostataka ubijenih u Drugome Svjetskom Ratu i poraču te doprinos vicepostulature postupka mučeništva "Fra Leo Petrović i 65 subraće" [in Croatian]. on 3rd Symposium "Stopama Pobjenih", ŠirokiBrijeg (Oct 2, 2011). Available from: <http://www.pobjeni.info/izdavastvo/izdanje/24>. Accessed: May 14, 2014.
- Davoren J, Vanek D, Konjhodžić R, Crews J, Huffine E, Parsons TJ. Highly effective DNA extraction method for nuclear short tandem repeat testing of skeletal remains from mass graves. *Croat Med J*. 2007;48:478-85. [Medline:17696302](#)
- Applied Biosystems. *Quantifiler®Kits User's Manual*. Applied Biosystems; 2012.
- Promega Corporation. *GenePrint®PowerPlex ESI system instruction for use of Product*. Promega Corporation; 2014.
- Promega Corporation. *GenePrint®PowerPlex Y23 System Instruction for use of Product*. Promega Corporation; 2015.
- Qiagen Companies. *QIAmp DNA Mini Kit Handbook*. Qiagen; 2012.
- Schanfield MS, Primorac D, Marjanović D. Identification of missing persons and forensic DNA analysis and statistics. In: Primorac D, Schanfield MS, editors. *Forensic DNA Application: An interdisciplinary perspective*. London, UK: CRC Press; 2014. p. 277-92.
- Zupanić-Pajnić I, Gornjak-Pogorelc B, Balažić J, Zupanc T, Stefanić B. Highly efficient nuclear DNA typing of the World War II skeletal remains using three new autosomal short tandem repeat amplification kits with the extended European Standard Set of loci. *Croat Med J*. 2012;53:17-23. [Medline:22351574](#) [doi:10.3325/cmj.2012.53.17](#)
- Zupanić-Pajnić I, Gornjak-Pogorelc B, Balažić J. Molecular genetic identification of skeletal remains from the Second world war

- konfin I mass grave in Slovenia. *Int J Legal Med.* 2010;124:307-17. [Medline:20217112](#) [doi:10.1007/s00414-010-0431-y](#)
- 18 Primorac D, Andelinovic S, Definis-Gojanovic M, Drmic I, Rezic B, Baden MM, et al. Identification of war victims from mass graves in Croatia and Bosnia and Herzegovina through the use of DNA typing and standards forensic methods. *J Forensic Sci.* 1996;41:891-4. [Medline:8789853](#)
- 19 Alonso A, Andelinovic S, Martin P, Sutlovic D, Erceg I, Huffine E, et al. DNA typing from skeletal remains: evaluation of multiplex and megaplex str systems on DNA isolated from bone and teeth samples. *Croat Med J.* 2001;42:260-6. [Medline:11387635](#)
- 20 Palo JU, Hedman M, Solderholm N, Sajantila A. Repatriation and identification of Finnish World War II soldiers. *Croat Med J.* 2007;48:528-35. [Medline:17696308](#)