

Marked increase in rat red blood cell membrane protein glycosylation by one-month treatment with a cafeteria diet

Laia Oliva¹, Cristian Baron¹, José-Antonio Fernández-López^{1,2,3}, Xavier Remesar^{1,2,3} and Marià Alemany^{1,2,3}

¹ Department of Nutrition and Food Science, Faculty of Biology, University of Barcelona, Barcelona, Spain

² Institute of Biomedicine of the University of Barcelona, Barcelona, Spain

³ CIBER OBN, Barcelona, Spain

ABSTRACT

Background and Objectives. Glucose, an aldose, spontaneously reacts with protein amino acids yielding glycosylated proteins. The compounds may reorganize to produce advanced glycosylation products, which regulatory importance is increasingly being recognized. Protein glycosylation is produced without the direct intervention of enzymes and results in the loss of function. Glycosylated plasma albumin, and glycosylated haemoglobin are currently used as index of mean plasma glucose levels, since higher glucose availability results in higher glycosylation rates. In this study we intended to detect the early changes in blood protein glycosylation elicited by an obesogenic diet.

Experimental Design. Since albumin is in constant direct contact with plasma glucose, as are the red blood cell (RBC) membranes, we analyzed their degree or glycosylation in female and male rats, either fed a standard diet or subjected to a hyper-energetic self-selected cafeteria diet for 30 days. This model produces a small increase in basal glycaemia and a significant increase in body fat, leaving the animals in the initial stages of development of metabolic syndrome. We also measured the degree of glycosylation of hemoglobin, and the concentration of glucose in contact with this protein, that within the RBC. Glycosylation was measured by colorimetric estimation of the hydroxymethylfurfural liberated from glycosyl residues by incubation with oxalate.

Results. Plasma glucose was higher in cafeteria diet and in male rats, both independent effects. However, there were no significant differences induced by sex or diet in either hemoglobin or plasma proteins. Purified RBC membranes showed a marked effect of diet: higher glycosylation in cafeteria rats, which was more marked in females (not in controls). In any case, the number of glycosyl residues per molecule were higher in hemoglobin than in plasma proteins (after correction for molecular weight). The detected levels of glucose in RBC were lower than those of plasma, even when expressed in molal units, and were practically nil in cafeteria-diet fed rats compared with controls; there was no effect of sex.

Conclusions. RBC membrane glycosylation is a sensitive indicator of developing metabolic syndrome-related hyperglycemia, more sensitive than the general

Submitted 27 March 2015

Accepted 22 June 2015

Published 16 July 2015

Corresponding author

Marià Alemany, malemany@ub.edu

Academic editor

Daniela Foti

Additional Information and
Declarations can be found on
page 11

DOI 10.7717/peerj.1101

© Copyright
2015 Oliva et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

measurement of plasma or RBC protein glycosylation. The extensive glycosylation of blood proteins does not seem to be markedly affected by sex; and could be hardly justified from an assumedly sustained plasma hyperglycemia. The low levels of glucose found within RBC, especially in rats under the cafeteria diet, could hardly justify the extensive glycosylation of hemoglobin and the lack of differences with controls, which contained sizeable levels of intracellular glucose. Additional studies are needed to study the dynamics of glucose *in vivo* in the RBC to understand how such extensive protein glycosylation could take place.

Subjects Cell Biology, Diabetes and Endocrinology, Hematology, Metabolic Sciences

Keywords Erythrocyte, Protein glycosylation, Glycemia, Glycosylated hemoglobin, Blood cell membrane, Cafeteria diet, Glycosylation

INTRODUCTION

Glucose, in addition to being the main intercellular energy staple, is a reducing aldose. Thus, it may easily react with a number of chemical groups in proteins and other biological compounds. The direct condensation with protein free amino groups (Maillard reactions) (*John & Lamb, 1993*) is fairly common, to the degree that a significant proportion of circulating plasma proteins are glycosylated (*Gragnoli et al., 1982*), as well as proteins in the red blood cell (RBC) membrane (*Miller, Gravallesse & Bunn, 1980*) and the hemoglobin they contain (*Bunn, Gabbay & Gallop, 1978*). The proportion of hemoglobin glycosylated in the terminal valine of chain B (Hb_{1AC}) is currently used as an index of overall exposure to free plasma glucose over time (*Siu & Yuen, 2014*). Glycosylation products may undergo Amadori reorganizations, producing a number of complex compounds known as advanced glycosylation products (AGP) (*Henning et al., 2011*), which play a significant role in the control of substrate utilization (*Wu et al., 2011*), cell function (*Guo et al., 2012*) and inflammation (*Poulsen et al., 2014*).

The chemical reactivity of glucose is often overlooked because of its overwhelming function in energy supply and rapid turnover, but direct non-enzymatic glycosylation remains a common mechanism of alteration of protein function and interference in signaling pathways (*Asahi et al., 2000; Itkonen & Mills, 2013*). It is commonly accepted that higher sustained circulating levels of glucose, as in diabetes, result in increased proportions of glycosylated proteins in plasma, RBCs and endothelial cells, Hb_{1AC} being a case in point (*Carson et al., 2010*). In fact, equations based on the correlation between mean estimated plasma glucose concentration and Hb_{1AC} proportion are currently in use (*Borg & Kuenen, 2009; Nathan et al., 2008*).

The self-selected cafeteria diets (*Sclafani & Springer, 1976*) are essentially hyperlipidic (*Prats et al., 1989*), and its consumption by rats causes hyperphagia, insulin resistance and obesity (*Correa Pinto & Monteiro Seraphim, 2012; Prats et al., 1989*). Exposure for one month of young adult rats to a cafeteria diet induces a number of metabolic changes that are in the limit of normalcy and correspond to the initial stages of the metabolic syndrome

(Romero *et al.*, 2010). The effects are more marked in male than in female rats (Romero *et al.*, 2012), probably because of the anti-inflammatory effects of estrogen (Thomas *et al.*, 2003); but, in any case, the obesity is already patent. Short-term treatment with cafeteria diets induce a mild hyperglycemia and hyperinsulinemia (Romero *et al.*, 2010), but not frank diabetes, which is more developed after prolonged exposure (Castell-Auvi *et al.*, 2012; Correa Pinto & Monteiro Seraphim, 2012).

In the present study, we have analyzed whether the glycosylation degree of total plasma or RBC proteins, as well as those of RBC membranes, are a direct correlate of their prolonged contact with plasma glucose in an early stage of development of hyperglycemia. We wanted, also, to check whether sex exerts any influence on the glycosylation response to comparable glucose concentrations.

MATERIALS AND METHODS

Animals and animal handling

All animal handling procedures were carried out in accordance with the norms of the European, Spanish and Catalan Governments. The study was specifically approved (DMAH-5483) by the Animal Ethics Committee of the University of Barcelona.

Wistar adult male and female (9 week-old) rats were used (Harlan Laboratories Models, Sant Feliu de Codines, Spain). The rats were adapted to the Animal House environment for at least one week prior to the beginning of the experiment, and were fed a standard (Harlan, type 2014) chow. The rats were kept in solid-bottomed adjoining collective 2-rat cages, with wood shavings as bedding material. Half of the rats in each group were subjected to an energy-rich limited-item cafeteria diet (Ferrer-Lorente *et al.*, 2005) for a month. The items of cafeteria diet (plain cookies spread with liver pâté, bacon, standard chow, water and whole cow's milk containing 300 g/L sucrose and a mineral and vitamin supplement) were renewed daily. Food consumption per cage and rat weights were recorded every day.

The four experimental groups ($N = 6$ for each) were: female-control (FC), female-cafeteria, (FCAF) male-control (MC) and male-cafeteria (MCAF). On day 29, a small sample of blood was taken from a cut in the rat tail's tip, centrifuged in capillary tubes, and the plasma was frozen for later measurement of glucose levels.

At the end of the experiment (day 30), the rats were anaesthetized with isoflurane and killed by exsanguination (blood drawn from the aorta using a dry-heparinized syringe). Part of the blood was centrifuged immediately (at $1,300\times g$ for 25 min and $2-4^{\circ}\text{C}$). Plasma and packed cells were frozen and kept at -20°C . A sample of fresh blood was deproteinized with 0.5 volumes of 6.7 M perchloric acid, mixed, neutralized with 4.5 M KOH containing 1.55 M potassium bicarbonate, centrifuged again at the same speed (at 4°C), and the supernatants used for the measurement of total blood glucose.

Packed cell volume was estimated from the weight of blood before centrifugation, that of plasma obtained after that centrifugation and the (redundant) weight of packed cells sedimented. Since the densities of cells and packed cells were known, and the proportion of packed cell volume was a direct correlate of time and acceleration generated during centrifugation, we used the previously described graphs, obtained under the same

conditions (Romero *et al.*, 2012) to estimate the actual proportion of plasma trapped between the cells, and thus determine the real packed cell volume.

A known weight of frozen packed cells was suspended in 10 volumes of chilled pure water. After gentle shaking for 20 min at 4 °C, the suspension was centrifuged 10 min at 2,000× *g* and 2–4 °C. The clear supernatant (hemoglobin and cytosolic RBC proteins) was used for the analyses of total and glycosylated protein.

RBC membrane separation

About 0.5 g samples of frozen packed cells were weighed and suspended in 3.5 mL of chilled tris–HCl buffer 10 mM pH 7.4, the cells were gently stirred until a uniform solution was obtained. Then, 4 mL of chilled 250 mM glucose were added and gently mixed. After standing 15 min (Tomoda *et al.*, 1984), the suspension was coarse-filtered through a small wad of glass fiber to remove debris, and then was centrifuged for 3 min at 8,000× *g* in the cold (2–4 °C). The fluffy precipitate was suspended in medium, and centrifuged again. A small translucent sediment of RBC membranes was obtained; it was weighed, and used for the analysis of protein, total phosphate and glycosylation.

Chemical analyses

Glucose in plasma and deproteinized fresh blood was measured with a glucose oxidase kit (Biosystems, Barcelona, Spain), supplemented with mutarotase (490 nkat/mL of reagent) (Calzyme, San Luis Obispo, CA, USA). Mutarotase was added to speed up epimerization equilibrium of α - and β -D-glucose and thus facilitate the oxidation of β -D-glucose by glucose oxidase (Miwa *et al.*, 1972). The enzyme addition was complemented with a precise control of the time (15 min) and temperature (30 °C) conditions of development of the reaction, in order to make sure all glucose in the sample was oxidized to gluconate. Protein content was estimated with a variant of the Lowry method (Lowry *et al.*, 1951) using fatty acid-free bovine serum albumin (Sigma, St Louis MO, USA) as standard.

RBC membranes were mineralized with perchloric acid (700 g/L) in 15 mL Teflon-stoppered glass tubes, in a dry block heater, at 150 °C for 24 h (Stein & Smith, 1982). Aliquots of the clear mineralized samples were used, after centrifugation, for the estimation of phosphate using the phosphomolybdate reaction using sodium mono-phosphate as standard (Gomori, 1942; Stein & Smith, 1982). A standard of phosphatidyl-choline (Sigma) was processed along with the samples. The measurements of phosphate from the phosphatidyl-choline standards proved that mineralization was complete (98–101%). Each batch of samples was corrected using their own standards, ran in parallel.

The degree of glycosylation was estimated by direct measurement of the 5-hydroxymethylfurfural (HMF) liberated by treatment of the samples with 1 N oxalic acid at 100 °C for 24 h (Gabbay *et al.*, 1979) in 15 mL Teflon-stoppered tubes set in a dry heating block. After cooling, trichloroacetic acid was added (final concentration 100 g/L), and the tubes were shaken and centrifuged for 15 min at 5,000× *g*. The precipitate was discarded. The amount of HMF released was measured through the condensation of HMF with 50 mM thiobarbituric acid (Sigma) (Gabbay *et al.*, 1979). After 20 min at 37 °C for development of color, the OD was measured at 443 nm, using blanks and pure HMF

Table 1 Body weight changes, energy intake and plasma glucose of Wistar rats fed control or cafeteria diet for 30 days. The data are the mean \pm sem of 6 animals per group. Plasma glucose was measured on day 29. Statistical significance of the differences between means were determined using a 2-way ANOVA program.

Parameter	Units	Male		Female		P values		
		Control	Cafeteria	Control	Cafeteria	Sex	Diet	Interaction
Initial weight	g	394 \pm 9	379 \pm 3	238 \pm 5	217 \pm 4	<0.0001	NS	NS
Final weight	g	474 \pm 10	511 \pm 3.5	275 \pm 11	290 \pm 8	0.0068	<0.0001	NS
Weight increase	g/30 d	82 \pm 10	137 \pm 4	41 \pm 5	74 \pm 7	<0.0001	<0.0001	NS
Eenergy intake	MJ/30 d	8.62 \pm 0.04	21.4 \pm 1.5	6.32 \pm 0.39	18.0 \pm 1.0	0.0055	<0.0001	NS
	W	3.33 \pm 0.01	8.26 \pm 0.59	2.44 \pm 0.15	6.93 \pm 0.38			
Plasma glucose	mM	7.58 \pm 0.32	9.13 \pm 0.15	6.83 \pm 0.26	8.53 \pm 0.12	0.0082	<0.0001	NS

(Sigma) standards, and was used to determine the HMF (i.e., unaltered glycosyl residues in proteins) in each sample.

Blood cell glucose estimation

Blood glucose is the composite of the glucose carried by the cells and that in plasma using the common (Higgins, Garlick & Bunn, 1982) formula:

$$\text{blood glucose} = \text{plasma glucose} \times (1 - \text{PCV}) + \text{cell glucose} \times \text{PCV}$$

where PCV (Packed Cell Volume) is the net cell volume fraction (i.e., discounting trapped plasma volume) of total blood volume (in this case = 1). In that equation, we had, for each rat, the PCV value as well as plasma and blood glucose. Crude cell-transported glucose was derived from these data. Since it was assumed that trapped plasma glucose concentration was the same than in plasma obtained by centrifugation, the glucose present in that plasma fraction was discounted from the total packed cell glucose (and added to the final data for “plasma glucose”). These calculations were carried out for each individual rat, thus all data used for the calculations were homologous.

Statistics

Statistic comparison between groups was carried out using 2- and 3-way ANOVA analyses, and the Bonferroni *post-hoc* test for further differences between specific groups (Prism 5 program; GraphPad Software, La Jolla, CA, USA).

RESULTS

Table 1 presents the changes in body weight experienced by the rats during one month of exposure to a cafeteria diet. The initial weight difference between female and male rats widened with time, since control males increased about 20% of their weight, compared with 15% of females; cafeteria diet increased body weight 35% in males and 34% in females. Males ate more energy than females: 36% (control diet) or 19% (cafeteria diet). Males' food (expressed as energy content) intake was 2.5-fold higher in cafeteria than in control diet; the value for females was 2.8 \times .




Figure 1 Distribution of blood glucose in plasma and cell compartments of Wistar rats fed control or cafeteria diet during 30 days. Data are the mean \pm sem of 6 animals per group (killed under isoflurane anesthesia), and were calculated from the data presented in Table 2. Statistical significance (two-way ANOVA) of the differences between groups: No differences were found for “sex”, but “diet” showed $P < 0.0001$ for cells and was not significant for plasma. Blood cell data for cafeteria diet were not statistically different from zero.

Standard plasma glucose (measured on day 29) showed both an effect of sex (female levels being lower) and diet (cafeteria diet data being higher).

Table 2 shows the data obtained from the analysis of blood extracted under isoflurane anesthesia. In this case, all plasma glucose data were higher than those obtained on day 29 under basal conditions, and there were no statistical differences between the groups attributable to sex or diet. Total blood glucose values were lower than those of plasma, and showed neither differences by sex or diet. However, the estimated data for cell glucose showed a clear effect of diet (Fig. 1). In both groups of cafeteria rats, the levels were minimal, not statistically different from zero, while those of rats under the control diet were lower than in plasma but clearly positive, the differences being not significant for “sex” but significant for “diet”. In control rats, when water content of plasma (about 92%) and packed cells (about 70%) was included in the calculations, the molal concentrations of cell glucose were in the range of 1/3rd of those of plasma; female rats presented similar values. Cafeteria diet-fed rats showed values in the range of only 4–7%.

The proportions of glycosylated protein, both in RBC and in plasma, are presented in Fig. 2. No significant differences were observed between the groups for “sex” and “diet”. However, cell protein was more heavily glycosylated than plasma proteins. In the case of cells, since most of the protein (>95%) is hemoglobin, it can be assumed that most glycosyl residues were bound to this protein; since its molecular weight (tetramer) is about 64,000, the molar ratio of HMF to hemoglobin was about 320, i.e., about 80 glycosyl residues per hemoglobin subunit. This value is about six-fold higher than the number of sites representing 7% Hb_{1AC}, which is limited to the terminal chains of hemoglobin. In the case of plasma, since albumin makes about 55% of plasma proteins and its molecular weight is close to 66,500, we obtain about 90 glycosyl residues per molecule. Evidently, this is only an imprecise approximation but shows that under the particular conditions of this




Figure 2 Degree of glycosylation expressed in nmol HMF per mg total protein in the cells and plasma of Wistar rats fed control or cafeteria diet during 30 days. Data are the mean \pm sem of 6 animals per group. Statistical significance (three-way ANOVA) of the differences between groups: No differences were found for “sex” and “diet”, but the differences between “compartments” (i.e., blood cells vs. plasma) was $P < 0.0001$.

Table 2 Blood glucose and packed cell volume of Wistar rats fed control or cafeteria diet for one month, after exsanguination under isoflurane anesthesia on day 30. The data are the mean \pm sem of 6 animals per group. Packed cell volume data were corrected for trapped plasma as explained in the text. Statistical significance of the differences between means were determined using a 2-way ANOVA program.

Parameter	Units	Male		Female		P values		
		Control	Cafeteria	Control	Cafeteria	Sex	Diet	Interaction
Blood glucose	mM	6.83 \pm 0.13	5.93 \pm 0.37	6.43 \pm 0.14	6.31 \pm 0.31	NS	NS	NS
Plasma glucose	mM	10.41 \pm 0.33	10.90 \pm 0.64	10.71 \pm 0.63	10.84 \pm 0.64	NS	NS	NS
Packed cell volume	% blood volume	45.7 \pm 0.9	43.8 \pm 0.7	43.1 \pm 1.1	42.7 \pm 1.8	NS	NS	NS
Blood cell glucose ^a	μ mol/g	2.75 \pm 0.32	0.53 \pm 0.96*	2.76 \pm 0.56	0.33 \pm 0.27*	NS	<0.0001	NS

Notes.

^a Blood cells' glucose concentration was calculated for each animal from glucose data (whole blood and plasma) and the net packed cell volume.

* Statistically not different from zero.

experiment, protein glycosylation was significant and about 3.5 times more intensive in cells (on a molar ratio) than in plasma proteins as a whole.

Figure 3 depicts the rate of glycosylation observed in purified membranes of blood cells. Since purification of membranes is not even close to quantitative, we could not determine in which proportion RBC membranes were glycosylated. In fact, we were not able to ascertain the degree of the purity of samples. Thus, membrane proteins could be contaminated by hemoglobin (in spite of the appearance of total elimination at the expense of dwindling recovery of membranes), spectrin or other molecules. Thus, we decided to also relate the degree of glycosylation to phospholipid, an exclusive membrane component in RBC. The molar ratio of released HMF to phospholipid phosphate (Fig. 3) showed an image quite different from that of Fig. 2. There were no statistical differences between groups attributable to “sex”. This was clear for control diet, but the post-hoc test showed a significant ($P < 0.05$) sex-related difference in cafeteria-fed rats. The effect of “diet” was significant, with several-fold higher values in cafeteria- than in control-fed rats.




Figure 3 Degree of glycosylation of blood cell membranes, expressed as mmoles of HMF per mmol of phospholipid *P* or unit of membrane protein weight, of Wistar rats fed control or cafeteria diet during 30 days. Data are the mean \pm sem of 6 animals per group. Statistical significance (two-way ANOVA) of the differences between groups: When analyzed for HMF vs. protein, no statistical differences were found. When analyzed for HMF vs. phospholipid *P*. No differences were found for “sex” but the difference for “diet” was $P < 0.0001$. There was a significant interaction between sex and diet. Glycosylation was higher ($P < 0.05$, Bonferroni post-hoc test) in the female cafeteria rats, compared with males.

Presentation of the data of HMF per mg of membrane preparation protein shown in Fig. 3, yields almost the same pattern, but statistical significance was lower because the individual variation of data was higher.

DISCUSSION

In the development of this apparently simple study, we tried to maintain a close control of methodology, since the problems of glycosylation of blood components have generated a sizeable number of studies, but their integrated analysis is scarce, in a way that only limited comparisons have been studied. We intended to present homologous data for plasma and RBC proteins, including also samples of RBC membranes, and using a model in which the metabolic syndrome pathologies, especially insulin resistance and hyperglycemia, were not fully set in.

The problem of anesthesia as hyperglycemic agent (Arola et al., 1981; Zuurbier et al., 2008) has not been solved; we opted by using this avenue to obtain sufficient blood to carry out all the compartmentation and membrane experiments in the same samples. Consequently we had to obtain separate plasma samples to compare the basal results with previous studies (Palou et al., 1980). The changes elicited by cafeteria diet agree with previously published studies (Ferrer-Lorente et al., 2005). We assumed that the brief isoflurane anesthesia-induced hyperglycemia (Zuurbier et al., 2008) (less than 5 min from start to exsanguination) changes plasma glucose levels, but its effects on RBC glucose (if any) would be at least partly buffered. In any case, it is highly improbable that these changes would affect differentially the rats depending on their diet. The uniformity of the data

obtained seem to support this weak point of our experimental setup. We have not been able to circumvent the problem within the ethical standards of our Laboratory.

The lower blood *vs.* plasma glucose levels, more marked in cafeteria diet-fed rats, attest directly to a lower cell compartment glucose content (there were minimal differences in packed cell volume). The accuracy of the calculations used to quantify the cell glucose content notwithstanding, do not change the fact that cafeteria rats had higher glucose content in the blood plasma fraction compared with that of cells; precisely the glucose in direct contact with hemoglobin.

Metabolic syndrome, diabetes and in general, high exposure to inflammation and hyperglycemia increase the glycosylation of plasma proteins (*Matsuura, Hughes & Khamashta, 2008; Roohk & Zaidi, 2008*). In fact, glycosylated albumin has been proposed as an indicator of maintained hyperglycemia (i.e., exposure of plasma proteins to higher aldose levels for long periods) (*Abe & Matsumoto, 2008*). However, the most used indicator of long-time maintained hyperglycemia is the measurement of glycosylated hemoglobin (*Siu & Yuen, 2014*), which initially was applied to whole RBC hemoglobin (*Carson et al., 2010*), but soon was focused on the terminal amino groups of hemoglobin (Hb_{1AC}) alone, giving rise to a much more sensitive (and extended) assay methodology (*Little et al., 2008; Weykamp et al., 2008*). The study of glycosylated hemoglobin (Hb_{1AC}) has become one of the standard elements for the evaluation of diabetes (metabolic syndrome) and, in general, sustained hyperglycemia (*Ong et al., 2010*). The critical point, however, is that all hemoglobin is contained within the RBC, and is not in direct contact with plasma glucose. This obvious circumstance would make *a priori* glycosylated albumin a more acceptable indicator of hyperglycemia. The conundrum of a marker of glycosylation not in direct contact with the parameter it measures has not been sufficiently explained so far. Nevertheless, its widespread use and clinical reliability are powerful reasons in favor of its continued use despite the largely unexplained nature of its origins.

In mammals, the direct permeability of the RBC membrane to glucose is low, if any (*Britton, 1964; Rich et al., 1967*), however, interchange of plasma and RBC glucose is active thanks to a facilitated-diffusion transport system (*Levine, Oxender & Stein, 1965*). The transport has been attributed, mainly to GLUT1 (*Graybill et al., 2006*), which function may be regulated by insulin, glucocorticoids and other factors (*Kahn & Flyer, 1990*). However, no differences in glucose transport through erythrocyte membranes were found between diabetic and euglycemic children (*Mortensen & Brahm, 1985*). There is, also, a high variability in the permeability of RBC membranes to glucose, due to species differences, individual factors and transporter modulation/saturation (*Khera et al., 2008*).

Compartmentation of blood glucose between plasma and cells may be an important regulatory factor by itself (*Palou et al., 1980*), since glucose carried by blood cells is rapidly interchanged with tissues (*Jacquez, 1984*). This is in overt contradiction with the slow velocity of glucose interchange of RBC when measured *in vitro* (*Sen & Widdas, 1962*). In addition, given the glycolytic nature of mammalian RBC, it can be expected that a sizeable part of the glucose entering the cell is rapidly glycolyzed to lactate, a process that is the only significant source of ATP for the cell. This inefficient mechanism converts blood in a

sizeable source of lactate, which implies that a variable part of glucose will be converted to hexoses-P on entering the cell, and thus (at least in part, when isomerized to ketose-P) lose its aldose-related glycosylating capacity.

The high proportions of Hb_{1AC} found under conditions of assumed sustained hyperglycemia (*Giuffrida et al., 2010*) contrast with the physical existence of barriers between hyperglycemic plasma and hemoglobin. We could not explain why Hb_{1AC} is so highly correlated with hyperglycemia, since total hemoglobin glycosylation does not reflect only hyperglycemia (*Adams et al., 2009; Chan et al., 2014*), which is in agreement with our data. In addition, diabetogenic conditions, such as those presented here and those found in the literature (*Koga et al., 2007; Miyashita et al., 2007; Wakabayashi, 2012*) do not show the expected changes in hemoglobin glycosylation. Our data on the lack of significant changes elicited by diet on plasma protein glycosylation do not agree with the common occurrence of increased glycosylated proteins in plasma of humans and rodents alike under already settled metabolic syndrome or its associated pathologies (*Gornik & Lauc, 2008*). The probable differences lie in the fact that in our model of initial stages of metabolic syndrome, the pathologic markers have not been yet developed fully, as we have previously found (*Ferrer-Lorente et al., 2005*). It must be also taken into account that metabolic syndrome-induced modifications on plasma proteins (*Marliss et al., 2006; Welle et al., 1992*) and RBC (*Cohen, Franco & Joiner, 2004; Manodori & Kuypers, 2002*) increase their cell turnover rates which compounds the problem and makes more difficult the comparisons unless the data maintain their homology.

The elevated degree of glycosylation found in RBC membranes, however, shows that even the small differences in basal glycemia found in our model are enough to already induce several-fold changes in the glycosylative activity of plasma glucose. Probably, other factors so far not identified, may help explain the increased glycosylation observed even at early stages of the development of metabolic syndrome. The relationship with high-energy (lipid) diet is clear, but the common assumption that these changes are a correlate of hyperglycemia remain unproven, and largely based only on indirect evidence.

In our experiment, the degree of glycosylation of hemoglobin was high, even under conditions in which practically no free glucose was found within the RBC. We only measured glycosyl residues, not those recombined by Amadori rearrangements, i.e., those able to liberate hydroxymethylfurfural. However, the ratio of up to 90 glycosyl residues per subunit of hemoglobin is close to the level of saturation of glycosylable sites. This was necessarily achieved in at most one month, a time shorter than the mean rat RBC half-life, 45–50 days (*Burwell, Brickley & Finch, 1953*), a value considerably decreased in rats with metabolic syndrome (*Kung, Tseng & Wang, 2009*). However, compared with albumin which median life span is close to 2 days (*Reed et al., 1988*), the differences can be better explained, since exposure of hemoglobin was 15-fold higher (30 days out of 45–50) than that of plasma proteins (2 days, assuming a behavior comparable to that of albumin). The shorter exposure was predictably more intense (as that of RBC membrane protein) because plasma proteins were in constant contact with plasma glucose.

The levels of glycosylation of plasma proteins and hemoglobin observed do not reflect the (limited) changes in plasma glucose, however, RBC membranes do. The results we obtained are puzzling; they agree with the known fact that exposure to hyperglycemia results in increased protein glycosylation, as shown by membrane proteins' differences, but not observed in plasma proteins; this may be due to their shorter half-lives and limited span of glucose level change.

On the other side, the low free glucose levels observed in cafeteria diet-fed rat RBC, agree with a slower rate of uptake (*Prats et al., 1987*) compared with plasma, but cannot directly explain how the overall glycosylation of hemoglobin was unaffected by one month of consumption of a hyper-energetic obesogenic diet.

CONCLUSIONS

We conclude that blood glucose compartmentation, as previously indicated, may play a role, in the regulation of plasma/blood versus tissue glucose transport/transfer, more important than usually assumed, but also, that glycosylation of blood proteins widely affects non-diabetic young experimental animals, both under standard or hyper-energetic diet conditions. This extensive glycosylation does not seem to be markedly affected by sex; and could be hardly justified from an assumedly sustained plasma hyperglycemia. More detailed—and comprehensive—analyses should be carried out to study the dynamics of glucose *in vivo* in the RBC to understand how so extensive protein glycosylation as that found here could take place, including an special emphasis on the hormonal regulation of RBC glucose transporters.

We have also found that RBC membrane glycosylation is a sensitive indicator of developing metabolic syndrome-related hyperglycemia, more sensitive than the general measurement of plasma or RBC protein glycosylation.

ACKNOWLEDGEMENTS

At the time this investigation took place, Christian Baron was an undergraduate student. Laia Oliva worked *pro bono*.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was done with the partial support of grants of the Plan Nacional de Investigación en Biomedicina (SAF2012-34895) and the Plan Nacional de Ciencia y Tecnología de los Alimentos (AGL-2011-23635) of the Government of Spain, and assistance from the CIBER OBN. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Plan Nacional de Investigación en Biomedicina: SAF2012-34895.
Plan Nacional de Ciencia y Tecnología de los Alimentos: AGL-2011-23635.
CIBER OBN.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Laia Oliva performed the experiments, prepared figures and/or tables, reviewed drafts of the paper.
- Cristian Baron performed the experiments.
- José-Antonio Fernández-López analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, reviewed drafts of the paper.
- Xavier Remesar analyzed the data, contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Marià Alemany conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All animal handling procedures were carried out in accordance with the norms of European, Spanish and Catalan Governments. The study was specifically approved (DMAH-5483) by the Animal Ethics Committee of the University of Barcelona.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1101#supplemental-information>.

REFERENCES

- Abe M, Matsumoto K. 2008.** Glycated hemoglobin or glycated albumin for assessment of glyemic control in hemodialysis patients with diabetes? *Nature Clinical Practice Nephrology* 4:482–483 DOI 10.1038/ncpneph0881.
- Adams RJ, Appleton SL, Hill CL, Wilson DH, Taylor AW, Chittleborough CR, Gill TK, Ruffin RE. 2009.** Independent association of HbA_{1c} and incident cardiovascular disease in people without diabetes. *Obesity* 17:559–563 DOI 10.1038/oby.2008.592.
- Arola L, Alemany M, Herrera E, Palou A, Remesar X. 1981.** Effect of ether, sodium pentobarbital and chloral hydrate anesthesia on rat plasma metabolite concentrations. *Revista Española de Fisiología* 37:379–385.

- Asahi K, Ichimori K, Nakazawa H, Izuhara Y, Inagi R, Watanabe T, Miyata T, Kurokawa K. 2000.** Nitric oxide inhibits the formation of advanced glycation end products. *Kidney International* **58**:1780–1787.
- Borg R, Kuenen J. 2009.** HbA_{1c} levels can be expressed as an estimated Average glucose (eAG). *Clinical Laboratory International* **33**:14–16.
- Britton HG. 1964.** Permeability of the human red cell to labelled glucose. *Journal of Physiology* **170**:1–20 DOI [10.1113/jphysiol.1964.sp007310](https://doi.org/10.1113/jphysiol.1964.sp007310).
- Bunn HF, Gabbay KH, Gallop PM. 1978.** The glycosylation of hemoglobin: relevance to diabetes mellitus. *Science* **200**:21–27 DOI [10.1126/science.635569](https://doi.org/10.1126/science.635569).
- Burwell EL, Brickley BA, Finch CA. 1953.** Erythrocyte life span in small animals. Comparison of two methods employing radioiron. *American Journal of Physiology* **172**:718–724.
- Carson AP, Reynolds K, Fonseca VA, Muntner P. 2010.** Comparison of A1C and fasting glucose criteria to diagnose diabetes among US adults. *Diabetes Care* **33**:95–97 DOI [10.2337/dc09-1227](https://doi.org/10.2337/dc09-1227).
- Castell-Auví A, Cedó L, Pallarès V, Blay M, Ardévol A, Pinent M. 2012.** The effects of a cafeteria diet on insulin production and clearance in rats. *British Journal of Nutrition* **108**:1155–1162 DOI [10.1017/S0007114511006623](https://doi.org/10.1017/S0007114511006623).
- Chan CL, McFann K, Newnes L, Nadeau KJ, Zeitler PS, Kelsey M. 2014.** Hemoglobin A_{1c} assay variations and implications for diabetes screening in obese youth. *Pediatric Diabetes* **15**:557–563 DOI [10.1111/pedi.12132](https://doi.org/10.1111/pedi.12132).
- Cohen RM, Franco RS, Joiner CH. 2004.** Is poor glycemic control associated with reduced red blood cell lifespan? *Diabetes Care* **27**:1013–1014 DOI [10.2337/diacare.27.4.1013](https://doi.org/10.2337/diacare.27.4.1013).
- Correa Pinto DA, Monteiro Seraphim P. 2012.** Cafeteria diet intake for fourteen weeks can cause obesity and insulin resistance in Wistar rats. *Revista de Nutrição* **25**:313–319 DOI [10.1590/S1415-52732012000300001](https://doi.org/10.1590/S1415-52732012000300001).
- Ferrer-Lorente R, Cabot C, Fernández-López JA, Alemany M. 2005.** Combined effects of oleoyl-estrone and a β_3 -adrenergic agonist (CL316,243) on lipid stores of diet-induced overweight male Wistar rats. *Life Sciences* **77**:2051–2058 DOI [10.1016/j.lfs.2005.04.008](https://doi.org/10.1016/j.lfs.2005.04.008).
- Gabbay KH, Sosenko JM, Banuchi GA, Mininsohn MJ, Flückiger R. 1979.** Glycosylated hemoglobins: increased glycosylation of Hemoglobin A in diabetic patients. *Diabetes* **28**:337–340 DOI [10.2337/diab.28.4.337](https://doi.org/10.2337/diab.28.4.337).
- Giuffrida FMA, Sallum CFC, Gabbay MAL, Gomes MB, Pires AC, Dib SA. 2010.** Relationship between glycated hemoglobin and metabolic syndrome of type 1 and type 2 diabetes. A factor analysis study. *Diabetes Care* **33**:e80–e80 DOI [10.2337/dc09-2280](https://doi.org/10.2337/dc09-2280).
- Gomori G. 1942.** A modification of the colorimetric phosphorus determination for use with the photoelectric colorimeter. *Journal of Laboratory and Clinical Medicine* **27**:955–960.
- Gornik O, Lauc G. 2008.** Glycosylation of serum proteins in inflammatory diseases. *Disease Markers* **25**:267–278 DOI [10.1155/2008/493289](https://doi.org/10.1155/2008/493289).
- Gagnoli G, Tanganelli I, Signorini AM, Tarli P, Paoli C. 1982.** Non-enzymatic glycosylation of serum proteins as an indicator of diabetic control. *Acta Diabetologica Latina* **19**:161–166 DOI [10.1007/BF02581152](https://doi.org/10.1007/BF02581152).
- Graybill C, Van Hoek AN, Desai D, Carruthers AM, Carruthers A. 2006.** Ultrastructure of human erythrocyte GLUT1. *Biochemistry* **45**:8096–8107 DOI [10.1021/bi060398x](https://doi.org/10.1021/bi060398x).
- Guo CX, Zeng XJ, Song JJ, Zhang M, Wang HX, Xu XW, Du FH, Chen BX. 2012.** A soluble receptor for advanced glycation end-products inhibits hypoxia/reoxygenation-induced apoptosis in rat cardiomyocytes via the mitochondrial pathway. *International Journal of Molecular Sciences* **13**:11923–11940 DOI [10.3390/ijms130911923](https://doi.org/10.3390/ijms130911923).

- Henning C, Smuda M, Girndt M, Ulrich C, Glomb MA. 2011. Molecular basis of Maillard amide-advanced glycation end product (AGE) formation *in vivo*. *Journal of Biological Chemistry* 286:44350–44356 DOI 10.1074/jbc.M111.282442.
- Higgins PJ, Garlick RL, Bunn HF. 1982. Glycosylated hemoglobin in human and animal red cells: role of glucose permeability. *Diabetes* 31:743–748 DOI 10.2337/diab.31.9.743.
- Itkonen HM, Mills IG. 2013. N-Linked glycosylation supports cross-talk between receptor tyrosine kinases and androgen receptor. *PLoS ONE* 8:e65016 DOI 10.1371/journal.pone.0065016.
- Jacquez JA. 1984. Red blood cell as glucose carrier: significance for placental and cerebral glucose transfer. *American Journal of Physiology* 246:R289–R298.
- John WG, Lamb EJ. 1993. The Maillard or browning reaction in diabetes. *Eye* 7:230–237 DOI 10.1038/eye.1993.55.
- Kahn BB, Flyer JS. 1990. Regulation of glucose-transporter gene expression *in vitro* and *in vivo*. *Diabetes Care* 13:548–564 DOI 10.2337/diacare.13.6.548.
- Khera PK, Joiner CH, Carruthers A, Lindsell CJ, Smith EP, Franco RS, Holmes YR, Cohen RM. 2008. Evidence for interindividual heterogeneity in the glucose gradient across the human red blood cell membrane and its relationship to hemoglobin glycation. *Diabetes* 57:2445–2452 DOI 10.2337/db07-1820.
- Koga M, Otsuki M, Matsumoto S, Saito H, Mukai M, Kasayama S. 2007. Negative association of obesity and its related chronic inflammation with serum glycated albumin but not glycated hemoglobin levels. *Clinica Chimica Acta* 378:48–52 DOI 10.1016/j.cca.2006.10.013.
- Kung CM, Tseng ZL, Wang HL. 2009. Erythrocyte fragility increases with level of glycosylated hemoglobin in type 2 diabetic patients. *Clinical Hemorheology and Microcirculation* 43:345–351.
- Levine M, Oxender DL, Stein W. 1965. The substrate-facilitated transport of the glucose carrier across the human erythrocyte membrane. *Biochimica et Biophysica Acta* 109:151–163 DOI 10.1016/0926-6585(65)90099-3.
- Little RR, Rohlfing CL, Hanson S, Connolly S, Higgins T, Weykamp CW, D'Costa M, Luzzi V, Owen WE, Roberts WL. 2008. Effects of hemoglobin (Hb) E and HbD traits on measurements of glycated Hb (HbA_{1C}) by 23 methods. *Clinical Chemistry* 54:1277–1282 DOI 10.1373/clinchem.2008.103580.
- Lowry OH, Rosebrough RW, Farr AL, Randall RJ. 1951. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry* 193:265–275.
- Manodori AB, Kuypers FA. 2002. Altered red cell turnover in diabetic mice. *Journal of Laboratory and Clinical Medicine* 140:161–165 DOI 10.1067/mlc.2002.126504.
- Marliss E, Chevalier S, Gougeon R, Morais J, Lamarche M, Adegoke O, Wu G. 2006. Elevations of plasma methylarginines in obesity and ageing are related to insulin sensitivity and rates of protein turnover. *Diabetologia* 49:351–359 DOI 10.1007/s00125-005-0066-6.
- Matsuura E, Hughes GR, Khamashta MA. 2008. Oxidation of LDL and its clinical implication. *Autoimmunity Reviews* 7:558–566 DOI 10.1016/j.autrev.2008.04.018.
- Miller JA, Gravallesse E, Bunn HF. 1980. Nonenzymatic glycosylation of erythrocyte membrane proteins. Relevance to diabetes. *Journal of Clinical Investigation* 65:896–901 DOI 10.1172/JCI109743.
- Miwa I, Maeda K, Okuda J, Okuda G. 1972. Mutarotase effect on colorimetric determination of blood-glucose with β -D-glucose oxidase. *Clinica Chimica Acta* 37:538–340 DOI 10.1016/0009-8981(72)90483-4.

- Miyashita Y, Nishimura R, Morimoto A, Matsudaira T, Sano H, Tajima N. 2007. Glycated albumin is low in obese, type 2 diabetic patients. *Diabetes Research and Clinical Practice* 78:51–55 DOI 10.1016/j.diabres.2007.02.021.
- Mortensen HB, Brahm J. 1985. Glucose transport in erythrocytes of diabetic and healthy children as related to hemoglobin A_{1c}. *Clinical Chemistry* 31:1387–1389.
- Nathan DM, Kuenen J, Borg R, Zheng H, Schoenfeld D, Heine RJ. 2008. Translating the A1C Assay into estimated average glucose values. *Diabetes Care* 31:1473–1478 DOI 10.2337/dc08-0545.
- Ong KL, Tso AWK, Lam KSL, Cherny SS, Sham PC, Cheung BMY. 2010. Using glycosylated hemoglobin to define the metabolic syndrome in United States adults. *Diabetes Care* 33:1856–1858 DOI 10.2337/dc10-0190.
- Palou A, Remesar X, Arola L, Alemany M. 1980. Blood and plasma glucose relationships during pregnancy, the breeding cycle and development in the rat. *Diabete et Metabolisme* 6:271–275.
- Poulsen MW, Bak MJ, Andersen JM, Monosik R, Giraudi-Futin AC, Holst JJ, Nielsen J, Lauritzen L, Larsen LH, Bugel S, Dragsted LO. 2014. Effect of dietary advanced glycation end products on postprandial appetite, inflammation, and endothelial activation in healthy overweight individuals. *European Journal of Nutrition* 53:661–672 DOI 10.1007/s00394-013-0574-y.
- Prats E, Monfar M, Argilés JM, Alemany M. 1987. Effects of starvation and a high-energy diet on rat blood compartmentation of injected radioactive alanine and glucose. *Biochemistry International* 14:95–101.
- Prats E, Monfar M, Iglesias R, Castellà J, Alemany M. 1989. Energy intake of rats fed a cafeteria diet. *Physiology and Behavior* 45:263–272 DOI 10.1016/0031-9384(89)90128-5.
- Reed RG, Davidson L, Burrington C, Peters T. 1988. Non-resolving jaundice: bilirubin covalently attached to serum albumin circulates with the same metabolic half-life as albumin. *Clinical Chemistry* 34:1992–1994.
- Rich G, Sha'afi R, Barton T, Solomon A. 1967. Permeability studies on red cell membranes of dog, cat, and beef. *The Journal of General Physiology* 50:2391–2405 DOI 10.1085/jgp.50.10.2391.
- Romero MM, Fernández-López JA, Alemany M, Esteve M. 2010. Gene expression modulation of liver energy metabolism by oleoyl-estrone in overweight rats. *Bioscience Reports* 20:81–89.
- Romero MM, Fernández-López JA, Remesar X, Alemany M. 2012. Effect of sex and prior exposure to a cafeteria diet on the distribution of sex hormones between plasma and blood cells. *PLoS ONE* 7(3):e34381 DOI 10.1371/journal.pone.0034381.
- Roohk HV, Zaidi AR. 2008. A review of glycated albumin as an intermediate glycation index for controlling diabetes. *Journal of Diabetes Science and Technology* 2:1114–1121 DOI 10.1177/193229680800200620.
- Sclafani A, Springer D. 1976. Dietary obesity in adult rats: similarities to hypothalamic and human obesity syndromes. *Physiology and Behavior* 17:461–471 DOI 10.1016/0031-9384(76)90109-8.
- Sen A, Widdas W. 1962. Determination of the temperature and pH dependence of glucose transfer across the human erythrocyte membrane measured by glucose exit. *The Journal of physiology* 160:392–403 DOI 10.1113/jphysiol.1962.sp006854.
- Siu PM, Yuen QS. 2014. Supplementary use of HbA_{1c} as hyperglycemic criterion to detect metabolic syndrome. *Diabetology and Metabolic Syndrome* 6:Article 119 DOI 10.1186/1758-5996-6-119.

- Stein J, Smith G. 1982.** Phospholipids. In: *Techniques in lipid and membrane biochemistry*. County Clare: Elsevier/North Holland, 1–15.
- Thomas TN, Rhodin JA, Clark L, Garces A, Bryant M. 2003.** A comparison of the anti-inflammatory activities of conjugated estrogens and 17- β estradiol. *Inflammation Research* **52**:452–460 DOI [10.1007/s00011-003-1198-0](https://doi.org/10.1007/s00011-003-1198-0).
- Tomoda A, Kodaira K, Taketo A, Tanimoto K, Yoneyama Y. 1984.** Isolation of human erythrocyte membranes in glucose solution. *Analytical Biochemistry* **140**:386–390 DOI [10.1016/0003-2697\(84\)90182-9](https://doi.org/10.1016/0003-2697(84)90182-9).
- Wakabayashi I. 2012.** Obesity-independent inverse association between regular alcohol consumption and hemoglobin A_{1C}. *Obesity Facts* **5**:60–67 DOI [10.1159/000336067](https://doi.org/10.1159/000336067).
- Welle S, Barnard R, Statt M, Amatruda J. 1992.** Increased protein turnover in obese women. *Metabolism* **41**:1028–1034 DOI [10.1016/0026-0495\(92\)90133-U](https://doi.org/10.1016/0026-0495(92)90133-U).
- Weykamp C, John WG, Mosca A, Hoshino T, Little R, Jeppsson J-O, Goodall I, Miedema K, Myers G, Reinauer H. 2008.** The IFCC reference measurement system for HbA_{1c}: a 6-year progress report. *Clinical Chemistry* **54**:240–248 DOI [10.1373/clinchem.2007.097402](https://doi.org/10.1373/clinchem.2007.097402).
- Wu CH, Huang HW, Huang SM, Lin JA, Yeh CT, Yen GC. 2011.** AGE-induced interference of glucose uptake and transport as a possible cause of insulin resistance in adipocytes. *Journal of Agricultural and Food Chemistry* **59**:7978–7984 DOI [10.1021/jf201271y](https://doi.org/10.1021/jf201271y).
- Zuurbier CJ, Keijzers PJ, Koeman A, Van Wezel HB, Hollmann MW. 2008.** Anesthesia's effects on plasma glucose and insulin and cardiac hexokinase at similar hemodynamics and without major surgical stress in fed rats. *Anesthesia and Analgesia* **106**:135–142 DOI [10.1213/01.ane.0000297299.91527.74](https://doi.org/10.1213/01.ane.0000297299.91527.74).