

CORRECTION


Correction: EBV induces persistent NF- κ B activation and contributes to survival of EBV-positive neoplastic T- or NK-cells

Honami Takada, Ken-Ichi Imadome, Haruna Shibayama, Mayumi Yoshimori, Ludan Wang, Yasunori Saitoh, Shin Uota, Shoji Yamaoka, Takatoshi Koyama, Norio Shimizu, Kouhei Yamamoto, Shigeyoshi Fujiwara, Osamu Miura, Ayako Arai

There is an error in affiliation 4 for Yasunori Saitoh, Shin Uota, and Shoji Yamaoka. Affiliation 4 should be: Department of Molecular Virology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University (TMDU), Bunkyo-ku, Tokyo, Japan.

Reference

1. Takada H, Imadome K-I, Shibayama H, Yoshimori M, Wang L, Saitoh Y, et al. (2017) EBV induces persistent NF- κ B activation and contributes to survival of EBV-positive neoplastic T- or NK-cells. PLoS ONE 12(3): e0174136. <https://doi.org/10.1371/journal.pone.0174136> PMID: [28346502](https://pubmed.ncbi.nlm.nih.gov/28346502/)



OPEN ACCESS

Citation: Takada H, Imadome K-I, Shibayama H, Yoshimori M, Wang L, Saitoh Y, et al. (2017) Correction: EBV induces persistent NF- κ B activation and contributes to survival of EBV-positive neoplastic T- or NK-cells. PLoS ONE 12(8): e0182682. <https://doi.org/10.1371/journal.pone.0182682>

Published: August 1, 2017

Copyright: © 2017 Takada et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.