

# Asbestos and multi-walled carbon nanotubes generate distinct oxidative responses in inflammatory cells

Satomi Funahashi,<sup>1</sup> Yasumasa Okazaki,<sup>1</sup> Daiki Ito,<sup>1</sup> Atsushi Asakawa,<sup>2</sup> Hiroataka Nagai,<sup>1</sup> Masafumi Tajima<sup>2</sup> and Shinya Toyokuni<sup>1,\*</sup>

<sup>1</sup>Department of Pathology and Biological Responses, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan

<sup>2</sup>ATTO Corporation, 3-2-2 Motoasakusa, Taito-ku, Tokyo 111-0041, Japan

(Received 6 August, 2014; Accepted 26 August, 2014; Published online 28 November, 2014)

Asbestos exposure is considered a social burden by causing mesothelioma. Despite the use of synthetic materials, multi-walled carbon nanotubes (MWCNTs) are similar in dimension to asbestos and produce mesothelioma in animals. The role of inflammatory cells in mesothelial carcinogenesis remains unclear. Here, we evaluated the differences in inflammatory cell responses following exposure to these fibrous materials using a luminometer and L-012 (8-amino-5-chloro-7-phenylpyrido[3,4-*d*]pyridazine-1,4-(2*H*,3*H*)dione) to detect reactive oxygen species (ROS). Rat peripheral blood or RAW264.7 cells were used to assess the effects on neutrophils and macrophages, respectively. Crocidolite and amosite induced significant ROS generation by neutrophils with a peak at 10 min, whereas that of chrysotile was ~25% of the crocidolite/amosite response. MWCNTs with different diameters (~15, 50, 115 and 145 nm) and different carcinogenicity did not induce significant ROS in peripheral blood. However, the MWCNTs induced a comparable amount of ROS in RAW264.7 cells to that following asbestos treatment. The peaks for MWCNTs (0.5–1.5 h) were observed earlier than those for asbestos (1–5 h). Apocynin and superoxide dismutase significantly inhibited ROS generation for each fiber, suggesting an involvement of NADPH oxidase and superoxide. Thus, asbestos and MWCNTs induce different oxidative responses in inflammatory cells, indicating the importance of mesothelial cell evaluation for carcinogenesis.

**Key Words:** asbestos, multi-walled carbon nanotubes, neutrophil, macrophage, NADPH oxidase

Novel materials may generate an unexpected health risk to the human society. Asbestos, a natural fibrous mineral, is an example. Asbestos presents excellent durability, such as resistance to acid, heat and friction, versatility for mixing with other materials and huge economic merits depending on mining, and because of these beneficial properties, it was abundantly used in industries worldwide during the last century. Therefore, it was copiously released into the air surrounding mines and factories. It took a few decades of epidemiological studies to declare that asbestos is the cause of not only asbestosis but also a rare and aggressive malignant tumor, mesothelioma.<sup>(1–3)</sup> Due to mesothelioma's extremely long incubation period of 30–40 years, the peak incidence of mesothelioma in Japan is predicted to occur in 2025 with 100,000 new patients being diagnosed in the coming 40 years.<sup>(4)</sup>

Multi-walled carbon nanotubes (MWCNTs) are novel synthetic materials consisting only of carbon,<sup>(5)</sup> and due to its superior properties, are already used in industries for production of semi-conductors, fuel cells and structural materials. However, the physical dimensions and the biopersistence of MWCNTs were

found to be similar to asbestos, and they indeed have revealed asbestos-like pathogenicity,<sup>(6,7)</sup> including mesothelial carcinogenesis in rodents.<sup>(8–11)</sup> In both cases, mesothelial cells, which are closely associated with foreign body-induced inflammation and the associated local iron overload, are the predominant carcinogenic cells.<sup>(10,12–15)</sup> Previous studies have reported the generation of reactive oxygen species (ROS) with luminol by macrophages or isolated neutrophils exposed to asbestos,<sup>(16–22)</sup> and the toxicity of MWCNTs to macrophages.<sup>(23)</sup> However, there are limited data available on the oxidative responses of inflammatory cells to MWCNTs.

Our recent studies revealed that MWCNT diameter and rigidity are critical factors in mesothelial injury and carcinogenesis.<sup>(10)</sup> Here, we compared neutrophil and macrophage responses to asbestos and MWCNTs of various defined diameters by measuring ROS generation, and thus studied the contribution of inflammation in fiber-induced mesothelial carcinogenesis. There has been a recent progress in the luminometer and its probe as well as protocols, thus allowing peripheral blood to be used as the source of neutrophils without performing separation procedures. In the present study, we found that asbestos and MWCNTs generate distinct responses in the inflammatory cells.

## Materials and Methods

**Materials.** We obtained asbestos (crocidolite, amosite and chrysotile) from the Union for International Cancer Control (UICC; Geneva, Switzerland) and suspended it in 0.9% saline (5 mg/ml). We obtained MWCNTs and suspended them in 0.5% bovine serum albumin (BSA) (5 mg/ml in saline) as described.<sup>(10,11)</sup> The carbon nanotubes were distinguished as CNT-50, CNT-115, CNT-145 and CNT-tngl to represent the diameter of each nanotube (Table 1). Zymosan (Sigma-Aldrich, St. Louis, MO) and lipopolysaccharide (LPS) from *E. coli* 0111:B4 (Sigma-Aldrich) were used to initiate inflammation. Deferoxamine mesylate (DFO) from Sigma-Aldrich and nitrilotriacetate disodium salt from Nakalai Tesque (Kyoto, Japan) were used at a final concentration of 40  $\mu$ M.

**Antioxidants.** Cu,Zn superoxide dismutase from bovine erythrocytes (SOD1; EC1.15.1.1), catalase from bovine liver (EC1.11.1.6) and apocynin (APO; a NADPH oxidase inhibitor) were purchased from Sigma-Aldrich and were used at final concentrations of 1,000 units/L, 1,000 units/L and 1 mM, respectively. NaN<sub>3</sub> (sodium azide; an inhibitor of catalase, peroxidase

\*To whom correspondence should be addressed.  
E-mail: toyokuni@med.nagoya-u.ac.jp

**Table 1.** Characteristics of asbestos and MWCNTs

	Fibers	Structural formula	Diameter (nm)	Length ( $\mu\text{m}$ )	*Mesothelial carcinogenicity (50% incidence, days)
Asbestos	Crocidolite	$\text{Na}_2(\text{Fe}^{2+})_3(\text{Fe}^{3+})_2\text{Si}_8\text{O}_{22}(\text{OH})_2$	40–150	4.54	~600
	Amosite	$(\text{Fe-Mg})_7\text{Si}_8\text{O}_{22}(\text{OH})_2$	60–350	5.45	~600
	Chrysotile	$\text{Mg}_3(\text{Si}_2\text{O}_5)(\text{OH})_4$	20–80	3.87	~400
MWCNTs	CNT-50	$\text{C}_n$	**52.40 $\pm$ 0.72	**4.60 $\pm$ 0.10	~280
	CNT-115	$\text{C}_n$	116.25 $\pm$ 1.58	4.88 $\pm$ 0.10	Not determined
	CNT-145	$\text{C}_n$	143.5 $\pm$ 1.56	4.34 $\pm$ 0.08	~320***
	CNT-tngl	$\text{C}_n$	~15	Not applicable	No carcinogenicity <sup>(36)</sup>

\*Data are based on 10 mg intraperitoneal injection to F1 rats between Fischer-344 and Brown-Norway. \*\*Means  $\pm$  SEM.<sup>(10,14)</sup> \*\*\*Low carcinogenicity (17%) at the dose of 1 mg intraperitoneal injection at day 350.<sup>(10)</sup> MWCNTs, multi-walled carbon nanotubes.

and cytochrome oxidase) was from Wako and was used at a final concentration of 1 mM.

**Peripheral blood and macrophage cell line.** Male Sprague-Dawley rats 15 weeks old (Shizuoka Laboratory Center, Hamamatsu, Japan) were used ( $n = 3$  for each group). The animals were anesthetized with pentobarbital, and the blood was collected from the inferior vena cava with heparinization immediately before use. The animal experiment committee of Nagoya University Graduate School of Medicine approved this experiment. We used the murine macrophage cell line RAW264.7 (DS Pharma Biomedical, Osaka, Japan).

**Determination of ROS generated from inflammatory cells.** We measured ROS with a luminometer (AB-2280; Atto Corporation, Tokyo, Japan; detection range, 350–900 nm) using L-012 (8-amino-5-chloro-7-phenylpyrido[3,4-*d*]pyridazine-1,4-(2*H*,3*H*) dione sodium salt; Wako Pure Chem. Co., Ltd., Osaka, Japan) as a chemiluminescent probe. L-012 develops strong chemiluminescence with a  $\lambda_{\text{max}}$  of 458 nm when it reacts with ROS, including superoxide ( $\text{O}_2^{\cdot-}$ ), hypochlorite ( $\text{HClO}^-$ ) and hydroxyl radical ( $\cdot\text{OH}$ ), among which  $\cdot\text{OH}$  causes the highest chemiluminescence.<sup>(24)</sup>

In the peripheral blood experiments, blood (20  $\mu\text{l}$ ) and glucose (5  $\mu\text{l}$ ; final concentration 10 mM) were incubated at 37°C for 3 min. L-012 (20  $\mu\text{l}$ ; final concentration 2  $\mu\text{M}$ ), a material sample (final concentration 1.0, 2.0 or 4.0 mg/ml) and an antioxidant (10  $\mu\text{l}$ ) were combined and adjusted to a total volume of 250  $\mu\text{l}$  with 10 mM phosphate-buffered saline (PBS; pH 7.4). After ample pipetting and vortexing, we started each measurement. Measurements were performed on the luminometer for 10 s and were repeated 99 times every 30 s for a period of ~50 min. Zymosan (1 mg/ml) was used as a positive control, and 0.9% NaCl and 0.5% BSA in saline were used as a negative controls for asbestos and MWCNTs, respectively.

For the macrophage experiments, RAW264.7 cells ( $1 \times 10^6$ ) were incubated in a 6-well plate in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum (FBS) and antibiotic/antimycotic (Life Technologies, Carlsbad, CA) at 37°C in 5%  $\text{CO}_2$  for 24 h. Asbestos (5  $\mu\text{g}/\text{cm}^2$ ) or MWCNTs (5  $\mu\text{g}/\text{cm}^2$ ) were then added to the culture, and the cells were further incubated for up to 7 h. The cells were removed with a scraper and recovered by centrifuge at 720  $\times g$ . New medium with FBS (230  $\mu\text{l}$ ) was then added, followed by incubation at 37°C for 3 min. Then, L-012 (20  $\mu\text{l}$ ; final concentration 2  $\mu\text{M}$ ) was added for a total volume of 250  $\mu\text{l}$ , and the measurements were performed as described above. LPS (1.2  $\mu\text{g}/\text{ml}$ ) was used as a positive control for macrophage stimulation, and 0.9% NaCl and 0.5% BSA in saline were used as negative controls for asbestos and MWCNTs, respectively.

**Hemolysis.** Heparinized blood (100  $\mu\text{l}$ ) and a fibrous material (5 mg/ml; in PBS or 0.5% BSA in PBS) were mixed and incubated at 37°C for 4 h. Thereafter, samples were centrifuged at 1,500  $\times g$  for 5 min, and the collected supernatant was measured

for absorbance at 540 nm (hemoglobin) using a spectrophotometer (ND-2000, Thermo, Japan). The hemolysis percentage (HP) was calculated using the following equation as described previously:<sup>(25)</sup>

$$\text{HP (\%)} = (\text{D}_t - \text{D}_{\text{nc}}) / (\text{D}_{\text{pc}} - \text{D}_{\text{nc}}) \times 100$$

$\text{D}_t$  is the absorbance of the test samples;  $\text{D}_{\text{pc}}$  and  $\text{D}_{\text{nc}}$  are the absorbances of the positive and negative control, respectively. The results are shown as the average of three independent measurements.

**Time-lapse microscopic observation.** BZ-9000 (Keyence, Osaka, Japan) was used for time-lapse video microscopy of RAW264.7 cells up to 5 h.

**Statistics.** The peak values of chemiluminescence during the observation period were analyzed by one-way ANOVA with Dennett's multiple comparison test through Prism5 (GraphPad Software Inc., San Diego, CA). Means  $\pm$  SEM are shown.

## Results

### ROS from neutrophils increased with asbestos treatment but not with MWCNT treatment.


Using zymosan as a positive control, we confirmed that the whole system works well (average  $\text{RLU} = 40.6 \times 10^3$ ). The peak time (~20 min) after addition (data not shown) also confirmed that we observe the function of neutrophils in the peripheral blood. All asbestos treatments significantly increased ROS generation in a dose-dependent manner with a peak increase at ~10 min (Fig. 1A–C). ROS generation by crocidolite and amosite were significantly higher than that of chrysotile (~25% of crocidolite/amosite;  $\text{RLU} < 1,000$ ). Amosite induced the highest ROS generation, followed by crocidolite and chrysotile (amosite > crocidolite >>> chrysotile). In contrast, MWCNTs of all diameters (Table 1) did not induce significant ROS generation under the same experimental conditions (Fig. 1D and data not shown).

SOD1, catalase (crocidolite only), sodium azide and apocynin significantly inhibited the ROS generation induced by crocidolite and amosite, indicating the involvement of  $\text{O}_2^{\cdot-}$ ,  $\text{H}_2\text{O}_2$ , cytochrome oxidase and NADPH oxidase. DFO also inhibited ROS generation, whereas nitrotriacetate promoted it. Inhibitory experiments were not performed for chrysotile due to its relatively low ROS generation.

### ROS from macrophages increased with both asbestos and MWCNT treatment.

LPS-mediated ROS increased in a concentration-dependent manner (0.12 < 1.2 < 12  $\mu\text{g}/\text{ml}$ ; data not shown), revealing that the system worked. Stimulation with 1.2  $\mu\text{g}/\text{ml}$  LPS caused a continual ROS generation for more than 24 h in RAW264.7 cells (Fig. 2A and data not shown). Furthermore, the peak time was different for each type of asbestos: crocidolite was 3 h, amosite was 2 h and chrysotile was >5 h (Fig. 2A–C).

Similar to asbestos, MWCNTs of various diameters consistently induced ROS generation in the macrophage cells (Fig. 2D–F). The peak time for MWCNTs occurred much earlier than that



**Fig. 1.** ROS generation by neutrophils was different following asbestos and MWCNT exposure. A luminometer and L-012 were used to measure ROS generation from neutrophils (rat whole blood) after stimulation by each type of fiber. (A) crocidolite; (B) amosite; (C) chrysotile; (D) CNT-50 (CNT-115, CNT-145 and CNT-tngl showed similar results). The inset in A and B shows the results of the inhibition experiments using antioxidants and iron modulators. Refer to the text for further details. *P* values were determined by one-way ANOVA with Dennett's multiple comparison test ( $*p < 0.05$ ,  $**p < 0.01$ ,  $***p < 0.001$ , ns, not significant vs each fiber;  $##p < 0.01$ ;  $###p < 0.001$  vs control; means  $\pm$  SEM from at least three independent samples). APO, apocynin; CAT, catalase; CNT, carbon nanotube; Cont, control; DFO, deferoxamine mesylate; MWCNT, multi-walled carbon nanotube; NTA, nitritoltriacetate; RLU, relative luminescence unit; ROS, reactive oxygen species; SOD, superoxide dismutase.

observed for asbestos: CNT-50 and CT-115 occurred at 1.5 h, CNT-145 at 0.5 h and CNT-tngl at 1 h (Fig. 2D–F).


SOD1 and apocynin consistently inhibited ROS generation, but catalase did not work in all experiments.  $\text{NaN}_3$  inhibited the ROS generated by CNT-115, CNT-145 and CNT-tngl. NTA significantly promoted ROS generation only with CNT-50 treatment, whereas DFO inhibited it for crocidolite, CNT-115 and CNT-tngl. Of note, DFO promoted ROS generation only with chrysotile (Fig. 2C and F).

**Hemolysis was induced by chrysotile.** Among the asbestos and MWCNTs used, only chrysotile caused massive hemolysis, which was 75% after a 4 h incubation.


**Distinct motion of macrophages after exposure to asbestos and MWCNTs.** We observed RAW264.7 cells using time-lapse microscopy analysis after exposure to either asbestos or MWCNTs. In the case of asbestos, we observed cell movements toward the fibers, leading to the isolation of fibers from media by groups of cells (Fig. 3A and B). In contrast, the cells remained dispersed with any MWCNTs (Fig. 3C and D).

## Discussion

The biological assessment of novel synthetic materials is important to evaluate human health risk. We compared the inflammatory response in neutrophils and macrophages following expo-



**Fig. 2.** ROS generation by macrophages after exposure to asbestos and MWCNTs. A luminometer and L-012 were used to measure ROS generation from macrophages (RAW264.7) after incubation with each fiber. Refer to the text for further details. (A–C) asbestos; (D–F) MWCNTs. LPS, lipopolysaccharide. Refer to the legend of Fig. 1 for abbreviations.



**Fig. 3.** Time-lapse microscopic analysis of cellular movements by macrophages after exposure to each fiber. Cellular movements were distinct between asbestos and MWCNT exposures. (A) crocidolite; (B) chrysotile; (C) CNT-50; (D) CNT-145. The number indicates h and min. Differences between asbestos and MWCNTs are indicated by the areas with disrupted circles (cellular aggregation). Refer to the text for further details (bar = 80  $\mu$ m).


sure to asbestos and MWCNTs with various diameters *ex vivo*. There was no difference observed in the chemiluminescence emitted by whole blood and with neutrophil isolation with L-012 probe.<sup>(24)</sup> Furthermore, the present *ex vivo* system worked well as demonstrated by the use of positive controls (zymosan and LPS). We found, for the first time to our knowledge, that ROS generation in neutrophils was completely different between asbestos and MWCNTs exposure (Fig. 1). Of note, we did not observe neutrophil stimulation by any of the MWCNTs used, indicating that the response was independent of the diameter (Fig. 1D). We believe that this effect is associated with the formulation of MWCNTs. MWCNTs consist only of carbon,<sup>(5,15)</sup> an element in the backbones of most biomolecules, whereas asbestos is a fibrous crystal made of silicon, oxygen and minerals.<sup>(1)</sup> The results indicate that the acute neutrophilic inflammation following exposure to MWCNTs may be minimal compared to other similar fibrous materials, which may call for medical attention.

Indeed, neutrophils reacted to all the types of asbestos tested within 10 min. The ROS generation was much higher with amosite and crocidolite treatment than with chrysotile. This finding is consistent with the direct catalytic activity of each type of asbestos for Fenton reactions observed by electron spin resonance analysis.<sup>(26)</sup> There are two indications on the results: amosite and crocidolite were found to contain large amounts of iron (27.3% and 28.5%, respectively), and chrysotile caused massive hemolysis. The presence of surface iron may facilitate ROS generation, and conversely, hemoglobin and heme in the reaction

mixture may delay or inhibit ROS generation by their toxicity.<sup>(27–29)</sup> The ROS generated were  $O_2^{\cdot-}$ ,  $H_2O_2$ , and  $\cdot OH$ , and based on the inhibition experiments, their generation was associated with cytochrome oxidase and NADPH oxidase. It is known that DFO blocks catalytic iron and nitrilotriacetate promotes it.<sup>(30–32)</sup> The results indicate that catalytic iron is also involved in the ROS generation from neutrophils.

Macrophages are the second cells following neutrophils to arrive at the site of inflammation and play a major role in chronic inflammation when the inflammatory stimulus is not quickly eliminated.<sup>(33)</sup> Both asbestos and MWCNTs induced ROS generation in RAW264.7 cells. Following asbestos exposure, the peak time observed in macrophages was later than that observed for neutrophils. Furthermore, the peak time occurred earlier following MWCNTs exposure (0.5–1.5 h) compared to asbestos (2–5 h) (Fig. 2A, B, D and E). This result suggests that different mechanisms exist for sensing the presence of different fibrous materials that have similar dimensions. Using video microscopy, we observed that the isolation activity for fibrous materials by macrophages, indicated by cell gathering, is stronger for asbestos than for MWCNTs (Fig. 3).

At the same time, the results observed for macrophages treated with antioxidants and iron chelators were much different from the results observed for neutrophils and also for each fiber. NADPH oxidase and  $O_2^{\cdot-}$  were consistently involved as indicated by the effects observed with apocynin and SOD treatment. This is consistent with recent reports on the involvement of NLRP3



**Fig. 4.** Summary of the differences in oxidative response of inflammatory cells to each fiber.

inflammasome activation by nanomaterials.<sup>(34,35)</sup> Catalase did not work for any of the fibers, suggesting that H<sub>2</sub>O<sub>2</sub> is not involved or was present in an unapproachable fashion. DFO worked for crocidolite and CNT-115, and NTA promoted ROS generation only in CNT-50 (Fig. 2C and F). We recently reported different mechanisms for the uptake of asbestos and MWCNTs by mesothelial cells. The former was phagocytosis, whereas the latter was penetration/piercing, which was most prominent for CNT-50.<sup>(10)</sup> The effect of NTA may be associated with the intracellular localization of fibers in macrophages, which may induce catalytic iron in the cell. The enhancement in ROS generation in chrysotile-exposed macrophages with DFO was unexpected (Fig. 2C). Among the types of asbestos used in the study, chrysotile displays the highest toxicity and inflammogenicity.<sup>(10)</sup> Thus, iron removal by DFO may have stimulated the macrophages. The time required for macrophage activation was different between asbestos and MWCNTs (Fig. 2A and D). The identification of molecules on the outer membrane and inside of neutrophils and macrophages that are associated with the present results is an interesting issue to pursue in the near future.

In our previous experiments of mesothelial carcinogenesis in rats, the MWCNTs used in the present study revealed that CNT-50 is potentially carcinogenic,<sup>(10)</sup> CNT-145 is less carcinogenic and CNT-tngl was not carcinogenic (Table 1).<sup>(36)</sup> However, in the studies presented here, neutrophil responses were subtle and macrophage responses were basically the same among the MWCNTs used (Fig. 4). Therefore, we believe that mesothelial cell injury is a more important indicator than the responses induced by neutrophils and macrophages in fiber-induced mesothelial carcinogenesis. On the other hand, neutrophils and macrophages may play a role in the inflammatory conditions often observed after airway exposure, such as bronchitis, pneumonitis and pulmonary fibrosis. Whether MWCNTs cause different types of pulmonary disease from those caused by asbestos should be carefully monitored in workers involved in MWCNT production.

## References

- Roggli VL, Oury TD, Sporn TA. *Pathology of Asbestos-Associated Diseases*. 2nd ed. New York: Springer Verlag, 2004.
- IARC, WHO. Asbestos (chrysotile, amosite, crocidolite, tremolite, actinolite, and anthophyllite). IARC Monographs on the Evaluation of Carcinogenic

## Acknowledgments

We would like to thank Nobuyuki Misawa for excellent technical assistance. This work was supported in part by a grant-in-aid for research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan (24390094; 221S0001-04; 24108001).

## Abbreviations

Amo	amosite (brown asbestos)
ANOVA	analysis of variance
APO	apocynin
BSA	bovine serum albumin
CAT	catalase
Chry	chrysotile (white asbestos)
CNT	carbon nanotube
Cro	crocidolite (blue asbestos)
DFO	deferoxamine mesylate
FBS	fetal bovine serum
L-012	8-amino-5-chloro-7-phenylpyrido[3,4- <i>d</i> ]pyridazine-1,4-(2 <i>H</i> ,3 <i>H</i> ) dione
LPS	lipopolysaccharide
MWCNT(s)	multi-walled carbon nanotube(s)
PBS	phosphate-buffered saline
RLU	relative luminescence unit
ROS	reactive oxygen species
SEM	standard error of means
SOD	superoxide dismutase

## Conflict of Interest

No potential conflicts of interest were disclosed.

Risks to Humans. A Review of Human Carcinogens; Part C: Arsenic, Metals, Fibres, and Dusts. France: Lyon, 2012; 219–309.

- Aierken D, Okazaki Y, Chew SH, *et al.* Rat model demonstrates a high risk of tremolite but a low risk of anthophyllite for mesothelial carcinogenesis.

- Nagoya J Med Sci* 2014; **76**: 149–160.
- 4 Robinson BW, Lake RA. Advances in malignant mesothelioma. *N Engl J Med* 2005; **353**: 1591–1603.
  - 5 Iijima S. Helical Microtubules of Graphitic Carbon. *Nature* 1991; **354**: 56–58.
  - 6 Poland CA, Duffin R, Kinloch I, et al. Carbon nanotubes introduced into the abdominal cavity of mice show asbestos-like pathogenicity in a pilot study. *Nat Nanotechnol* 2008; **3**: 423–428.
  - 7 Murphy FA, Schinwald A, Poland CA, Donaldson K. The mechanism of pleural inflammation by long carbon nanotubes: interaction of long fibres with macrophages stimulates them to amplify pro-inflammatory responses in mesothelial cells. *Part Fibre Toxicol* 2012; **9**: 8.
  - 8 Takagi A, Hirose A, Nishimura T, et al. Induction of mesothelioma in p53+/- mouse by intraperitoneal application of multi-wall carbon nanotube. *J Toxicol Sci* 2008; **33**: 105–116.
  - 9 Sakamoto Y, Nakae D, Fukumori N, et al. Induction of mesothelioma by a single intrascrotal administration of multi-wall carbon nanotube in intact male Fischer 344 rats. *J Toxicol Sci* 2009; **34**: 65–76.
  - 10 Nagai H, Okazaki Y, Chew S, et al. Diameter of multi-walled carbon nanotubes is a critical factor in mesothelial injury and subsequent carcinogenesis. *Proc Natl Acad Sci U S A* 2011; **108**: E1330–E1338.
  - 11 Toyokuni S. Genotoxicity and carcinogenicity risk of carbon nanotubes. *Adv Drug Deliv Rev* 2013; **65**: 2098–2110.
  - 12 Toyokuni S. Mechanisms of asbestos-induced carcinogenesis. *Nagoya J Med Sci* 2009; **71**: 1–10.
  - 13 Nagai H, Toyokuni S. Biopersistent fiber-induced inflammation and carcinogenesis: lessons learned from asbestos toward safety of fibrous nanomaterials. *Arch Biochem Biophys* 2010; **502**: 1–7.
  - 14 Jiang L, Akatsuka S, Nagai H, et al. Iron overload signature in chrysotile-induced malignant mesothelioma. *J Pathol* 2012; **228**: 366–377.
  - 15 Nagai H, Toyokuni S. Differences and similarities between carbon nanotubes and asbestos fibers during mesothelial carcinogenesis. *Cancer Sci* 2012; **103**: 1378–1390.
  - 16 Donaldson K, Cullen RT. Chemiluminescence of asbestos-activated macrophages. *Br J Exp Pathol* 1984; **65**: 81–90.
  - 17 Nyberg P. Polyvinylpyridine-N-oxide and carboxymethyl cellulose inhibit mineral dust-induced production of reactive oxygen species by human macrophages. *Environ Res* 1991; **55**: 157–164.
  - 18 Doll NJ, Stankus RP, Goldbach S, Salvaggio JE. *In vitro* effect of asbestos fibers on polymorphonuclear leukocyte function. *Int Arch Allergy Appl Immunol* 1982; **68**: 17–21.
  - 19 Korkina LG, Durnev AD, Suslova TB, Cheremisina ZP, Daugel-Dauge NO, Afanas'ev IB. Oxygen radical-mediated mutagenic effect of asbestos on human lymphocytes: suppression by oxygen radical scavengers. *Mutat Res* 1992; **265**: 245–253.
  - 20 Ishizaki T, Yano E, Urano N, Evans PH. Crocidolite-induced reactive oxygen metabolites generation from human polymorphonuclear leukocytes. *Environ Res* 1994; **66**: 208–216.
  - 21 Kinnula VL, Raivio KO, Linnainmaa K, Ekman A, Klockars M. Neutrophil and asbestos fiber-induced cytotoxicity in cultured human mesothelial and bronchial epithelial cells. *Free Radic Biol Med* 1995; **18**: 391–399.
  - 22 Iwata T, Yano E. Reactive oxygen metabolite production induced by asbestos and glass fibers: effect of fiber milling. *Ind Health* 2003; **41**: 32–38.
  - 23 Chen B, Liu Y, Song WM, Hayashi Y, Ding XC, Li WH. *In vitro* evaluation of cytotoxicity and oxidative stress induced by multiwalled carbon nanotubes in murine RAW 264.7 macrophages and human A549 lung cells. *Biomed Environ Sci* 2011; **24**: 593–601.
  - 24 Imada I, Sato EF, Miyamoto M, et al. Analysis of reactive oxygen species generated by neutrophils using a chemiluminescence probe L-012. *Analyt Biochem* 1999; **271**: 53–58.
  - 25 Meng ZX, Zheng W, Ding MH, et al. Fabrication and characterization of elastomeric polyester/carbon nanotubes nanocomposites for biomedical application. *J Nanosci Nanotechnol* 2011; **11**: 3126–3133.
  - 26 Jiang L, Nagai H, Ohara H, et al. Characteristics and modifying factors of asbestos-induced oxidative DNA damage. *Cancer Sci* 2008; **99**: 2142–2151.
  - 27 Nagai H, Ishihara T, Lee WH, et al. Asbestos surface provides a niche for oxidative modification. *Cancer Sci* 2011; **102**: 2118–2125.
  - 28 Kubo Y, Takenaka H, Nagai H, Toyokuni S. Distinct affinity of nuclear proteins to the surface of chrysotile and crocidolite. *J Clin Biochem Nutr* 2012; **51**: 221–226.
  - 29 Kumar S, Bandyopadhyay U. Free heme toxicity and its detoxification systems in human. *Toxicol Lett* 2005; **157**: 175–188.
  - 30 Toyokuni S, Sagripanti JL. Iron-mediated DNA damage: sensitive detection of DNA strand breakage catalyzed by iron. *J Inorg Biochem* 1992; **47**: 241–248.
  - 31 Toyokuni S, Sagripanti JL. DNA single- and double-strand breaks produced by ferric nitrilotriacetate in relation to renal tubular carcinogenesis. *Carcinogenesis* 1993; **14**: 223–227.
  - 32 Toyokuni S. Role of iron in carcinogenesis: cancer as a ferrototoxic disease. *Cancer Sci* 2009; **100**: 9–16.
  - 33 Kumar V, Abbas AK, Aster JC. *Robbins Basic Pathology*. 9th ed. Philadelphia, PA: Elsevier Saunders, 2013.
  - 34 Martinon F. Signaling by ROS drives inflammasome activation. *Eur J Immunol* 2010; **40**: 616–619.
  - 35 Sun B, Wang X, Ji Z, Li R, Xia T. NLRP3 inflammasome activation induced by engineered nanomaterials. *Small* 2013; **9**: 1595–1607.
  - 36 Nagai H, Okazaki Y, Chew SH, et al. Intraperitoneal administration of tangled multiwalled carbon nanotubes of 15 nm in diameter does not induce mesothelial carcinogenesis in rats. *Pathol Int* 2013; **63**: 457–462.