

RESEARCH ARTICLE

Informing the Historical Record of Experimental Nonhuman Primate Infections with Ebola Virus: Genomic Characterization of USAMRIID Ebola Virus/H.sapiens-tc/COD/1995/Kikwit-9510621 Challenge Stock “R4368” and Its Replacement “R4415”

Jeffrey R. Kugelman¹, Cynthia A. Rossi¹, Michael R. Wiley¹, Jason T. Ladner¹, Elyse R. Nagle¹, Bradley P. Pfeffer¹, Karla Garcia¹, Karla Prieto¹, Jiro Wada², Jens H. Kuhn², Gustavo Palacios^{1*}

1 United States Army Medical Research Institute of Infectious Diseases (USAMRIID), Fort Detrick, Frederick, Maryland, United States of America, **2** Integrated Research Facility at Fort Detrick (IRF-Frederick), National Institute of Allergy and Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, Maryland, United States of America

* gustavo.f.palacios.ctr@mail.mil



CrossMark
click for updates

OPEN ACCESS

Citation: Kugelman JR, Rossi CA, Wiley MR, Ladner JT, Nagle ER, Pfeffer BP, et al. (2016) Informing the Historical Record of Experimental Nonhuman Primate Infections with Ebola Virus: Genomic Characterization of USAMRIID Ebola Virus/H.sapiens-tc/COD/1995/Kikwit-9510621 Challenge Stock “R4368” and Its Replacement “R4415”. *PLoS ONE* 11(3): e0150919. doi:10.1371/journal.pone.0150919

Editor: Alexander N Freiberg, University of Texas Medical Branch, UNITED STATES

Received: October 7, 2015

Accepted: February 14, 2016

Published: March 22, 2016

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All sequence data will be available from GenBank (accession numbers KT582109, KT762961, and KT762962).

Funding: The content of this publication does not necessarily reflect the views or policies of the US Department of Defense, the US Department of Health and Human Services, the US Department of the Army, or the institutions and companies affiliated with the authors. This work was funded in part through

Abstract

The creation of licensed medical countermeasures against Select Agents such as Ebola virus (EBOV) is critically dependent on the use of standardized reagents, assays, and animal models. We performed full genome reconstruction, population genomics, contaminant analysis, and characterization of the glycoprotein gene editing site of historical United States Army Medical Research Institute of Infectious Diseases (USAMRIID) nonhuman-primate challenge stock Ebola virus Kikwit “R4368” and its 2014 replacement “R4415.” We also provide characterization of the master stock used to create “R4415.” The obtained data are essential to understanding the quality of the seed stock reagents used in pivotal animal studies that have been used to inform medical countermeasure development. Furthermore, these data might add to the understanding of the influence of EBOV variant populations on pathogenesis and disease outcome and inform attempts to avoid the evolution of EBOV escape mutants in response to current therapeutics. Finally, as the primary challenge stocks have changed over time, these data will provide a baseline for understanding and correlating past and future animal study results.

Introduction

Ebola virus disease (EVD) is a frequently lethal human viral hemorrhagic fever caused by four distinct ebolaviruses (Bundibugyo virus, Ebola virus, Sudan virus, and Tai Forest virus). EVD

Battelle Memorial Institute's prime contract with the US National Institute of Allergy and Infectious Diseases (NIAID) under Contract No. HHSN272200700016I (J.H.K.). A subcontractor to Battelle Memorial Institute who performed this work is: J.H.K., an employee of Tunnell Government Services, Inc. Tunnell Government Services provided support in the form of salaries, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. All work under contract HHSN272200700016I is performed under the agreement/understanding that it will belong to the client and will be affiliated directly to the NIH/NIAID Integrated Research Facility at Fort Detrick. The specific roles of these authors are articulated in the 'author contributions' section.

Competing Interests: The authors have the following interests: Jens H. Kuhn is employed by Tunnell Government Services, Inc. There are no patents, products in development or marketed products to declare. This employment does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors.

occurs sporadically and usually affects no more than several hundred people during an outbreak (reviewed in [1]). In 2014, Ebola virus (EBOV) was identified as the etiological agent of an unprecedented EVD outbreak that started in Western Africa in late 2013 and has since caused 28,637 cases and 11,315 deaths (as of January 3, 2016) [2].

Taxonomically, EBOV is the only member of the species *Zaire ebolavirus* in the genus *Ebolavirus* (*Mononegavirales: Filoviridae*) [3]. The EBOV genome is rather conserved over time and geographic distances, which may be due to genetic bottlenecks in the yet-unidentified host reservoir [1, 4]. For instance, the EBOV variant that caused the 2013–present outbreak in Western Africa (Makona [5]) differs from those that caused EVD outbreaks in Zaire in 1976 (Yambuku [6, 7]) and Zaire's successor country Democratic Republic of the Congo in 1995 (Kikwit [8]) by less than 3% over the entire ≈ 19 kb genome [9].

The development, evaluation, and final licensure of medical countermeasures (MCMs) against EVD in the US is critically dependent on standardized animal models of filovirus infection and standardized assays and reagents [10–12], including well-characterized virus stocks. Ebola virus/H.sapiens-tc/COD/1976/Yambuku-Mayinga (EBOV/Yam-May) is the best characterized EBOV isolate and has been used for the majority of *in vitro* experiments [1, 7, 13]. Ebola virus/H.sapiens-tc/COD/1995/Kikwit-9510621 (EBOV/Kik-9510621), on the other hand, has become the most used EBOV isolate for animal, and in particular nonhuman primate (NHP), experimentation in the United States (US) [1, 13, 14]. Unfortunately, this isolate has been passaged/maintained by different procedures in different locations and even within the same institutes. These variable procedures result in NHP challenge stocks of different quality and possibly in different experimental outcomes upon use. In addition, only one of these NHP challenge stocks has been genomically characterized to assess potential mutations relative to the published consensus genome sequence [14].

All ebolaviruses make use of co-transcriptional editing of their glycoprotein-encoding *GP* genes to access three partially overlapping open reading frames (ORF) [15, 16]. Editing occurs at a distinct editing site in the *GP* gene that typically consists of seven consecutive uridylyls (7U). Read-through results in the transcription of a cognate 7 adenylyl mRNA (7A mRNA) and thereby in the expression of pre-sGP, a protein precursor that is proteolytically processed to a secreted homodimeric glycoprotein (sGP) and a probably monomeric secreted peptide (Δ -peptide). Stuttering of the ebolavirus RNA-dependent RNA polymerase (L) at the editing site results in transcription of mRNAs that contain various numbers of adenylyls. The majority of non-7A mRNAs contain eight adenylyls, leading to the expression of the homotrimeric ebolavirus spike glycoprotein GP_{1,2}, or six or nine adenylyls, leading to expression of the small soluble glycoprotein (ssGP) [15–20]. The functions of sGP, ssGP, and Δ -peptide are largely unknown but a role for sGP in immune evasion has been described [21]. Editing appears to be tightly controlled, with the ratio of proteins expressed from a 7U virus of roughly sGP:GP_{1,2}:ssGP = 75%:25%:5% [17, 22].

EBOV, Sudan virus (SUDV), and possibly other ebolaviruses adapt to different *in vitro* and *in vivo* environments [14, 23–25]. These adaptations include changes in the EBOV and SUDV *GP* gene editing sites. For instance, during serial passage of a 7U EBOV/Yam-May isolate in Vero E6 cells, a viral population evolved that predominantly contained an 8U editing site. The opposite occurs *in vivo*: 8U EBOV/Yam-May populations converted to 7U populations in guinea pigs [10]. These changes may be related to selective advantages linked to the controlled expression of GP_{1,2} and/or sGP. Preliminary evaluations indicate that a 7U→8U mutation in EBOV does not alter pathogenesis in guinea pigs or nonhuman primates [25]. However, the observations described above indicate that there may be selective advantages associated with the ratio of expressed sGP:GP_{1,2} in different environments (8U-containing viruses predominantly express GP_{1,2} rather than sGP). In addition, a recent study reported subtle, but

significant, differences in disease course and severity between nonhuman primates exposed to 7U or 8U virus stock variants [26]. All of these findings underscore the importance of fully characterizing viral stocks used in MCM development.

Here, we report the coding-complete genome sequence (see [27] for sequencing terminology), population characterization, and contaminant analysis of an EBOV/Kik-9510621 challenge stock “R4368” (passage 4, 8U) that was used in the past for NHP studies at the United States Army Medical Research Institute of Infectious Diseases (USAMRIID) at Fort Detrick, Maryland. We further report the complete genome of sequence at the same level of characterization of challenge stock “R4415” (passage 3, 7U), which replaced “R4368” (passage 4, 8U) in 2014, and the master stock used to establish this new stock (“R4414” (passage 2, 7U)).

History of USAMRIID EBOV/Kik-9510621 challenge stocks

A clinical specimen (designated Centers for Disease Control and Prevention Special Pathogens Branch Log [CDC SPBLOG] 9510621) was obtained from an EBOV-infected 65-year old female patient during an EVD outbreak that occurred in 1995 around Kikwit, Zaire (today Democratic Republic of the Congo, COD). The patient’s disease onset was recorded as April 29, 1995. She was hospitalized on May 1, 1995, and died on May 5, 1995. The clinical specimen, most likely plasma or serum, was obtained from the patient on May 4, 1995. How the patient became infected and what medical care she may have received is unclear. Unfortunately, chain-of-custody records that further detail the origin of 9510621 or its shipment to the CDC are not available anymore, and EBOV titration was not attempted from this specimen prior to cell-culture passage.

The first passage of EBOV/Kik-9510621, designated “virus seed pool (VSP) 807223” (Fig 1), was conducted at the CDC using grivet (*Chlorocebus aethiops*) Vero E6 cells (ATCC #CRL-1586). The multiplicity of infection (MOI) used for this passage is unknown. Virus was harvested after a 6-day incubation period on May 19, 1995, but the method of harvest (initial freeze/thaw, clarification by centrifugation etc.) is not indicated in the available records. A second passage of virus, was conducted at the CDC, again using Vero E6 cells, and resulted in VSP 807224 (MOI unknown). Virus was harvested after an 8-day incubation, using a “freeze/thaw” method, on May 29, 1995. A titer of $3.2E+06$ was determined for “VSP 807224” in January 2008 using the TCID₅₀/Reed-Munch viral titration method. “VSP 807224” was transferred from CDC between late May and late June 1995 to USAMRIID.

At USAMRIID, “VSP 807224” was stored frozen at -70°C and designated there as “135” (passage 2). After an additional passage in Vero E6 cells, the virus, now designated “134” (passage 3), was sequenced by Chain *et al.* using classical dideoxynucleotide sequencing. “134” (passage 3) is cell culture-adapted, consisting predominantly of viruses with an 8 uridylyl (8U) glycoprotein (GP) gene editing site. A coding-complete genome for “134” was deposited on July 24, 2003, in GenBank under accession #AY354458. “135” (passage 2) was amplified at USAMRIID at least three more times (and in the process was depleted), leading to three separate NHP challenge stocks: “R4367” (passage 3, MOI 0.001), “R4369” (passage 3, MOI 0.01), and “16502” (passage 3, MOI not available). The genomic characterization of stock “16502” (passage 3) and its evolution in a nonhuman primate study over several days was previously described [14] (Fig 1). “R4367” (passage 3) was passaged on Vero E6 cells at an MOI of 0.001, collected, and clarified on day 5 (June 26, 2011), and then passaged one more time at an MOI of 0.01 on Vero E6 cells to yield “R4368” (passage 4). “R4368” (passage 4) was collected and clarified on day 4 (July 24, 2011). “R4369” (passage 3) was passaged on Vero E6 cells at an MOI of 0.01 and collected and clarified on day 4 (October 29, 2011). Titers of “R4367” ($3.31E+06$ plaque-forming units (pfu)/ml), “R4368” ($4.56E+06$ pfu/ml), and “R4369” ($7.01E+06$ pfu/ml) were established using




Fig 1. History of Ebola virus variant Kikwit (isolate 9510621) challenge stocks used at the United States Army Medical Research Institute of Infectious Diseases (USAMRIID), CDC, Centers for Disease Control and Prevention; NHP, nonhuman primate; SPBLOG, Special Pathogens Branch Log; UTMB, University of Texas Medical Branch; VSP, virus seed pool.

doi:10.1371/journal.pone.0150919.g001

an agarose-based plaque assay [21]. Each of these three preparations was harvested after 2–3+ cytopathic effect (CPE) had developed.

Independently at UTMB, "VSP 807223" was passaged one time on Vero E6 cells, harvested after a 10-day incubation period on February 21, 2012, and stored in 1.0-ml aliquots as "WRC000121" (passage 2; MOI unknown). A titer of 1.8E+07 pfu/ml was determined (titration method unknown). On May 1, 2012, UTMB transferred "WRC000121" to USAMRIID,

where the virus was stored at -70°C and designated as “R4414” (passage 2). This material was passaged at an MOI of 0.001 on Vero E6 cells to prepare NHP challenge stock “R4415” (passage 3), which was harvested on June 11, 2013, after a 12-day incubation period (time to develop 2–3+ CPE) with a titer of $1.31\text{E}+06$ pfu/ml determined by agarose-based plaque assay [21]. The primary USAMRIID NHP challenge stock “R4368” (passage 4, 8U) was replaced by “R4415” (passage 3, 7U) in December of 2014 and is now USAMRIID’s primary NHP challenge stock.

Materials and Methods

Vials (1 ml) of EBOV/Kik-9510621 challenge stocks were stored and maintained at USAMRIID at -70°C . “R4368” (passage 4) was stored since July 24, 2011, and thawed on March 2, 2012; “R4414” (passage 2) and “R4415” (passage 3) were stored since June 11, 2013, and were thawed in January 2014. An aliquot of 100 μl of each thawed virus stock was placed in 3:1 TRIzol (Life Technologies, Carlsbad, CA). Nucleic acids were isolated from TRIzol-treated material, and genome sequence was determined on Illumina technology (MiSeq or HiSeq) with EBOV-specific oligonucleotides following sample preparation performed as described in [14]. The consensus genomes were generated via reference alignment to EBOV/Kik-9510621 challenge stock “134” (passage 3) (GenBank #AY354458) using SeqMan nGen (DNASTAR). The resulting “R4368” (passage 4), “R4414” (passage 2), and “R4415” (passage 3) sequences were deposited in GenBank under accession numbers KT582109, KT762961, and KT762962, respectively. The population genetics assembly files and sequence-independent, single-primer amplification (SISPA) raw sequence used for contaminant analysis are available in BioSample database under SRS1037973, SRS1041428, and SRS1041442.

For rapid amplification of cDNA ends (RACE), “R4415” (passage 3) RNAs were extracted with Zymo Direct-Zol (Zymo Research Corporation, Irvine, CA) from cell-culture supernatant in TRIzol according to the manufacturers' instructions. SMARTer RACE 5'/3' kit (Clontech Laboratories, Inc., Mountain View, CA) was used to amplify both 5' and 3' untranslated regions (UTRs) of the virus genome from the extracted RNAs. The kit's two-stage nested PCR protocol was found to be optimal. The gene-specific primers for each RACE experiment are as follows 5' RACE (outer primer): ATTACCAGAGTTGATTAGTGTG; 5' RACE (inner primer): TTAAATAACGAAAGGAGTC; 3' RACE (outer primer): TGAATCTCCAATCCTCTAAGTA; 3' RACE (inner primer): AAGGGATTTTCAACTGAGCACACT. Amplification primer removal, duplicate removal, low average quality exclusion ($\leq Q30$), and quality trimming was performed. Viral assemblies were completed in DNASTar Lasergene nGen (Madison, WI) with $\approx 4 \times 10^5$ reads. Only single nucleotide polymorphism (SNPs) present in the population above the 2% threshold are presented in this report (however, the alignment files are provided in SRA if a less conservative approach is desired). Considering this threshold, a target depth of 200 requires an SNP to have ≥ 4 supporting reads prior to being called a SNP. The depths are reported in the tables presented in this text for all samples (below 200 depth, calls should be viewed with increasing skepticism). A consensus change is defined here as a change relative to the published sequence for EBOV/Kik-9510621 “134” (GenBank accession # AY354458) present in $\geq 50\%$ of the population. Below that threshold, SNPs are considered subclonal substitutions and part of a minority subpopulation of the virus.

Ethics Statement

Research has been reviewed for compliance with dual-use guidelines and approved for publication by the USAMRIID Institute Biosafety Committee (IBC) and the Operational Security office.

Table 1. Intrahost single nucleotide variants (iSNVs) of EBOV/Kik-9510621 “R4368” (passage 4) comprising $\geq 2\%$ of total population as compared to “134” (GenBank #AY354458).

| Reference position (“134” (passage 3)) | Reference base | SNP base | SNP % | Codon | Gene | Depth |
|--|----------------|----------|-------|-----------------------|------|-------|
| 5,878 | T | g | 2.84 | | | 1374 |
| 6,139 | C | t | 2.07 | P (CCA) @34 L (CtA) | GP | 1595 |
| 6,179 | G | t | 8.57 | E (GAG) @47 D (GAT) | GP | 1541 |
| 7,298 | T | c | 3.98 | Synonymous | GP | 804 |
| 7,327 | C | T | 99.61 | P (CCA) @430 L (CTA) | GP | 507 |
| 10,833 | G | a | 2.35 | R (AGA) @163 K (AaA) | VP24 | 1533 |
| 16,365 | A | g | 2.76 | Q (CAG) @1595 R (CgG) | L | 1560 |

SNP, single nucleotide polymorphism.

doi:10.1371/journal.pone.0150919.t001

Results

Compared to EBOV/Kik-9510621 “134” (passage 3), “R4368” (passage 4) acquired only one consensus-level substitution (nucleotide position 7,327). Six intrahost single nucleotide variants (iSNVs) ($\geq 2\%$ of the population) were detected and are reported in Table 1. Compared to EBOV/Kik-9510621 “134” (passage 3), “R4414” (passage 2) acquired three consensus-level substitutions (nucleotide positions 6,179, 7,327, 10,833). Eleven iSNVs ($\geq 2\%$ of the population) were detected and are reported in Table 2. An additional passage-acquired mutation was observed in the challenge stock “R4415” (passage 3) (nucleotide position 7,669) when compared to “R4414” (passage 2). Thirteen iSNVs ($\geq 2\%$ of the population) were detected and are reported in Table 3. No significant contaminants were detected. The genotype ratios at the GP editing site for all three stocks are detailed in Table 4. As expected based on the “134” (passage 3) progenitor sequence, “R4368” (passage 4) predominantly encodes the 8U genotype (85.0%). “R4414” (passage 2) and “R4415” (passage 3) predominantly encode the 7U genotype.

Table 2. Intrahost single nucleotide variants (iSNVs) of EBOV/Kik-9510621 “R4414” (passage 2) comprising $\geq 2\%$ of total population as compared to “134” (GenBank #AY354458).

| Reference position (“134” (passage 3)) | Reference base | SNP base | SNP % | Codon | Gene | Depth |
|--|----------------|----------|--------|---------------------|------|-------|
| 1,401 | G | a | 2.08 | G:GGT @ 311 → D:GaT | NP | 1,008 |
| 5,830 | T | a | 2.18 | | VP40 | 1,789 |
| 6,179 | G | t | 100.00 | E:GAG @ 47 → D:GAT | GP | 2,887 |
| 6,231 | T | c | 3.97 | S:TCA @ 65 → P:cCA | GP | 3,956 |
| 6,384 | C | a | 4.33 | P:CCT @ 116 → T:aCT | GP | 5,264 |
| 7,327 | C | t | 99.90 | P:CCA @ 430 → L:CtA | GP | 995 |
| 7,669 | C | t | 33.00 | T:ACA @ 544 → I:AtA | GP | 912 |
| 10,344 | C | a | 4.60 | | VP24 | 5,108 |
| 10,833 | G | a | 99.60 | R:AGA @ 163 → K:AaA | VP24 | 3,422 |
| 11,283 | A | c | 2.89 | | VP24 | 3,361 |
| 11,498 | G | a | 6.05 | | | 1,123 |
| 12,065 | G | a | 4.21 | G:GGT @ 162 → S:aGT | L | 4,798 |
| 12,153 | G | t | 5.38 | W:TGG @ 191 → L:TtG | L | 5,246 |
| 14,184 | C | t | 2.13 | S:TCG @ 868 → L:TtG | L | 1,880 |

SNP, single nucleotide polymorphism.

doi:10.1371/journal.pone.0150919.t002

Table 3. Intrahost single nucleotide variants (iSNVs) of EBOV/Kik-9510621 “R4415” (passage 3) comprising $\geq 2\%$ of total population as compared to “134” (GenBank #AY354458).

| Reference position (“134” (passage 3)) | Reference base | SNP base | SNP % | Codon | Gene | Depth |
|--|----------------|----------|--------|----------------------|------|--------|
| 520 | T | c | 10.60 | S:TCT @ 17 → S:TCc | NP | 4,330 |
| 530 | T | c | 10.15 | Y:TAC @ 21 → H:cAC | NP | 4,403 |
| 542 | T | c | 9.84 | L:TTG @ 25 → L:cTG | NP | 4,483 |
| 606 | T | c | 10.70 | V:GTA @ 46 → A:GcA | NP | 4,831 |
| 1,274 | A | g | 7.83 | R:AGG @ 269 → G:gGG | NP | 5,62 |
| 5,830 | T | c | 3.20 | | VP40 | 2,871 |
| 6,179 | G | t | 99.90 | E:GAG @ 47 → D:GAt | GP | 8,337 |
| 6,384 | C | a | 3.4 | P:CCT @ 116 → T:aCT | GP | 11,383 |
| 7,327 | C | t | 99.940 | P:CCA @ 430 → L:CtA | GP | 1,048 |
| 7,669 | C | t | 98.70 | T:ACA @ 544 → I:AtA | GP | 1,486 |
| 10,344 | C | a | 4.20 | | VP24 | 9,735 |
| 10,833 | G | a | 100.00 | R:AGA @ 163 → K:AaA | VP24 | 4,642 |
| 11,283 | A | c | 3.91 | | VP24 | 1,844 |
| 11,498 | G | a | 5.15 | | | 466 |
| 12153 | G | a | 4.95 | W:TGG @ 191 →.:TaG | L | 4,910 |
| 13994 | C | a | 4.65 | Q:CAA @ 805 → K:aAA | L | 7,614 |
| 16247 | T | c | 2.19 | S:TCA @ 1556 → P:cCA | L | 2,838 |

SNP, single nucleotide polymorphism.

doi:10.1371/journal.pone.0150919.t003

Discussion

We have provided here a concise report on the history and genomic characterization of the USAMRIID Ebola virus/H.sapiens-tc/COD/1995/Kikwit-9510621 NHP challenge stocks “R4368” (passage 4) and “R4415” (passage 3), as well as the “R4415” (passage 3) predecessor, “R4414 (passage 2).” “R4368” (passage 4) was used between July 2011 and December 2014 in both *in vitro* and *in vivo* Ebola virus experimentation, including pathogenesis studies and candidate medical countermeasure evaluation. “R4415” (passage 3) replaced “R4368” (passage 4) in December 2014 and has been used for most pathogenesis and medical countermeasure evaluation research since then. This work provides a framework for genomic comparison between past experiments as challenge stocks are replaced to address propagation issues and depletion. Characterization of these NHP challenge stocks was completed to the level of “Coding Complete” plus population-level characterization in the case of “R4368” (passage 4) and “Finished” in the case of “R4415” (passage 3) [27]. This characterization includes genome reconstruction (excluding determination of the 3’ and 5’ UTRs in the case of “R4414” (passage 2) and “R4415” (passage 3), characterization of intrahost variants (iSNVs), and determination of absence of contaminants. Studies to determine the role of the identified iSNVs in interactions with the

Table 4. GP gene editing site composition in EBOV/Kik-9510621 NHP challenge stocks.

| | 6U/9U (ssGP phenotype) | 7U (sGP phenotype) | 8U (GP _{1,2} phenotype) |
|---------------------|------------------------|--------------------|----------------------------------|
| “R4368” (passage 4) | 3.8% (373) | 11.2% (1,090) | 85.0% (8,300) |
| “R4414” (passage 2) | 0.6% (149) | 97.5% (23222) | 1.8% (439) |
| “R4415” (passage 3) | 0.3% (40) | 88.8% (11231) | 10.9% (1378) |

GP, glycoprotein; sGP, soluble glycoprotein; ssGP, small secreted glycoprotein; 7–9U, 7–9 uridylyl glycoprotein (GP) gene editing site.

doi:10.1371/journal.pone.0150919.t004

host (e.g., in the immune response) are being considered for expansion of this body of work. This level of characterization is crucial for studies evaluating the possibility of EBOV escape from candidate therapeutics or vaccines, as minority variants can play an important phenotypic role in viral escape [28].

Acknowledgments

The authors thank Dr. Pierre E. Rollin (CDC, Atlanta, GA) and Dr. Thomas W. Geisbert (UTMB) for providing the background information on EBOV/Kik-951062 stocks, the USAMRIID virus production team of Brian Kearney, Scott Olschner, Priscilla Williams, Dr. Anna N. Honko, and Joshua C. Johnson for experimental support, and Laura Bollinger (IRF-Frederick) for critically editing the manuscript.

Author Contributions

Conceived and designed the experiments: GP. Performed the experiments: JRK CAR MRW JTL ERN BPP KG KP. Analyzed the data: JRK JTL JW JHK GP. Wrote the paper: JRK CAR JHK GP.

References

1. Kuhn JH. Filoviruses. A compendium of 40 years of epidemiological, clinical, and laboratory studies. Archives of Virology Supplementum, vol. 20. Vienna, Austria: SpringerWienNewYork; 2008.
2. World Health Organization. Ebola Situation Reports. Available: <http://apps.who.int/ebola/ebola-situation-reports>. Geneva, Switzerland: 2016.
3. Kuhn JH, Becker S, Ebihara H, Geisbert TW, Jahrling PB, Kawaoka Y, et al. Family Filoviridae. In: King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ, editors. Virus Taxonomy—Ninth Report of the International Committee on Taxonomy of Viruses. London, UK: Elsevier/Academic Press; 2011. p. 665–71.
4. Wahl-Jensen V, Radoshitzky SR, de Kok-Mercado F, Taylor SL, Bavari S, Jahrling PB, et al. Role of Rodents and Bats in Human Viral Hemorrhagic Fevers. In: Singh SK, Ruzek D, editors. Viral Hemorrhagic Fevers. Boca Raton, Florida, USA: Taylor & Francis/CRC Press; 2013. p. 99–127.
5. Kuhn JH, Andersen KG, Baize S, Bao Y, Bavari S, Berthet N, et al. Nomenclature- and database-compatible names for the two Ebola virus variants that emerged in Guinea and the Democratic Republic of the Congo in 2014. Viruses. 2014; 6(11):4760–99. doi: [10.3390/v6114760](https://doi.org/10.3390/v6114760) PMID: [25421896](https://pubmed.ncbi.nlm.nih.gov/25421896/); PubMed Central PMCID: [PMC4246247](https://pubmed.ncbi.nlm.nih.gov/PMC4246247/).
6. Kuhn JH, Lofts LL, Kugelman JR, Smither SJ, Lever MS, van der Groen G, et al. Reidentification of Ebola virus “E718” and “ME” as Ebola virus/H.sapiens-tc/COD/1976/Yambuku-Ecran. Genome Announc. 2014; 2(6):e01178–14. PubMed Central PMCID: [PMC4192379](https://pubmed.ncbi.nlm.nih.gov/PMC4192379/).
7. Kuhn JH, Andersen KG, Bao Y, Bavari S, Becker S, Bennett RS, et al. Filovirus RefSeq entries: evaluation and selection of filovirus type variants, type sequences, and names. Viruses. 2014; 6(9):3663–82. doi: [10.3390/v6093663](https://doi.org/10.3390/v6093663) PMID: [25256396](https://pubmed.ncbi.nlm.nih.gov/25256396/); PubMed Central PMCID: [PMC4189044](https://pubmed.ncbi.nlm.nih.gov/PMC4189044/).
8. Kuhn JH, Bao Y, Bavari S, Becker S, Bradfute S, Brister JR, et al. Virus nomenclature below the species level: a standardized nomenclature for natural variants of viruses assigned to the family Filoviridae. Arch Virol. 2013; 158(1):301–11. doi: [10.1007/s00705-012-1454-0](https://doi.org/10.1007/s00705-012-1454-0) PMID: [23001720](https://pubmed.ncbi.nlm.nih.gov/23001720/); PubMed Central PMCID: [PMC3535543](https://pubmed.ncbi.nlm.nih.gov/PMC3535543/).
9. Baize S, Pannetier D, Oestereich L, Rieger T, Koivogui L, Magassouba N, et al. Emergence of Zaire Ebola virus disease in Guinea. N Engl J Med. 2014; 371(15):1418–25. doi: [10.1056/NEJMoa1404505](https://doi.org/10.1056/NEJMoa1404505) PMID: [24738640](https://pubmed.ncbi.nlm.nih.gov/24738640/).
10. Korch GW Jr, Niemi SM, Bergman NH, Carucci DJ, Ehrlich SA, Gronvall GK, et al. Animal models for assessing countermeasures to bioterrorism agents. Washington, DC, USA: Institute for Laboratory Animal Research (ILAR), Division on Earth and Life Studies, The National Academies of Sciences (NAS); 2011.
11. Hirschberg R, Ward LA, Kilgore N, Kumat R, Schiltz H, Albrecht MT, et al. Challenges, progress, and opportunities: proceedings of the filovirus medical countermeasures workshop. Viruses. 2014; 6(7):2673–97. doi: [10.3390/v6072673](https://doi.org/10.3390/v6072673) PMID: [25010768](https://pubmed.ncbi.nlm.nih.gov/25010768/); PubMed Central PMCID: [PMC4113787](https://pubmed.ncbi.nlm.nih.gov/PMC4113787/).
12. Kilgore N, Nuzum EO. An interagency collaboration to facilitate development of filovirus medical countermeasures. Viruses. 2012; 4(10):2312–6. doi: [10.3390/v4102312](https://doi.org/10.3390/v4102312) PMID: [23202465](https://pubmed.ncbi.nlm.nih.gov/23202465/); PubMed Central PMCID: [PMC3497053](https://pubmed.ncbi.nlm.nih.gov/PMC3497053/).

13. Kuhn JH, Dodd LE, Wahl-Jensen V, Radoshitzky SR, Bavari S, Jahrling PB. Evaluation of perceived threat differences posed by filovirus variants. *Biosecur Bioterror*. 2011; 9(4):361–71. Epub 2011/11/11. doi: [10.1089/bsp.2011.0051](https://doi.org/10.1089/bsp.2011.0051) PMID: [22070137](https://pubmed.ncbi.nlm.nih.gov/22070137/); PubMed Central PMCID: PMC3233913.
14. Kugelman JR, Lee MS, Rossi CA, McCarthy SE, Radoshitzky SR, Dye JM, et al. Ebola virus genome plasticity as a marker of its passaging history: a comparison of in vitro passaging to non-human primate infection. *PLoS One*. 2012; 7(11):e50316. doi: [10.1371/journal.pone.0050316](https://doi.org/10.1371/journal.pone.0050316) PMID: [23209706](https://pubmed.ncbi.nlm.nih.gov/23209706/); PubMed Central PMCID: PMC3509072.
15. Volchkov VE, Becker S, Volchkova VA, Ternovoj VA, Kotov AN, Netesov SV, et al. GP mRNA of Ebola virus is edited by the Ebola virus polymerase and by T7 and vaccinia virus polymerases. *Virology*. 1995; 214(2):421–30. PMID: [8553543](https://pubmed.ncbi.nlm.nih.gov/8553543/).
16. Sanchez A, Trappier SG, Mahy BW, Peters CJ, Nichol ST. The virion glycoproteins of Ebola viruses are encoded in two reading frames and are expressed through transcriptional editing. *Proc Natl Acad Sci U S A*. 1996; 93(8):3602–7. PMID: [8622982](https://pubmed.ncbi.nlm.nih.gov/8622982/); PubMed Central PMCID: PMC39657.
17. Mehedi M, Falzarano D, Seebach J, Hu X, Carpenter MS, Schnittler HJ, et al. A new Ebola virus non-structural glycoprotein expressed through RNA editing. *J Virol*. 2011; 85(11):5406–14. Epub 2011/03/18. doi: [10.1128/jvi.02190-10](https://doi.org/10.1128/jvi.02190-10) PMID: [21411529](https://pubmed.ncbi.nlm.nih.gov/21411529/); PubMed Central PMCID: PMC3094950.
18. Barrientos LG, Martin AM, Rollin PE, Sanchez A. Disulfide bond assignment of the Ebola virus secreted glycoprotein SGP. *Biochem Biophys Res Commun*. 2004; 323(2):696–702. doi: [10.1016/j.bbrc.2004.08.148](https://doi.org/10.1016/j.bbrc.2004.08.148) PMID: [15369806](https://pubmed.ncbi.nlm.nih.gov/15369806/).
19. Volchkova VA, Feldmann H, Klenk HD, Volchkov VE. The nonstructural small glycoprotein sGP of ebola virus is secreted as an antiparallel-orientated homodimer. *Virology*. 1998; 250(2):408–14. doi: [10.1006/viro.1998.9389](https://doi.org/10.1006/viro.1998.9389) WOS:000076652100016. PMID: [9792851](https://pubmed.ncbi.nlm.nih.gov/9792851/)
20. Volchkova VA, Klenk HD, Volchkov VE. Delta-peptide is the carboxy-terminal cleavage fragment of the nonstructural small glycoprotein sGP of Ebola virus. *Virology*. 1999; 265(1):164–71. doi: [10.1006/viro.1999.0034](https://doi.org/10.1006/viro.1999.0034) WOS:000084172500016. PMID: [10603327](https://pubmed.ncbi.nlm.nih.gov/10603327/)
21. Shurtleff AC, Biggins JE, Keeney AE, Zumbrun EE, Bloomfield HA, Kuehne A, et al. Standardization of the filovirus plaque assay for use in preclinical studies. *Viruses*. 2012; 4(12):3511–30. doi: [10.3390/v4123511](https://doi.org/10.3390/v4123511) PMID: [23223188](https://pubmed.ncbi.nlm.nih.gov/23223188/); PubMed Central PMCID: PMC3528277.
22. Mehedi M, Hoenen T, Robertson S, Ricklefs S, Dolan MA, Taylor T, et al. Ebola virus RNA editing depends on the primary editing site sequence and an upstream secondary structure. *PLoS Pathog*. 2013; 9(10):e1003677. doi: [10.1371/journal.ppat.1003677](https://doi.org/10.1371/journal.ppat.1003677) PMID: [24146620](https://pubmed.ncbi.nlm.nih.gov/24146620/); PubMed Central PMCID: PMC3798607.
23. Volchkova VA, Dolnik O, Martinez MJ, Reynard O, Volchkov VE. Genomic RNA editing and its impact on Ebola virus adaptation during serial passages in cell culture and infection of guinea pigs. *J Infect Dis*. 2011; 204 Suppl 3:S941–6. Epub 2011/10/19. doi: [10.1093/infdis/jir321](https://doi.org/10.1093/infdis/jir321) PMID: [21987773](https://pubmed.ncbi.nlm.nih.gov/21987773/).
24. Alfson KJ, Avena LE, Beadles MW, Menzie H, Patterson JL, Carrion R Jr., et al. Genetic Changes at the Glycoprotein Editing Site Associated With Serial Passage of Sudan Virus. *J Infect Dis*. 2015; 212 Suppl 2:S295–304. Epub 2015/04/30. doi: [10.1093/infdis/jiv216](https://doi.org/10.1093/infdis/jiv216) PMID: [25920319](https://pubmed.ncbi.nlm.nih.gov/25920319/).
25. Volchkova VA, Dolnik O, Martinez MJ, Reynard O, Volchkov VE. RNA Editing of the GP Gene of Ebola Virus is an Important Pathogenicity Factor. *J Infect Dis*. 2015; 212 Suppl 2:S226–33. Epub 2015/07/04. doi: [10.1093/infdis/jiv309](https://doi.org/10.1093/infdis/jiv309) PMID: [26138826](https://pubmed.ncbi.nlm.nih.gov/26138826/).
26. Trefry JC, Wollen SE, Nasar F, Shamblin JD, Kern SJ, Bearss JJ, et al. Ebola Virus Infections in Nonhuman Primates Are Temporally Influenced by Glycoprotein Poly-U Editing Site Populations in the Exposure Material. *Viruses*. 2015; 7(12):6739–54. doi: [10.3390/v7122969](https://doi.org/10.3390/v7122969) PMID: [26703716](https://pubmed.ncbi.nlm.nih.gov/26703716/); PubMed Central PMCID: PMC4690892.
27. Ladner JT, Beitzel B, Chain PS, Davenport MG, Donaldson EF, Frieman M, et al. Standards for sequencing viral genomes in the era of high-throughput sequencing. *MBio*. 2014; 5(3):e01360–14. doi: [10.1128/mBio.01360-14](https://doi.org/10.1128/mBio.01360-14) PMID: [24939889](https://pubmed.ncbi.nlm.nih.gov/24939889/); PubMed Central PMCID: PMC4068259.
28. Kugelman JR, Kugelman-Tonos J, Ladner JT, Pettit J, Keeton CM, Nagle ER, et al. Emergence of Ebola Virus Escape Variants in Infected Nonhuman Primates Treated with the MB-003 Antibody Cocktail. *Cell Rep*. 2015; 12(12):2111–20. doi: [10.1016/j.celrep.2015.08.038](https://doi.org/10.1016/j.celrep.2015.08.038) PMID: [26365189](https://pubmed.ncbi.nlm.nih.gov/26365189/).