

Open camera or QR reader and
scan code to access this article
and other resources online.



Bariatric Surgery in Patients with Existing Ostomy: A Preliminary Feasibility Study

Ray Portela, MD, Ahmet Vahibe, MD, Joseph N. Badaoui, MD, Omer U.I. Hassan, MD,
Travis J. Mckenzie, MD, Todd A. Kellogg, MD, and Omar M. Ghanem, MD, FACS

Background: Bariatric surgery is the most effective treatment for weight loss and obesity-related comorbidity resolution. However, bariatric surgery is not readily offered in specific populations due to the lack of data assessing its feasibility. This study intends to evaluate bariatric surgery in patients with an existing ostomy.

Methods: We conducted a retrospective case series to assess the safety of Roux-en-Y gastric bypass (RYGB) or sleeve gastrectomy (SG) in patients with an end ileostomy or colostomy. Patient demographics, including obesity-related comorbidities, overall health status (American Society of Anesthesiologists score), and short-term complications (up to 30 days postoperatively), were analyzed.

Results: Six patients were included. The mean age was 58 years, and the mean preoperative body mass index was 41.6. Three patients had a colostomy, and three had an ileostomy. The mean time of ostomy before surgery was 11 years. Two ostomies were due to trauma, two due to inflammatory bowel disease, one due to cancer, and one due to scleroderma. Mean postoperative follow-up was 23 months. No patient had increased ostomy output or infusion center visit. One patient had an ED visit, one had a short-term complication, and one had SG conversion to RYGB.

Conclusions: Bariatric surgery is technically feasible in selected patients with ileostomy/colostomy with a reasonable short-term safety profile.

Keywords: obesity, bariatrics, gastrointestinal, minimally invasive surgery

KEY POINTS

1. Feasible and safe to perform bariatric procedures in selected patients with an ileostomy or colostomy.
2. Patients with preexisting ostomy neither have a higher incidence of dehydration nor have an increased ostomy output after bariatric surgery.

Introduction

THE OBESITY EPIDEMIC affects more than 1 billion people worldwide and is one of the major contributors to health care costs.¹ Bariatric surgery is proven to be the most effective and durable treatment for obesity.^{2,3} Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG) are the most common procedures performed in the United States.⁴⁻⁶ Both RYGB and SG are offered to a wide array of patients

Department of Surgery, Mayo Clinic, Rochester, Minnesota, USA.

© Ray Portela et al., 2022; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons License [CC-BY] (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been updated on May 31, 2022 after first online publication of March 21, 2022 to reflect Open Access, with copyright transferring to the author(s), and a Creative Commons License [CC-BY] (<http://creativecommons.org/licenses/by/4.0>). BARIATRIC SURGICAL PRACTICE AND PATIENT CARE Mary Ann Liebert, Inc. DOI: 10.1089/bari.2021.0131.

since the risk-benefit profile associated with these procedures is usually favorable. However, particular situations may arise where the safety and feasibility of bariatric surgery have not been studied enough. This lack of data in the scientific literature puts the bariatric surgeon in a challenging position if they were ever faced with a particular situation; an example, patients with an existing ostomy who also suffer from obesity.

Are they adequate surgical candidates? Patients with stoma history have undergone previous abdominal surgeries and thus are believed to have higher complication rates.^{6,7} In addition, a certain bowel length is bypassed in RYGB, and there is increased gastric emptying in sleeve patients^{8,9}; two variables that can potentially lead to higher ostomy output and eventually dehydration. However, should this be enough to deprive these patients of a bariatric procedure that can positively impact their overall health and quality of life? Beyond one case report portraying safety, there exist no data on patients undergoing bariatric surgery in the setting of pre-existing ostomy.¹⁰ In this series, we intend to assess the technical feasibility and safety profile of bariatric surgery in patients with a previous colostomy or ileostomy.

Methods

Study design and population

After Institutional Review Board approval, patients from January 1, 2015, to May 31, 2021, with a pre-existing ostomy before bariatric surgery were included in this retrospective case series spanning across the Mayo Clinic Enterprise.

The inclusion criteria were age ≥ 18 years and presence of an ileostomy or colostomy before bariatric surgery. Colostomy was identified using the International Classification of Diseases (ICD) 10 code Z93.3 and ileostomy using the ICD 10 code Z93.2. Not to miss any patients, all stomach and bypass procedures were included and identified using the Current Procedural Terminology (CPT) codes 43621, 43632, 43633, 43644, 43645, 43659, 43774, 43775, 43840, 43846, 43848, 43860, 43887, and 43999. Patients were observed for short- and long-term complications.

Surgical technique

The SG was tailored using a bougie (40-F) starting 4–6 cm proximal to the pylorus and toward the stomach body until about 0.5–1 cm away from the angle of His. For the RYGB, the pouch was 25–30 mL in size, and the Roux limb lengths ranged between 100 and 125 cm. Although the common channel measurement was not documented in all cases, the goal was to maintain a total alimentary limb length (Roux and common channel) of at least 400 cm.

Data collection

Clinical baseline information, including the precursor to ostomy, smoking status, preoperative diabetes mellitus type II, systemic arterial hypertension, dyslipidemia, gastroesophageal reflux disease (GERD), history of myocardial infarction, atrial fibrillation, previous pulmonary embolism, chronic obstructive pulmonary disease, immunosuppression status, and sleep apnea.

Intraoperative notes were screened for intraoperative complications. For short-term complications, all records were screened up to 30 days after surgery, including emergency department visits and infusion center visits. Long-term com-

TABLE 1. DEMOGRAPHICS AND CLINICAL BASELINE

Factor	Overall
<i>N</i>	6
Age, mean (SD)	58 (\pm 11.86)
BMI, mean (SD)	41.59 (\pm 16.34)
Gender: female, <i>n</i> (%)	4 (66.67)
Race, <i>n</i> (%)	
White	5 (83.34)
Other	1 (16.67)
Diabetes mellitus type II	4 (66.67)
Hyperlipidemia	4 (66.67)
Hypertensive requiring medication	4 (66.67)
Current smoker within 1 year	0 (0)
Obstructive sleep apnea	3 (50.00)
Therapeutic anticoagulation	2 (33.34)
Immunosuppression	2 (33.34)
GERD requiring medication	5 (83.34)
ASA class, <i>n</i> (%)	
3	5 (83.34)
4	1 (16.67)
Year of operation, <i>n</i> (%)	
2018	3 (50.00)
2020	1 (16.67)
2021	2 (33.34)
Surgical procedure, <i>n</i> (%)	
Roux-en-Y gastric bypass	2 (33.34)
Sleeve gastrectomy	4 (66.67)

ASA, American Society of Anesthesiologists; BMI, body mass index; GERD, gastroesophageal reflux disease; SD, standard deviation.

plications included reoperation, anastomotic stricture, hernia, marginal ulceration, gastrointestinal leakage, dumping syndrome, short bowel syndrome, and small intestinal bacterial overgrowth. In addition, the nutritional profile and ostomy output were also observed.

Results

Baseline characteristics

Matching the ICD and CPT codes resulted in 59 patients. However, after a thorough chart review, only six patients were found to have an ileostomy or colostomy constructed before bariatric surgery. All cases were operated laparoscopically. Demographics and obesity-related comorbidities are summarized in Table 1. Four patients (66%) were women, the mean

TABLE 2. OSTOMY

Factor	Overall
<i>N</i>	6
Ileostomy, <i>n</i> (%)	3 (50.00)
Colostomy, <i>n</i> (%)	3 (50.00)
Years of ostomy before surgery, mean (SD)	11.67 (\pm 8.98)
Ended ostomy, <i>n</i> (%)	6 (100)
Parastomal hernia, <i>n</i> (%)	4 (66.67)
Cause of ostomy, <i>n</i> (%)	
Trauma	2 (33.34)
IBD	2 (33.34)
Cancer	1 (16.67)
Scleroderma/fecal incontinence	1 (16.67)

IBD, inflammatory bowel disease.

TABLE 3. POSTOPERATIVE FOLLOW-UP

Factor	Overall
<i>N</i>	4
Mean follow-up (SD)	23 months (16.43)
Mean percentage of weight loss (SD)	15.50% (11.77)
Upto 30 days complications, <i>n</i> (%)	
Postoperative ED visit	1 (25.00)
Conversion to open	0 (00)
Visits to an infusion center	0 (00)
Upto 3 years complications, <i>n</i> (%)	
Bariatric revision—SG to RYGB	1 (25.00)
Dumping	1 (25.00)
Anastomotic stricture	1 (25.00)

ED, emergency department; RYGB, Roux-en-Y gastric bypass; SG, sleeve gastrectomy.

age was 58 (\pm 11.86) years, and the mean preoperative body mass index was 41.58 (\pm 16.34) kg/m². SG was performed in four (66%) patients and RYGB in two (33%).

Ostomy

Three patients had a colostomy, and three had an ileostomy, as summarized in Table 2. Mean time from ostomy creation to surgery was 11.67 (\pm 8.98) years. All patients had a closed-ended ostomy. From two patients who underwent RYGB, one had an ileostomy and one a colostomy. Of the SG group, two patients had an ileostomy, and the other two had a colostomy. From four patients with a parastomal hernia, only one had to be addressed intraoperatively at the time of bariatric surgery (primary repair). For the cause of ostomy in the SG, two were due to trauma, one due to Crohn's disease, and the other one due to ulcerative colitis. In the RYGB group, one ostomy was due to colon cancer, and the other one was due to scleroderma leading to rectal prolapse with severe fecal incontinence.

Postoperative follow-up and complications

Table 3 summarizes the postoperative follow-up and complications. The mean follow-up was 23 (\pm 16.43) months. The mean percentage of total weight loss (%TWL) on the follow-up period was 15.50% (\pm 11.77).

Up to 30 days of follow-up, there were no readmissions due to dehydration or outpatient fluid infusions, and only one patient had an emergency department visit for acrochordon removal. Long-term complications included one case of intractable reflux leading to conversion from SG to RYGB. Moreover, one RYGB patient presented with anastomotic stricture and dumping. The stricture was treated with one endoscopic dilation after the surgery and the dumping with dietary counseling. None of the patients had increased postoperative ostomy output, dehydration, or needed fluid infusion. No malnutrition (assessed through routine postoperative bariatrician clinic notes) was noted during the follow-up period.

Discussion

In this report, no short-term complications were encountered in patients with a history of stoma undergoing bariatric surgery. In the long term, one patient had dumping syndrome and an anastomotic stricture managed with dilation, and one

patient underwent revisional surgery. There were no conversions to open surgery or episodes of dehydration requiring medical care. Interestingly, this particular population behaved similarly to patients without a stoma.¹¹

We had expected these patients to have a higher stoma output relative to their baseline, especially in the early postoperative period; however, we found no changes in the stomal output in the short-term evaluation.¹² A possible explanation could include the development of adaptation mechanisms because these patients had long-standing ostomies.¹³ In fact, patients develop hypertrophy and hyperplasia of the remaining intestinal length early after the procedure.¹⁴ Over time, slower intestinal transit leads to better absorption of nutrients,¹⁵ and small bowel electrolyte transport changes lead to a proportional response to fluids.¹⁶

Only a single case report with a patient undergoing RYGB with a pre-existing ostomy had an uneventful 12-month postoperative course.¹⁰ Similarly, our data showed a relatively low rate of adverse events mainly related to the procedure as only one patient eventually required surgical revision. Despite both procedures being safe and effective,^{4,5} careful decision-making to avoid multiple surgeries is more relevant to this population. In our report, a patient with mild self-reported GERD symptoms underwent SG. One of the factors affecting the decision-making process was the lack of data about the feasibility of RYGB in this particular population. This patient ended up requiring conversion to RYGB due to the severity of the reflux, and had an uneventful postoperative course, and witnessed amelioration in his reflux symptoms.^{17–20}

Obesity is related to an increase in the incidence of parastomal hernias.^{21,22} In addition, recurrence rates are higher after hernia repair in patients with obesity.^{23,24} In our report, four patients had parastomal hernias. Our approach to parastomal hernias is similar to our approach to any ventral hernia at the time of bariatric surgery. We attempt to leave the hernia content within the hernia sac if possible and deal with the hernia repair later once weight loss is achieved. However, if hernia contents were reduced, primary closure of the defect should be performed to prevent bowel incarceration postoperatively, thus leading to obstruction and potentially proximal leak.

This report has several limitations. Initially, the small sample size greatly reduces the chances of generalizability of our results. In addition, the lack of a control group introduces several biases. Finally, the relatively short follow-up with a mean follow-up of 23 months is a limitation. The fundamentally different underlying diagnosis that leads to ostomy also introduces biases. Further investigations with bigger numbers, stratified analysis, and a longer follow-up are required to better understand the postoperative behaviors pertaining to this specific population.

Conclusion

Bariatric surgery in the setting of pre-existing ostomy is technically feasible with a reasonable short-term safety profile. Further investigations with bigger numbers and a longer follow-up are required to better understand the postoperative behaviors pertaining to this specific population.

Author Disclosure Statement

No competing financial interests exist.

Funding Information

No specific funding was received for this project.

References

- Lee MJ, Kwon JS, Jiang HB, Choi EH, Park G, Kim KM. The antibacterial effect of non-thermal atmospheric pressure plasma treatment of titanium surfaces according to the bacterial wall structure. *Sci Rep* 2019;9:1–13.
- Maciejewski ML, Arterburn DE, van Scoyoc L, Smith VA, Yancy WS, Weidenbacher HJ, *et al.* Bariatric surgery and long-term durability of weight loss. *JAMA Surg* 2016;151:1046–1055.
- Appel LJ, Clark JM, Yeh HC, Wang NY, Coughlin JW, Daumit G, *et al.* Comparative effectiveness of weight-loss interventions in clinical practice. *N Engl J Med* 2011;365:1959–1968.
- Christou NV, Sampalis JS, Liberman M, Liberman M, Look D, Auger S, *et al.* Surgery decreases long-term mortality, morbidity, and health care use in morbidly obese patients. *Ann Surg* 2004;240:416–424.
- Schauer DP, Arterburn DE, Livingston EH, Fischer D, Eckman MH. Decision modeling to estimate the impact of gastric bypass surgery on life expectancy for the treatment of morbid obesity. *Arch Surg* 2010;145:57–62.
- Stahl JM, Malhotra S. Obesity surgery indications and contraindications. *StatPearls* 2019. Available at www.ncbi.nlm.nih.gov/pubmed/30020657. Accessed January 12, 2022.
- Tevis SE, Cobian AG, Truong HP, Craven MW, Kennedy GD. Implications of multiple complications on the post-operative recovery of general surgery patients. *Ann Surg* 2016;263:1213–1218.
- Ivanics T, Nasser H, Leonard-Murali S, Genaw J. Dehydration risk factors and impact after bariatric surgery: an analysis using a national database. *Surg Obes Relat Dis* 2019;15:2066–2074.
- Vavricka SR, Greuter T. Gastroparesis and dumping syndrome: current concepts and management. *J Clin Med* 2019;8:1127.
- Khitaryan A, Miziev I, Mezhunts A, Look D, Auger S, McLean APH, *et al.* Roux-en-Y gastric bypass and parastomal hernia repair: case report of concurrent operation in comorbid patient. *Int J Surg Case Rep* 2020;71:360–363.
- Pomp A. Safety of bariatric surgery. *Lancet Diabetes Endocrinol* 2014;2:98–100.
- Kwiatt M, Kawata M. Avoidance and management of stomal complications. *Clin Colon Rect Surg* 2013;26:112–121.
- Rowe KM, Schiller LR. Ileostomy diarrhea: pathophysiology and management. *Baylor Univ Med Center Proc* 2020;33:218–226.
- Wright HK, Cleveland JC, Tilson MD, Herskovic T. Morphology and absorptive capacity of the ileum after ileostomy in man. *Am J Surg* 1969;117:242–245.
- Ladas SD, Isaacs PET, Murphy GM, Sladen GE. Fasting and postprandial ileal function in adapted ileostomates and normal subjects. *Gut* 1986;27:906–912.
- Huber FX, Lucas M, Stern J, Hinz U, Haack D, Heuschen U, *et al.* Changes in glucocorticoid and mineralocorticoid hormone levels due to compensation for ileostomy losses. *Int J Surg Invest* 2001;2:369–375.
- Lager CJ, Esfandiari NH, Subauste AR, Kraftson AT, Brown MB, Cassidy RB, *et al.* Roux-en-Y gastric bypass vs. sleeve gastrectomy: balancing the risks of surgery with the benefits of weight loss. *Obes Surg* 2017;27:154–161.
- Peterli R, Wolnerhanssen BK, Peters T, Vetter D, Kroll D, Borbely Y, *et al.* Effect of laparoscopic sleeve gastrectomy vs laparoscopic roux-en-y gastric bypass on weight loss in patients with morbid obesity the sm-boss randomized clinical trial. *JAMA* 2018;319:255–265.
- Parmar CD, Mahawar KK, Boyle M, Schroeder N, Balupuri S, Small PK. Conversion of sleeve gastrectomy to Roux-en-Y Gastric bypass is effective for gastro-oesophageal reflux disease but not for further weight loss. *Obes Surg* 2017;27:1651–1658.
- Raj PP, Bhattacharya S, Misra S, Kumar SS, Khan MJ, Gunasekaran SC, *et al.* Gastroesophageal reflux-related physiologic changes after sleeve gastrectomy and Roux-en-Y gastric bypass: a prospective comparative study. *Surg Obes Relat Dis* 2019;15:1261–1269.
- Sohn YJ, Moon SM, Shin US, Jee SH. Incidence and risk factors of parastomal hernia. *J Korean Soc Coloproctol* 2012;28:241–246.
- Tivenius M, Näsval P, Sandblom G. Parastomal hernias causing symptoms or requiring surgical repair after colorectal cancer surgery—a national population-based cohort study. *Int J Colorectal Dis* 2019;34:1267–1272.
- Nieto E, Ponz C, Ananin S, Vazquez E. Obesity as a risk factor for complications and recurrences after ventral hernia repair. *Int Jo Abdom Wall Hernia Surg* 2020;3:1.
- Tsereteli Z, Pryor BA, Heniford BT, Park A, Voeller G, Ramshaw BJ. Laparoscopic ventral hernia repair (LVHR) in morbidly obese patients. *Hernia* 2008;12:233–238.

Address correspondence to:
 Omar M. Ghanem, MD, FACS
 Department of Surgery
 Mayo Clinic
 200 1st St SW
 Rochester, MN 55905
 USA

E-mail: ghanem.omar@mayo.edu