

RESEARCH ARTICLE

Open Access

Electron transport in acetate-grown *Methanosarcina acetivorans*

Mingyu Wang¹, Jean-Francois Tomb² and James G Ferry^{1*}

Abstract

Background: Acetate is the major source of methane in nature. The majority of investigations have focused on acetotrophic methanogens for which energy-conserving electron transport is dependent on the production and consumption of H₂ as an intermediate, although the great majority of acetotrophs are unable to metabolize H₂. The presence of cytochrome *c* and a complex (Ma-Rnf) homologous to the Rnf (*R*hodobacter *n*itrogen *f*ixation) complexes distributed in the domain *Bacteria* distinguishes non-H₂-utilizing *Methanosarcina acetivorans* from H₂-utilizing species suggesting fundamentally different electron transport pathways. Thus, the membrane-bound electron transport chain of acetate-grown *M. acetivorans* was investigated to advance a more complete understanding of acetotrophic methanogens.

Results: A component of the CO dehydrogenase/acetyl-CoA synthase (CdhAE) was partially purified and shown to reduce a ferredoxin purified using an assay coupling reduction of the ferredoxin to oxidation of CdhAE. Mass spectrometry analysis of the ferredoxin identified the encoding gene among annotations for nine ferredoxins encoded in the genome. Reduction of purified membranes from acetate-grown cells with ferredoxin lead to reduction of membrane-associated multi-heme cytochrome *c* that was re-oxidized by the addition of either the heterodisulfide of coenzyme M and coenzyme B (CoM-S-S-CoB) or 2-hydroxyphenazine, the soluble analog of methanophenazine (MP). Reduced 2-hydroxyphenazine was re-oxidized by membranes that was dependent on addition of CoM-S-S-CoB. A genomic analysis of *Methanosarcina thermophila*, a non-H₂-utilizing acetotrophic methanogen, identified genes homologous to cytochrome *c* and the Ma-Rnf complex of *M. acetivorans*.

Conclusions: The results support roles for ferredoxin, cytochrome *c* and MP in the energy-conserving electron transport pathway of non-H₂-utilizing acetotrophic methanogens. This is the first report of involvement of a cytochrome *c* in acetotrophic methanogenesis. The results suggest that diverse acetotrophic *Methanosarcina* species have evolved diverse membrane-bound electron transport pathways leading from ferredoxin and culminating with MP donating electrons to the heterodisulfide reductase (HdrDE) for reduction of CoM-S-S-CoB.

Background

The decomposition of complex organic matter to methane (biomethanation) in diverse anaerobic habitats of Earth's biosphere involves an anaerobic microbial food chain comprised of distinct metabolic groups, the first of which metabolizes the complex organic matter primarily to acetate and also formate or H₂ that are growth substrates for two distinct methane-producing groups (methanogens) [1]. The methyl group of acetate

contributes most of the methane produced in the biomethanation process *via* the aceticlastic pathway whereas the remainder originates primarily from the reduction of CO₂ with electrons derived from the oxidation of formate or H₂ in the CO₂-reduction pathway [2,3]. Smaller, albeit significant, amounts of methane derive from the methyl groups of methanol, methylamines and dimethylsulfide [1].

Only two genera of aceticlastic methanogens have been described, *Methanosarcina* and *Methanosaeta* [2]. In both genera, the CO dehydrogenase/acetyl-CoA complex (Cdh) cleaves activated acetate into methyl and carbonyl groups. The methyl group is transferred to coenzyme M (HS-CoM) producing CH₃-S-CoM that is

* Correspondence: jgf3@psu.edu

¹Department of Biochemistry and Molecular Biology, Eberly College of Science, The Pennsylvania State University, University Park, Pennsylvania 16802-4500, USA

Full list of author information is available at the end of the article

reductively demethylated to methane with electrons donated by coenzyme B (HS-CoB). The heterodisulfide CoM-S-S-CoB is a product of the demethylation reaction that is reduced to the sulfhydryl forms of the cofactors by heterodisulfide reductase (Hdr). The proton gradient driving ATP synthesis is generated *via* a membrane-bound electron transport chain originating with oxidation of the carbonyl group of acetate by Cdh and terminating with reduction of CoM-S-S-CoB by Hdr. Although the pathway of carbon flow from the methyl group of acetate to methane is understood for both acetoclastic genera, the understanding of electron transport coupled to generation of the proton gradient is incomplete. The majority of investigations have focused on *Methanosarcina barkeri* and *Methanosarcina mazei* for which electron transport is dependent on the production and consumption of H₂ as an intermediate, although the great majority of *Methanosarcina* species [4] and all *Methanosaeta* species are unable to metabolize H₂.

In the H₂-metabolizing *Methanosarcina* species investigated, a ferredoxin accepts electrons from Cdh [5,6] and donates to a membrane-bound Ech hydrogenase complex that produces H₂ and generates a proton gradient for ATP synthesis [7-9]. A hypothesis has been advanced wherein H₂ is re-oxidized by another membrane-bound hydrogenase (Vho) that transfers electrons to methanophenazine (MP), a quinone-like electron carrier [9]. In the model, MP donates electrons to the heterodisulfide reductase HdrDE accompanied by translocation of protons which further contributes to ATP synthesis.

An electron transport chain has been hypothesized for the marine isolate *Methanosarcina acetivorans*, the only non-H₂-metabolizing acetotrophic methanogen for which the genome is sequenced. Although encoding Cdh, the genome does not encode Ech hydrogenase [10,11]. Furthermore, in contrast to all H₂-utilizing acetoclastic *Methanosarcina* species investigated [12], acetate-grown *M. acetivorans* synthesizes a six-subunit complex (Ma-Rnf) [13] encoded within a co-transcribed eight-gene (MA0658-0665) cluster with high identity to membrane-bound Rnf (**R**hodobacter **n**itrogen **f**ixation) complexes from the domain *Bacteria*. It is hypothesized that the Ma-Rnf complex plays an essential role in the electron transport chain, generating a sodium gradient that is exchanged for a proton gradient driving ATP synthesis [13]. Consistent with this idea, it was recently shown that the six-subunit Rnf complex from *Acetobacterium woodii* of the domain *Bacteria* couples electron transport from reduced ferredoxin to NAD⁺ with the generation of a sodium gradient [14]. Remarkably, the Ma-Rnf complex of *M. acetivorans* is co-transcribed with a gene (MA0658) encoding a multi-heme

cytochrome *c*, and another flanking gene (MA0665) encoding a hypothetical membrane integral protein with unknown function [13]. Indeed, the cytochrome *c* was shown to be synthesized in high levels of acetate-grown cells where it completely dominates the UV-visible spectrum of the purified membranes and is distinguishable from *b*-type cytochromes [13]. Furthermore, it was recently reported (A. M. Guss and W. W. Metcalf, unpublished results) that a six-subunit Ma-Rnf/cytochrome *c* (Δ MA0658-0665) deletion mutant of *M. acetivorans* fails to grow with acetate [15]. However, biochemical evidence necessary to support the hypothesized role of cytochrome *c* has not been forthcoming. The only other report of cytochromes *c* in methanogens is for the H₂-metabolizing species *Methanosarcina mazei* (*f. Methanosarcina* strain Gö1) grown with methanol [16].

The freshwater isolate *Methanosarcina thermophila* is the only non-H₂-metabolizing acetotrophic methanogen for which electron transport components have been investigated biochemically [17]. Like H₂-metabolizing *Methanosarcina* species, ferredoxin mediates electron transfer between Cdh and the membrane-bound electron transport chain in which a cytochrome *b* participates and dominates the UV-visible absorbance spectrum of membranes. It is also reported that MP is the electron donor to HdrDE [18]. Electron carriers other than cytochrome *b* that participate between ferredoxin and MP were not identified. Importantly, no evidence for participation of Ma-Rnf or cytochrome *c* was reported. Homologs encoding an Ma-Rnf complex and cytochrome *c* are absent in the sequenced genome of *Methanosaeta thermophila* suggesting yet another novel electron transport chain that functions in the conversion of acetate to methane in this non-H₂-metabolizing genus [19]. Clearly, diverse electron transport pathways have evolved in diverse acetotrophic methanogens necessitating biochemical investigations of representative species.

The absence of Ech hydrogenase and the demonstrated presence of the Ma-Rnf complex and cytochrome *c* that is elevated in acetate- *versus* methanol-grown cells [13] suggests that electron transport of the non-H₂-metabolizing marine isolate *M. acetivorans* is decidedly dissimilar from the genus *Methanosaeta* and H₂-metabolizing acetotrophic species of the genus *Methanosarcina*. However, a biochemical investigation essential to support the role of electron carriers has not been reported for *M. acetivorans*. Here we report evidence indicating roles for ferredoxin, cytochrome *c* and MP in electron transport of acetate-grown *M. acetivorans*. The results underscore the diversity of electron transport pathways in acetotrophic methanogens and contribute to a more complete understanding of acetotrophic methanogenesis.

Results

The electron acceptor for the CO dehydrogenase/acetyl-CoA complex of *M. acetivorans*

The Cdh from acetate-grown *M. acetivorans* was purified to ascertain the electron acceptor that initiates electron transport. The Cdh complex purified from the H₂-metabolizing acetotrophic species *Methanosarcina barkeri* contains five-subunits (CdhABCDE) [20] of which the CdhAE component oxidizes CO derived from the carbonyl group of acetate [21]. The genome of *M. acetivorans* is annotated with duplicate Cdh gene clusters [10], each encoding five subunits homologous to the Cdh subunits of *M. barkeri*. Previous proteomic analyses of acetate-grown *M. acetivorans* identified subunits CdhA, CdhB and CdhC from one cluster (MA1011-16) and CdhA, CdhB CdhC and CdhE from the other (MA3860-65) [22]. The purification was monitored by following the CO-dependent reduction of methyl viologen. SDS PAGE of the purified enzyme showed bands with molecular masses of 16 kDa and 85 kDa consistent with the predicted values for the CdhA and CdhE subunits encoded in the genome. Mass spectrometry of the protein bands identified the CdhA and CdhE subunits encoded by both Cdh gene clusters consistent with previous proteomic analyses that indicated up-regulation of both clusters in acetate- versus methanol-grown cells [22].

Ferredoxin from acetate-grown cells of *M. acetivorans* was purified as described in the Methods section to determine if it accepts electrons from the partially purified CdhAE components thereby initiating electron transport. Mass spectrometry analysis of the purified ferredoxin detected only one protein identified as the product of MA0431 previously annotated as a 2 × [4Fe-4S] ferredoxin [23]. The UV-visible absorption spectrum of the as-purified protein was typical of ferredoxins with an absorption maximum at 402 nm that decreased upon reduction with dithionite (Additional file 1, Figure S1). The genome of *M. acetivorans* is annotated with nine genes encoding ferredoxins, a phylogenetic analysis of which is shown in Additional file 2, Figure S2. The analysis revealed that the product of MA0431 is closely related to the 2 × [4Fe-4S] ferredoxin purified from acetate-grown cells of *M. thermophila* [24-27] and the ferredoxin up-regulated in acetate- versus methanol-grown *M. mazei* [28]. These three ferredoxins contain two CX₂CX₂CX₃CP motifs typical of 2 × [4Fe-4S] ferredoxins and share high identity within a distinct clade (Additional file 2, Figure S2). Figure 1 shows CO-dependent reduction of the purified *M. acetivorans* ferredoxin catalyzed by the CdhAE components purified from *M. acetivorans*. These results suggest that ferredoxin isolated initiates the electron transport chain in both *M.*





Figure 1 Reduction of ferredoxin by CdhAE. The 70- μ l reaction mixture consisted of 2.2 μ g of CdhAE and 28 μ M (final concentration) of ferredoxin contained in 50 mM MOPS buffer (pH 6.8) under 1 atm CO. The reaction was initiated with CdhAE. A, complete reaction mixture initial absorbance 0.61. B, reaction mixture minus CdhAE, initial absorbance 0.72. C, reaction mixture minus ferredoxin, initial absorbance 0.72. The reduction of ferredoxin was followed by the decrease in absorbance at 402 nm.

acetivorans and H₂-metabolizing acetotrophic *Methanosarcina* species.

Ferredoxin as the electron donor to the membrane-bound electron transport chain

The finding that ferredoxin is an electron acceptor for the CdhAE component of the Cdh complex of *M. acetivorans* raises the question whether it is the direct electron donor to membrane-bound electron carriers or if other soluble electron carriers are required to mediate electron transfer between ferredoxin and the membrane. This question was addressed in a system containing sucrose gradient-purified membranes and plant ferredoxin-NADPH reductase (FNR) to regenerate reduced ferredoxin that was purified from acetate-grown cells. The CO-dependent reduction of ferredoxin with CdhAE was not used to avoid binding of CO to high spin heme in cytochrome *c* and potentially inhibiting membrane-bound electron transport. The NADPH:CoM-S-S-CoB oxidoreductase activity was monitored by detecting the sulfhydryl groups of HS-CoM and HS-CoB (Figure 2). No significant activity was detected when each component of the reaction mixture was deleted individually including membranes. The dependence of the activity on purified membranes and the concentration of ferredoxin purified from acetate-grown *M. acetivorans* indicated a role for the ferredoxin in the direct transfer of electrons from CdhAE to the membrane-bound electron transport chain terminating with reduction of CoM-S-S-CoB by heterodisulfide reductase.




Role of cytochrome c in the membrane-bound electron transport chain

It was previously documented [13] that purified membranes of acetate-grown *M. acetivorans* contain a multi-heme cytochrome *c* that clearly dominates the UV-visible spectrum of membranes from acetate-grown *M. acetivorans* with the major peak centered at 554 nm (Figure 3B). Absorbance at 554 nm increased on incubation of the membrane fraction with the reduced ferredoxin regenerating system indicating reduction of cytochrome *c* that was dependent on ferredoxin (Figure 3). Addition of CoM-S-S-CoB oxidized the reduced cytochrome (Figure 4) indicating that it is a component of the membrane-bound electron transport chain terminating with reduction of the heterodisulfide. The re-oxidation was too rapid to determine a rate and incomplete, albeit greater than 50%. The explanation for incomplete re-oxidation is unknown, although the result is nearly identical to that reported for the re-oxidation of cytochromes in the membrane fraction of methanol-grown *M. mazei* that was rapid and reached 40% re-oxidation [16]. This is the first report of cytochrome *c* involvement in the conversion of acetate to methane.

Role of methanophenazine in the membrane-bound electron transport chain

The soluble analog of MP, 2-hydroxyphenazine, has been used to investigate the role of MP in methanogens



[18,29]. The 2-hydroxyphenazine that was reduced in the presence of membranes with the CO/CdhAE ferredoxin regenerating system was re-oxidized upon addition of CoM-S-S-CoB (Figure 5) consistent with a membrane-bound heterodisulfide reductase. The re-oxidation generated a total of 239 μ M free thiol groups in this representative experiment, a result that is in approximate agreement with the observed oxidation of 106 μ M 2-hydroxyphenazine. Assuming a two-electron




Figure 4 Oxidation of membrane-bound cytochrome *c* by CoM-S-S-CoB. The reduction of cytochrome *c* was performed as described in the caption to Figure 3. The 100- μ l reaction mixture consisted of membranes (400 μ g protein), 2 μ M ferredoxin, 1 μ g FNR and 1 mM NADPH. FNR was added at time zero and 0.32 mM (final concentration) CoM-S-S-CoB was added (arrow). Reduction and oxidation of cytochrome *c* was monitored by the absorbance at 554 nm.

transfer from the MP analog, 212 μ M free thiol groups would be expected. These results indicate that MP is a component of the membrane-bound electron transport chain terminating with reduction of CoM-S-S-CoB.




Figure 5 Reduction of 2-hydroxyphenazine and re-oxidation dependent on membranes and CoM-S-S-CoB. The 100- μ l reaction mixture consisted of membranes (107 μ g protein), 4 μ M ferredoxin, 100 μ M 2-hydroxyphenazine and CdhAE (40 μ g) in 50 mM MOPS (pH 6.8) under 1 atm CO. The reduction and oxidation of 2-hydroxyphenazine was followed by the absorbance at 475 nm ($\epsilon_{475} = 2.5 \text{ mM}^{-1} \text{ cm}^{-1}$). CdhAE was added to initiate the reduction at time zero. At point A the cuvette was flushed with 100% N₂ and 2 μ l of MOPS buffer (pH 6.8) was added. At points B and C, 2 μ l of MOPS buffer (pH 6.8) containing CoM-S-S-CoB was added to the reaction reaching final concentrations of 240 and 480 μ M.

The results implicating MP and cytochrome *c* in the membrane-bound electron transport chain presents the possibility of electron transfer between these carriers. The MP analog 2-hydroxyphenazine re-oxidized cytochrome *c* when added to membranes of acetate-grown cells previously reduced with ferredoxin (Figure 6). These results suggest that MP is either directly or indirectly linked to cytochrome *c*, a result further supporting the participation of MP and cytochrome *c* in the membrane-bound electron transport chain.

Discussion

The overwhelming majority of methanogens capable of growth *via* conversion of the methyl group of acetate to




Figure 6 Oxidation of membrane-bound cytochrome *c* by 2-hydroxyphenazine. The 100- μ l reaction mixture consisted of membranes (750 μ g protein), 4 μ M ferredoxin 1 mM NADPH and 1 μ g FNR contained in 50 mM MOPS buffer (pH 6.8). The reduction of cytochrome *c* was initiated by addition of FNR. The reduction and re-oxidation was monitored at 554 nm. When fully reduced, 200 μ M 2-hydroxyphenazine (2 μ l) was added (arrow). Panel A, time course for the reduction and re-oxidation by 2-hydroxyphenazine added at the arrow. Panel B, reduced minus oxidized UV-visible spectra of membranes before (lower trace) and after (upper trace) addition of 2-hydroxyphenazine.

methane do not metabolize H₂ suggesting they employ an electron transport pathway distinct from that proposed for the few acetotrophic methanogens in which H₂ is an obligatory intermediate. *M. acetivorans* is the ideal candidate to represent the majority of acetotrophic *Methanosarcina* species by virtue of its sequenced genome and published proteomic analyses leading to the hypothesis of a novel electron transport pathway for acetotrophic methanogens incapable of metabolizing H₂. This, the first biochemical investigation of electron transport in *M. acetivorans*, has established roles for electron carriers that reveal both commonalities and differences in electron transport pathways of diverse acetotrophic *Methanosarcina* species.

Figure 7 compares the current understanding of electron transport for acetate-grown *M. acetivorans* with that for H₂-metabolizing acetotrophic *Methanosarcina* species. In both pathways, the five-subunit CdhABCDE complex (not shown) cleaves the C-C and C-S bonds of acetyl-CoA releasing a methyl group and CO that is oxidized to CO₂ with electrons transferred to ferredoxin. The CdhAE component of *M. acetivorans* was isolated independently from the other subunits and both copies encoded in the genome were represented. Although it was not possible to determine which CdhAE component reduced ferredoxin, the high percent identities (CdhA, MA1016 vs. MA3860 = 84% and CdhE, MA1015 vs. MA3861 = 82%) suggests it is the electron acceptor for

either or both copies. In both pathways, ferredoxin is the electron donor to a membrane-bound electron transport chain that terminates with MP donating electrons to the heterodisulfide reductase HdrDE that catalyzes the reduction of CoB-S-S-CoM. Proteomic and genetic evidence [15,22] indicates that HdrDE functions in acetate-grown *M. acetivorans*. MP is the direct electron donor to HdrDE in acetate-grown cells of H₂-metabolizing *Methanosarcina* species and the non-H₂-metabolizing *M. thermophila* [18]. Thus, it is reasonable to postulate that MP is also the direct electron donor to HdrDE of *M. acetivorans*. However, the electron transport pathways of H₂-metabolizing and non-H₂-metabolizing species diverge significantly in electron transfer between ferredoxin and MP. In H₂-metabolizing species, ferredoxin donates electrons to the membrane-bound Ech hydrogenase. A H₂ cycling mechanism is postulated in which the H₂ generated by Ech hydrogenase is re-oxidized by the MP-reducing Vho-type hydrogenase further contributing to the proton gradient [8]. Although the genome of *M. acetivorans* contains homologs of genes encoding Vho-type hydrogenases they are not expressed during growth with acetate [4], a result consistent with the absence of Ech hydrogenase and inability to metabolize H₂. Instead, the results reported here support a role for cytochrome *c* mediating electron transport between ferredoxin and MP, although the identities of the direct electron donor and acceptor for cytochrome *c* remain




Figure 7 Comparison of electron transport pathways for *Methanosarcina mazei* and *Methanosarcina barkeri* versus *Methanosarcina acetivorans*. Panel A, *M. mazei* and *M. barkeri*. Panel B, *M. acetivorans*. Ech, Ech hydrogenase; Fd_r, ferredoxin reduced; Fd_o, ferredoxin oxidized; Vho, Vho hydrogenase; MP, methanophenazine; HdrDE, heterodisulfide reductase; CoM-SH, coenzyme M; CoB-SH, coenzyme B; Atp, ATP synthase; Cyt *c*, cytochrome *c*; Ma-Rnf, Rnf complex from *M. acetivorans*; Mrp, putative sodium/proton antiporter.

unknown. The membrane location of cytochrome *c* is unknown; however, if on the outer aspect as for multi-heme cytochromes *c* in the domain *Bacteria*, ferredoxin would be an unlikely electron donor. The most probable electron donor to cytochrome *c* is the Ma-Rnf complex that is also hypothesized to accept electrons from ferredoxin in analogy to homologous Rnf complexes from the domain *Bacteria* [13,30]. In the absence of biochemical evidence, the proposed role for Ma-Rnf in electron transport is at least consistent with up-regulation in acetate *vs.* methanol-grown cells and the reported failure of an Ma-Rnf-cytochrome *c* deletion mutant (Δ MA0658-0665) of *M. acetivorans* to grow with acetate [15]. The proposed interaction of Ma-Rnf with cytochrome *c* is supported by co-transcription of the encoding genes and up-regulation in acetate- *vs.* methanol-grown cells [13]. A role for cytochrome *c* in the electron transport chain is also supported by results showing re-oxidation of cytochrome *c* upon addition of the MP analog 2-hydroxyphenazine to ferredoxin-reduced membranes, although an unknown carrier mediating electron transfer between cytochrome *c* and MP cannot be ruled out.

It was recently shown that the Rnf complex from *A. woodii* translocates sodium ions coupled to electron transfer from ferredoxin to NAD⁺ [14]. In view of the potential sodium ion pumping function of Ma-Rnf, it is interesting to note that a multi-subunit sodium/proton antiporter (Mrp) is up-regulated in acetate-grown *M. acetivorans* and that the encoding genes are absent in H₂-metabolizing *Methanosarcina* species [13]. Thus, it is tempting to speculate that Ma-Rnf generates a sodium gradient (high outside) that is exchanged for a proton gradient by Mrp. The only other coupling site is the reduction and oxidation of MP generating a proton gradient as proposed for H₂-metabolizing *Methanosarcina* species (Figure 7). The role of a proton gradient driving ATP synthesis is consistent with the presence of a proton translocating ATP synthase in acetate-grown cells [13] recently shown to be the primary ATP synthase [31].

The available evidence indicates that the non-H₂-metabolizing freshwater isolate *M. thermophila* also utilizes ferredoxin as electron donor to a membrane-bound electron transport chain involving cytochrome *b* and culminating with MP donating electrons to HdrDE [17,18,32]; however, a role for cytochrome *c* is not evident and other electron carriers have not been reported. Thus, based on current evidence, it appears that all acetotrophic *Methanosarcina* species have in common ferredoxin as electron donor to a membrane-bound electron transport chain terminating with MP donating electrons to HdrDE, although differ widely in membrane components transferring electrons from ferredoxin to MP. The evidence for involvement of HdrDE in acetate-grown

cells is convincing; however, genes (MA2868, MA4236 and MA4237) homologous to those encoding the soluble HdrABC heterodisulfide reductase of CO₂-reducing methanogens were shown to be up regulated in acetate-*versus* methanol-grown *M. acetivorans* [33]. This result is consistent with the previously reported increased abundance of HdrA encoded by MA2868 in acetate-*versus* methanol-grown *M. acetivorans* [22] which opens the possibility that the electron transport chain may terminate with both the membrane HdrDE or a soluble HdrABC heterodisulfide reductase.

Of the nine putative 2 × [4Fe-4S] ferredoxins annotated for the genome of *M. acetivorans*, only the ferredoxin encoded by MA0431 was purified from acetate-grown cells. While it cannot be ruled out that other ferredoxins are synthesized in acetate-grown cells, the results suggest that the ferredoxin encoded by MA0431 is at least dominant in acetate-grown cells. Of the nine putative 2 × [4Fe-4S] ferredoxins, the one purified from *M. acetivorans* is most closely related to that isolated from acetate-grown *M. thermophila* [26], a result suggesting it is the preferred electron acceptor of CdhAE in acetate-grown *Methanosarcina* species.

Interestingly, genes encoding subunits of Ma-Rnf or Ech hydrogenase are absent in the genome of the acetate-utilizing isolate *Methanosaeta thermophila* [19] that is also incapable of metabolizing H₂ suggesting still other alternative electron transport pathways coupled to generation of ion gradients driving ATP synthesis in acetate-utilizing methanogens. The physiological significance of these diverse electron transport pathways is yet to be determined; however, it has been suggested that avoiding H₂ is advantageous to the marine isolate *M. acetivorans* since sulfate reducing species that dominate this environment outcompete methanogens for H₂ potentially disrupting electron transport [13]. It is important to note here that although *M. acetivorans* is incapable of growth with H₂/CO₂ it synthesizes all of the enzymes necessary for reduction of CO₂ to methane and is capable of robust growth *via* the CO₂-reduction pathway albeit with electrons derived from the oxidation of CO [34-36].

Comparative analysis of the *M. thermophila* genome

M. thermophila is an acetotrophic *Methanosarcina* species incapable of metabolizing H₂ [37,38]. Analysis of the genomic sequence revealed a gene cluster identical in arrangement and homologous to genes encoding the six subunits of Ma-Rnf and multi-heme cytochrome *c* of *M. acetivorans* with deduced sequence identities ranging from 86 to 98% (Additional file 3, Figure S3A). Alignments of the deduced sequences showed strict conservation of heme-binding, flavin binding and iron-sulfur binding motifs suggesting conserved functions

(Additional file 3, Figure S3B). Although not conclusive, these results are consistent with a role for the Ma-Rnf complex and multi-heme cytochrome *c* in the electron transport pathway of *M. thermophila* grown with acetate. Furthermore, the genome of *M. thermophila* contains a gene cluster (Additional file 4, Figure S4) homologous to genes encoding the seven subunits of the sodium/proton antiporter (Mrp) that is up-regulated in acetate- versus methanol-grown cells of *M. acetivorans* and absent in the sequenced genomes of acetotrophic *Methanosarcina* species capable of metabolizing H₂/CO₂ [22,39].

Conclusions

Although the majority of *Methanosarcina* species are unable to metabolize H₂, electron transport has only been investigated in the few species for which H₂ is an obligatory intermediate. *M. acetivorans* is proposed to utilize a fundamentally different electron transport pathway based on bio-informatic, proteomic and genetic approaches. However, the proposal has not been tested biochemically. The results indicate roles for ferredoxin, cytochrome *c* and MP in support of the proposed electron transport pathway. Further, this is the first report for involvement of a cytochrome *c* in acetotrophic methanogens. The results suggest that diverse acetotrophic *Methanosarcina* species have evolved diverse membrane-bound electron transport pathways leading from ferredoxin and culminating with MP donating electrons to HdrDE for reduction of CoM-S-S-CoB.

Methods

Materials

CoM-S-S-CoB was a kind gift of Dr. Jan Keltjens. 2-hydroxyphenazine was custom synthesized by Sigma-Aldrich (St. Louis, MO). All other chemicals were purchased from Sigma-Aldrich or VWR International (West Chester, PA). All chromatography columns, resins and pre-packed columns were purchased from GE Healthcare (Waukesha, WI).

Preparation of cell extract and membranes

M. acetivorans [40] was cultured with acetate as described previously [41] and the cell paste was frozen at -80°C. All solutions were O₂-free and manipulations were performed anaerobically in an anaerobic chamber (Coy Manufacturing, Ann Arbor, MI) containing 95% N₂ and 5% H₂. Frozen cells were thawed, re-suspended (1 g wet weight/ml buffer) in 50 mM MOPS buffer (pH 6.8) containing 10% (v/v) ethylene glycol and passed twice through a French pressure cell at 6.9×10^3 kPa. The lysate was centrifuged at $7,200 \times g$ for 15 min to pellet cell debris and unbroken cells. Membranes were purified from the cell extract using a discontinuous

sucrose gradient comprised of 2 ml 70% sucrose, 4 ml 30% sucrose and 1.5 ml 20% sucrose contained in 50 mM MOPS buffer (pH 6.8). A 2 ml volume of cell extract was overlaid on the gradient and centrifuged at $200,000 \times g$ for 2 h in a Beckman type 50 Ti rotor. The brown band containing membranes at the 30% and 70% sucrose interface was collected and stored at -80°C until use.

Purification of the α : component (CdhAE) of the CO dehydrogenase/acetyl-CoA synthase complex

All purification steps and biochemical assays were performed anaerobically in the anaerobic chamber. Crude cell extract of acetate-grown *M. acetivorans* was centrifuged at $200,000 \times g$ for 2 h to pellet the membrane fraction. The supernatant solution (200 mg of protein in 10-ml) containing the soluble fraction was loaded onto a Q-Sepharose FF column (50 ml bed volume) equilibrated with 50 mM MOPS (pH 6.8). The column was developed with 500 ml of a 0-1.0 M NaCl linear gradient. Each 10 ml fraction was assayed for CO dehydrogenase activity by monitoring the CO-dependent reduction of methyl viologen as previously described [42]. The pooled fractions from the peak with the highest specific activity were concentrated 10-fold with a Vivacell 70 protein concentrator equipped with a 10-kDa cut off membrane (Sartorius Group, Göttingen, Germany). A 1.0 M solution of (NH₄)₂SO₄ contained in 50 mM MOPS (pH 6.8) was added to the concentrated protein solution to final concentration of 900 mM and loaded onto a Phenyl-Sepharose FF (low sub) column (20-ml bed volume) equilibrated with 50 mM MOPS (pH 6.8) containing 1.0 M (NH₄)₂SO₄. The column was developed with 100 ml of a 1.0-0.0 M (NH₄)₂SO₄ decreasing linear gradient. Fractions from the peak of CO dehydrogenase activity were pooled and concentrated followed by addition of a volume of 50 mM MOPS (pH 6.8) to lower the (NH₄)₂SO₄ concentration to below 100 mM and then loaded on a HiTrap Q-Sepharose HP column (5 ml bed volume) equilibrated with 50 mM MOPS buffer (pH 6.8). The column was developed with 50 ml of a 0-1.0 M NaCl linear gradient. The peak containing CO dehydrogenase activity that eluted at approximately 0.3 M NaCl was collected and stored at -80°C until use.

Purification of ferredoxin

All purification steps and biochemical assays were performed anaerobically in the anaerobic chamber. Ferredoxin was assayed by the ability to couple CO oxidation by CdhAE to the reduction of metronidazole followed by the decrease in A₃₂₀ ($\epsilon_{320} = 9300 \text{ M}^{-1} \text{ cm}^{-1}$) similar to that described previously [27]. One unit of activity was the amount that reduced 1 μmol of metronidazole/

min. The reaction mixture (100 μ l) contained 100 μ M metronidazole and 1-3 μ g CdHAE in 50 mM Tris buffer (pH 8.0) to which 1-10 μ l of the column fraction was added. The reaction was contained in an anaerobic cuvette flushed with 100% CO.

The soluble fraction of cell extract from acetate-grown *M. acetivorans* was loaded onto a Q-sepharose FF column (20 ml bed volume) equilibrated with 50 mM MOPS (pH 6.8) containing 10% (v/v) ethylene glycol. The column was developed with 200 ml of a 0-1.0 M linear NaCl gradient. The fraction with the highest activity was then diluted 10-fold with 50 mM MOPS (pH 6.8) containing 10% (v/v) ethylene glycol. The solution was loaded on a Mono Q column (1.7 ml bed volume) to which 10 ml of a 0-1.0 M NaCl linear gradient was applied. The fraction containing ferredoxin that eluted at 600 mM NaCl was loaded on a Sephadex G-75 gel filtration column (100 ml bed volume) and developed with 50 mM MOPS (pH 6.8) containing 10% (v/v) ethylene glycol and 150 mM NaCl. The peak containing the purified ferredoxin was concentrated to $A_{402} > 0.2$ with a Vivacell 70 protein concentrator equipped with a 5-kDa cutoff membrane and stored at -80°C until use. The protein concentration was estimated by the ratio of absorbance at 230 and 260 nm as described [43].

Analytical

All protein concentrations except for ferredoxin were determined by the bicinchoninic acid assay using the reagent from Thermo Scientific, Inc.. Detection of the free sulfhydryl groups of CoM-SH and CoB-SH was performed as previously described [17]. The buffer used in the assay was 25 mM sodium acetate containing 1 mM DTNB (5,5'-dithiobis-(2-nitrobenzoic acid)). All assays in this study were performed anaerobically with vacuum degassed solutions contained in sealed cuvettes with the indicated atmosphere and at room temperature.

Nucleotide sequence accession number

The sequences of DNA encoding Rnf and Mrp of *M. thermophila* have been deposited in the GenBank database under accession number JN173061, JN173062, JN173063, JN173064, JN173065, JN173066, JN173067, JN173068, JN173069, JN173070, JN173071, JN173072, JN173073, JN173074, JN173075.

Additional material

Additional file 1: Figure S1. UV-visible absorption spectra of purified ferredoxin. As-purified (–), dithionite reduced (..). The protein concentration was 20 μ M.

Additional file 2: Figure S2. Phylogenetic analysis and sequence alignment of ferredoxins. The *M. mazei* and *M. acetivorans* sequences, labeled with the prefix MA, were derived from the CMR database [23]. The *M. thermophila* (Mt.) sequence is published [26]. The sequence of the $2 \times [4\text{Fe-4S}]$ *Clostridium pasteurianum* is published [44] and the

sequence of the 2Fe-2S *Spinacia oleracea* ferredoxin was obtained from the NCBI database (accession number O04683). Panel A, Phylogenetic analysis of ferredoxins. The tree was constructed by the neighbor-joining method with the MEGA4 program [45]. Bootstrap values are shown at the nodes. Bar, evolutionary distance of 0.2. Panel B, Sequence alignment of ferredoxins from *Methanosarcina* species. Motifs predicted to ligate two 4Fe-4S clusters are highlighted. The alignment was performed with ClustalX2 [46].

Additional file 3: Figure S3. Comparison of rnf genes between Methanosarcina thermophila and Methanosarcina acetivorans. Panel A. Organization of rnf genes in *Methanosarcina thermophila* versus *Methanosarcina acetivorans*. Numbers next to the arrows indicate deduced sequence identity. Panel B. Alignment of the deduced sequences of rnf genes between *Methanosarcina thermophila* (Mt) and *Methanosarcina acetivorans* (Ma). Highlighted are: conserved heme binding sites (CXXCH and CXXXCH) in Cyt c, the flavin binding motif (SGAT) in RnfG, and cysteine motifs binding iron-sulfur clusters in RnfC and RnfB.

Additional file 4: Figure S4. Alignment of mrp gene clusters between Methanosarcina thermophila and Methanosarcina acetivorans. Numbers next to the arrows indicate deduced sequence identity.

Acknowledgements

This work was supported by the National Science Foundation. We thank Dr. Jan Keltjens for generously supplying CoM-S-S-CoB and the Penn State-Hershey Core Research Facilities for mass spectrometry analyses.

Author details

¹Department of Biochemistry and Molecular Biology, Eberly College of Science, The Pennsylvania State University, University Park, Pennsylvania 16802-4500, USA. ²E. I. DuPont de Nemours Company, Central Research and Development, Experimental Station, Wilmington, Delaware 19880, USA.

Authors' contributions

MW carried out the biochemical studies, participated in sequence analysis and drafted the manuscript. J-F T carried out the genomic sequencing and sequence alignments. JGF conceived of the study, participated in its design and coordination, and finalized the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 13 April 2011 Accepted: 24 July 2011 Published: 24 July 2011

References

1. Liu Y, Whitman WB: Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. *Ann N Y Acad Sci* 2008, **1125**:171-189.
2. Ferry JG: How to make a living exhaling methane. *Annu Rev Microbiol* 2010, **64**:453-473.
3. Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R: Methanogenic archaea: ecologically relevant differences in energy conservation. *Nat Rev Microbiol* 2008, **6**:579-591.
4. Guss AM, Kulkarni G, Metcalf WW: Differences in hydrogenase gene expression between *Methanosarcina acetivorans* and *Methanosarcina barkeri*. *Journal of Bacteriology* 2009, **191**(8):2826-2833.
5. Meuer J, Kuettner HC, Zhang JK, Hedderich R, Metcalf WW: Genetic analysis of the archaeon *Methanosarcina barkeri* Fusaro reveals a central role for Ech hydrogenase and ferredoxin in methanogenesis and carbon fixation. *Proc Natl Acad Sci USA* 2002, **99**(8):5632-5637.
6. Fischer R, Thauer RK: Ferredoxin-dependent methane formation from acetate in cell extracts of *Methanosarcina barkeri* (strain MS). *FEBS Lett* 1990, **269**:368-372.
7. Meuer J, Bartoschek S, Koch J, Kunkel A, Hedderich R: Purification and catalytic properties of Ech hydrogenase from *Methanosarcina barkeri*. *Eur J Biochem* 1999, **265**(1):325-335.

8. Welte C, Kratzer C, Deppenmeier U: Involvement of Ech hydrogenase in energy conservation of *Methanosarcina mazei*. *FEBS J* 2010, **277**(16):3396-3403.
9. Welte C, Kallnik V, Grapp M, Bender G, Ragsdale S, Deppenmeier U: Function of Ech hydrogenase in ferredoxin-dependent, membrane-bound electron transport in *Methanosarcina mazei*. *Journal of Bacteriology* 2010, **192**(3):674-678.
10. Galagan JE, Nusbaum C, Roy A, Endrizzi MG, Macdonald P, FitzHugh W, Calvo S, Engels R, Smirnov S, Atnoor D, et al: The genome of *M. acetivorans* reveals extensive metabolic and physiological diversity. *Genome Res* 2002, **12**(4):532-542.
11. Nelson MJ, Ferry JG: Carbon monoxide-dependent methyl coenzyme M methylreductase in acetotrophic *Methanosarcina* spp. *Journal of Bacteriology* 1984, **160**:526-532.
12. Deppenmeier U, Muller V: Life close to the thermodynamic limit: how methanogenic archaea conserve energy. *Results Probl Cell Differ* 2008, **45**:123-152.
13. Li Q, Li L, Rejtar T, Lessner DJ, Karger BL, Ferry JG: Electron transport in the pathway of acetate conversion to methane in the marine archaeon *Methanosarcina acetivorans*. *J Bacteriol* 2006, **188**(2):702-710.
14. Biegel E, Müller V: Bacterial Na⁺-translocating ferredoxin:NAD⁺ oxidoreductase. *Proc Natl Acad Sci USA* 2010, **107**:18138-18142.
15. Buan NR, Metcalf WW: Methanogenesis by *Methanosarcina acetivorans* involves two structurally and functionally distinct classes of heterodisulfide reductase. *Mol Microbiol* 2010, **75**:843-853.
16. Kamlage B, Blaut M: Characterization of cytochromes from *Methanosarcina* strain Go1 and their involvement in electron transport during growth on methanol. *Journal of Bacteriology* 1992, **174**:3921-3927.
17. Peer CW, Painter MH, Rasche ME, Ferry JG: Characterization of a CO: heterodisulfide oxidoreductase system from acetate-grown *Methanosarcina thermophila*. *Journal of Bacteriology* 1994, **176**:6974-6979.
18. Murakami E, Deppenmeier U, Ragsdale SW: Characterization of the intramolecular electron transfer pathway from 2-hydroxyphenazine to the heterodisulfide reductase from *Methanosarcina thermophila*. *J Biol Chem* 2001, **276**:2432-2439.
19. Smith KS, Ingram-Smith C: *Methanosaeta*, the forgotten methanogen? *Trends Microbiol* 2007, **7**:150-155.
20. Grahame DA: Catalysis of acetyl-CoA cleavage and tetrahydroscarinapterin methylation by a carbon monoxide dehydrogenase-corrinoid enzyme complex. *J Biol Chem* 1991, **266**:22227-22233.
21. Gong W, Hao B, Wei Z, Ferguson DJ Jr, Tallant T, Krzycki JA, Chan MK: Structure of the a2e2 Ni-dependent CO dehydrogenase component of the *Methanosarcina barkeri* acetyl-CoA decarbonylase/synthase complex. *Proc Natl Acad Sci USA* 2008, **105**(28):9558-9563.
22. Li L, Li Q, Rohlin L, Kim U, Salmon K, Rejtar T, Gunsalus RP, Karger BL, Ferry JG: Quantitative proteomic and microarray analysis of the archaeon *Methanosarcina acetivorans* grown with acetate versus methanol. *J Proteome Res* 2007, **6**(2):759-771.
23. The Comprehensive Microbial Resource. *J Craig Venter Institute* 2011 [http://cmr.tigr.org/tigr-scripts/CMR/CMrHomePage.cgi].
24. Clements AP, Kilpatrick L, Lu WP, Ragsdale SW, Ferry JG: Characterization of the iron-sulfur clusters in ferredoxin from acetate-grown *Methanosarcina thermophila*. *Journal of Bacteriology* 1994, **176**:2689-2693.
25. Terlesky KC, Ferry JG: Purification and characterization of a ferredoxin from acetate-grown *Methanosarcina thermophila*. *J Biol Chem* 1988, **263**:4080-4082.
26. Clements AP, Ferry JG: Cloning, nucleotide sequence, and transcriptional analyses of the gene encoding a ferredoxin from *Methanosarcina thermophila*. *Journal of Bacteriology* 1992, **174**:5244-5250.
27. Terlesky KC, Ferry JG: Ferredoxin requirement for electron transport from the carbon monoxide dehydrogenase complex to a membrane-bound hydrogenase in acetate-grown *Methanosarcina thermophila*. *J Biol Chem* 1988, **263**:4075-4079.
28. Hovey R, Lentjes S, Ehrenreich A, Salmon K, Saba K, Gottschalk G, Gunsalus RP, Deppenmeier U: DNA microarray analysis of *Methanosarcina mazei* Go1 reveals adaptation to different methanogenic substrates. *Mol Genet Genomics* 2005, **273**:225-239.
29. Abken HJ, Tietze M, Brodersen J, Baumer S, Beifuss U, Deppenmeier U: Isolation and characterization of methanophenazine and the function of phenazines in membrane-bound electron transport of *Methanosarcina mazei* Go1. *Journal of Bacteriology* 1998, **180**:2027-2032.
30. Biegel E, Schmidt S, Gonzalez JM, Muller V: Biochemistry, evolution and physiological function of the Rnf complex, a novel ion-motive electron transport complex in prokaryotes. *Cell Mol Life Sci* 2011, **68**:613-634.
31. Saum R, Schlegel K, Meyer B, Muller V: The F1FO ATP synthase genes in *Methanosarcina acetivorans* are dispensable for growth and ATP synthesis. *FEMS Microbiology Letters* 2009, **300**(2):230-236.
32. Simianu M, Murakami E, Brewer JM, Ragsdale SW: Purification and properties of the heme- and iron-sulfur-containing heterodisulfide reductase from *Methanosarcina thermophila*. *Biochemistry* 1998, **37**(28):10027-10039.
33. Carbon-dependent control of electron transfer and central carbon pathway genes for methane biosynthesis in the Archaeon, *Methanosarcina acetivorans* strain C2A. .
34. Lessner DJ, Li L, Li Q, Rejtar T, Andreev VP, Reichlen M, Hill K, Moran JJ, Karger BL, Ferry JG: An unconventional pathway for reduction of CO₂ to methane in CO-grown *Methanosarcina acetivorans* revealed by proteomics. *Proc Natl Acad Sci USA* 2006, **103**:17921-17926.
35. Rother M, Oelgeschlager E, Metcalf WM: Genetic and proteomic analyses of CO utilization by *Methanosarcina acetivorans*. *Arch Microbiol* 2007, **188**(5):463-472.
36. Rother M, Metcalf WW: Anaerobic growth of *Methanosarcina acetivorans* C2A on carbon monoxide: an unusual way of life for a methanogenic archaeon. *Proc Natl Acad Sci USA* 2004, **101**:16929-16934.
37. Zinder SH, Mah RA: Isolation and characterization of a thermophilic strain of *Methanosarcina* unable to use H₂-CO₂ for methanogenesis. *Applied and Environmental Microbiology* 1979, **38**:996-1008.
38. Zinder SH, Sowers KR, Ferry JG: *Methanosarcina thermophila* sp. nov., a thermophilic, acetotrophic, methane-producing bacterium. *Int J Syst Bacteriol* 1985, **35**:522-523.
39. Li Q, Li L, Rejtar T, Lessner DJ, Karger BL, Ferry JG: Electron transport in the pathway of acetate conversion to methane in the marine archaeon *Methanosarcina acetivorans*. *Journal of Bacteriology* 2006, **188**:702-710.
40. Sowers KR, Baron SF, Ferry JG: *Methanosarcina acetivorans* sp. nov., an acetotrophic methane-producing bacterium isolated from marine sediments. *Applied and Environmental Microbiology* 1984, **47**:971-978.
41. Sowers KR, Nelson MJ, Ferry JG: Growth of acetotrophic, methane-producing bacteria in a pH auxostat. *Curr Microbiol* 1984, **11**:227-230.
42. Terlesky KC, Nelson MJ, Ferry JG: Isolation of an enzyme complex with carbon monoxide dehydrogenase activity containing a corrinoid and nickel from acetate-grown *Methanosarcina thermophila*. *Journal of Bacteriology* 1986, **168**:1053-1058.
43. Kalb VF, Bernlohr RW: A new spectrophotometric assay for protein in cell extracts. *Anal Biochem* 1977, **82**:362-371.
44. Graves MC, Mullenbach GT, Rabinowitz JC: Cloning and nucleotide sequence determination of the *Clostridium pasteurianum* ferredoxin gene. *Proc Natl Acad Sci* 1985, **82**:1653-1657.
45. Tamura K, Dudley J, Nei M, Kumar S: MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 2007, **24**(8):1596-1599.
46. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, et al: Clustal W and Clustal X version 2.0. *Bioinformatics* 2007, **23**(21):2947-2948.

doi:10.1186/1471-2180-11-165

Cite this article as: Wang et al.: Electron transport in acetate-grown *Methanosarcina acetivorans*. *BMC Microbiology* 2011 **11**:165.