

Enhancing parental knowledge of childhood and adolescence safety

An interventional educational campaign

Mohamad-Hani Temsah, MD, FRCPCH^{a,b,c}, Fadi Aljamaan, MD^{a,d}, Ali Alhaboob, MD^{a,b}, Badr Almosned, MD^b, Raghad Alsebil, MD^b, Reem Temsah^e, Abdulrahman Senjab^e, Abdulrahman Alarfaj^a, Talal Aljudi^a, Amr Jamal, MD^{a,f,g,*}, Alia Habash, MD^{a,h}, Fahad Alsohime, MD^{a,b}, Mohammed Almazayad, MD^{a,b}, Majed Alabdulhafid, MD^{a,b}, Gamal Hasan, MD^{i,j}, Raghad Asaad Assiri^k, Wa'ad Massoud Almunassir Alqahtani^k, Adi Alherbish, MD^{a,b}, Khalid Alhasan, MD^{a,b}, Ayman Al-Eyadhy, MD^{a,b}

Abstract

Background: Safeguarding children and adolescents from unintentional injuries is a significant concern for parents and caregivers. With them staying more at home during the coronavirus disease 2019 pandemic, more educational tools and valid educational programs are warranted to improve parental knowledge and awareness about childhood and adolescences' safety. This study aims to explore the effectiveness of childhood and adolescence safety campaigns on parents' knowledge and attitude toward preventable injuries.

Methods: This was a pre–post experimental study, in which the predesigned assessments were used as an evaluation tool before and after attending a childhood and adolescence safety campaign. The pre–post assessment question included questions to evaluate the socio-demographic status, followed by knowledge questions in line with the current childhood and adolescence safety campaign. The outcomes of interest were assessed before and after attending the campaign's stations.

Results: Three hundred eight parents volunteered to participate in this study. Their knowledge score improved from 36.2 [standard deviation (SD) 17.7] to 79.3 (SD 15.6) after attending the Campaign (t value=34.6, $P < .001$). Both, perceptions on the preventability of accidents and the parents' perceived usefulness of educational campaigns showed improvements, with (t value=6.3, $P < .001$) and (t value=3.097, $P < .001$), respectively.

Conclusion: The educational childhood and adolescence safety campaign for caregivers in Saudi Arabia resulted in a significant increase in the overall knowledge and attitudes toward childhood and adolescence's safety. As children and adolescents are currently staying at home more, additional educational tools and programs are warranted to promote safe practices among parents and caregivers.

Editor: Khaled Saad.

MHT and FA contributed equally to this work.

The authors are grateful to the Deanship of Scientific Research, King Saud University, for funding through the Vice Deanship of Scientific Research Chairs.

This research has been financially supported by Prince Abdullah Ben Khalid Celiac Disease Research Chair, under the Vice Deanship of Research Chairs, King Saud University, Riyadh, Kingdom of Saudi Arabia.

All authors gave their consent for publication.

All the data for this study will be made available upon reasonable request.

The authors have no conflicts of interest to disclose.

Supplemental Digital Content is available for this article.

The datasets generated during and/or analyzed during the current study are publicly available.

^a College of Medicine, King Saud University, Riyadh, Saudi Arabia, ^b Pediatrics Department, King Saud University Medical City, Riyadh, Saudi Arabia, ^c Prince Abdullah Ben Khalid Celiac Disease Research Chair, King Saud University, Riyadh, Saudi Arabia, ^d Critical care department, King Saud University Medical City, Riyadh, Saudi Arabia, ^e Al-Faisal University, Riyadh, Saudi Arabia, ^f Department of Family and Community Medicine, King Saud University Medical City, Riyadh, Saudi Arabia, ^g Evidence-Based Health Care & Knowledge Translation Research Chair, King Saud University, Riyadh, Saudi Arabia, ^h Emergency Medicine Department, King Saud University Medical City, Riyadh, Saudi Arabia, ⁱ Department of Pediatrics, Assiut University Children Hospital, Assiut University, Egypt, ^j Department of Pediatrics, Sheikh Shakhboub Medical City, Abu Dhabi, UAE, ^k College of Medicine, Imam Mohammad Ibn Saud Islamic University, Riyadh, Saudi Arabia.

* Correspondence: Amr Jamal, College of Medicine, King Saud University, PO Box 14135, Riyadh 11424, Saudi Arabia (e-mail: Med.researcher.2020@gmail.com).

Copyright © 2022 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Temsah MH, Aljamaan F, Alhaboob A, Almosned B, Alsebil R, Temsah R, Senjab A, Alarfaj A, Aljudi T, Jamal A, Habash A, Alsohime F, Almazayad M, Alabdulhafid M, Hasan G, Assiri RA, Alqahtani WMA, Alherbish A, Alhasan K, Al-Eyadhy A. Enhancing parental knowledge of childhood and adolescence safety: an interventional educational campaign. *Medicine* 2022;101:3(e28649).

Received: 5 October 2021 / Received in final form: 14 December 2021 / Accepted: 3 January 2022

<http://dx.doi.org/10.1097/MD.00000000000028649>

Abbreviations: COVID-19 = Coronavirus disease 2019, IT = information technology, KAP = knowledge, attitude and perceptions, MVA = motor vehicle accidents, SPSS = Statistical Package for the Social Sciences.

Keywords: child injury prevention, childhood and adolescence safety campaign, parental educational effects

1. Introduction

Children are naturally vulnerable and curious; therefore, they tend to explore anything and everything around them, unable to distinguish between harmful and safe objects. This might lead to unintentional injuries that most commonly occur in the home environment because they spend most of their time indoors. Some hazards that might cause injury to children include stairs, sharp objects, and toxic products.^[1] Moreover, many of these unintentional injuries can be prevented by the caregivers responsible for providing a safe, habitable, and healthy environment for children and by teaching them principal methods of self-protection. Thus, caregivers should be encouraged to have good knowledge about the right practices that promote childhood and adolescence safety.

Unintentional injuries amongst children are of primary concern all over the world. The consequences can be dire, as the injured child may develop permanent disabilities such as skin burns, amputations, fractures; and in the worst case, injuries may lead to death.^[2] A study conducted in the United States showed that 40% of children's deaths from ages 1 to 19 years were related to unintentional injuries^[1]; other studies in Bangladesh, Columbia, Egypt, and Pakistan showed that half of the severe injuries that required an emergency visit resulted in disability.^[3] Similarly, a study conducted in Columbia, Maryland, by Dershewitz et al^[4] found that the majority of child mortality between the ages of 1 and 14 are associated with accidental causes that account for approximately 2800 minor injuries, 97 major injuries, and 1 death in every 29,000 cases.

The statistics indicate that the frequency of injuries is high, as unintentional childhood injuries might happen anywhere, like at home, school, in the car, on the road, or in a public area. For instance, a study in Italy showed that most unintentional injuries occurred at home (45.4%), followed by on the road (24.3%), and then at sports facilities (20.3%).^[5]

However, many injuries, and their consequences, are preventable. To reduce the risk of accidents and injuries, parents and caregivers need to utilize different self-protection measures and provide a safe environment.^[6] Additionally, one of the most effective prevention techniques is through community awareness campaigns. For instance, in the United States, campaigns held to increase awareness about the use of child's booster seats in motor vehicles showed a significant increase in their use.^[7]

Despite that child safety is a high priority for caregivers, there was no study conducted in Saudi Arabia about the effect of childhood and adolescence safety campaigns on parents or communities. However, this study aimed to measure and analyze the effect of childhood and adolescence safety campaigns on the parents' knowledge and attitudes.

2. Methodology

This was a pre–post experimental study, in which the predesigned assessments were used as an evaluation tool during the childhood and adolescence safety campaign (March 14–17, 2017). The

study assessed its impact on knowledge, attitude, and intended practices among parents in Riyadh, Saudi Arabia.

The targeted population was parents who attended this campaign. The study sample consisted of parents (for children up to 17 years of age and living in Saudi Arabia for at least 1 year), excluding adults with no children living with them. As no previous data exists locally, a convenience sampling technique was used. We incorporated close-ended questions into an electronic format (using Survey Monkey). The questions were piloted among 10 parents to ensure clarity. Then the assessment questionnaire was modified accordingly and tested for validity and consistency before using it in the childhood and adolescence safety campaign.

The main components of the questionnaire were sociodemographic data information including parents' ages, parents' education, current employment, current relationship status, and the number of their children. This was followed by knowledge questions about childhood and adolescence safety that were in line with the current childhood and adolescence safety campaign and parents were tested before and after attending the campaign's stations for their knowledge in that regard, additionally the parents were assessed pre and post-campaign regarding their perceived usefulness of the safety campaigns and belief in childhood injuries preventability.

3. Data collection methods

Parents who attended the childhood and adolescence safety campaign were invited to participate in this survey. Those who consented to participate in this study were interviewed with the precampaign structured questionnaire-A (the bilingual Arabic/English version is available by contacting the corresponding author) (Appendix 1 Pre, Supplemental Digital Content, <http://links.lww.com/MD/G597>).

Questionnaire-A consisted of a series of questions that measured their essential demographic and economic characteristics, alongside their perceived self-rating on childhood and adolescence safety knowledge and their perceptions on the likelihood of preventing childhood and adolescence injuries.

The parents were also assessed for knowledge on childhood and adolescence safety using predesigned questions twice (at baseline and after attending the series of educational materials at the campaign).

Parents answered the postcampaign assessment tool (questionnaire-B, the bilingual Arabic/English version is available by contacting the corresponding author), evaluating the same previously assessed aspects, focusing on possible changes in knowledge and attitude toward childhood and adolescence safety (Appendix 2 Post, Supplemental Digital Content, <http://links.lww.com/MD/G598>). The knowledge questions were reshuffled and paraphrased with 2 additional questions to help alter the questions and avoid recall bias.

Each questionnaire was headed with a letter stating that participation was voluntary, and no identification data were

required. Ethical approval was granted by the Institutional Review Board of College of Medicine, King Saud University (Riyadh, Saudi Arabia).

The analysis was performed using Statistical Package for the Social Sciences v19 (Armonk, NY: IBM Corp). Basic descriptive analysis was utilized to calculate the frequency and proportion of study variables. Means and standard deviation were calculated to describe continuous variables.

Pre and postcampaign parental knowledge about childhood and adolescence main safety targets and goals were compared, parental belief regarding the usefulness of the safety campaigns in addition to their belief in childhood injuries preventability, were compared using paired *t* test, *P* value of $<.05$ was considered significant.

4. Results

A total of 308 parents volunteered to participate during the childhood and adolescence safety campaign and completed the first set of questions (Questionnaire A) before attending the campaign. Thereafter, the same sample were followed up with linked Questionnaire B.

Out of the participating 308 parents, the majority were mothers (68.8%), and the majority of them were aged above 40 years (79.2%) and married (93.8%). Most of them were educated with a college degree or higher (74%). Saudi nationals comprised the majority of the sample (67.2%) while many of them had monthly income greater than 10,000 Saudi Riyals per month (36%). The mean number of children for the whole sample was 3.3 children. Reportedly, the primary caregiver at home was the mother by most respondents (93.6%) (Table 1).

The “knowledge questions” used to assess participants’ information on childhood and adolescence safety are shown in Table 2.

The binomial test showed that a significant proportion of correctly answered questions were observed after the campaign compared to before for most of the used questions with a *P* value $<.001$, except to whether the first life-saving measure for a child with an ingested battery is to take them to the hospital immediately, *P* = .228.

Surveyed parents had significant improvement in their knowledge regarding most of the domains associated with childhood and adolescence safety postcampaign, whether life-threatening ones as poisoning or drowning, or issues related to electronics and information technology, and issues related to even skeletal growth and appropriate school bag size handling. For further details refer to Table 2.

For example, their knowledge of getting rid of expired medications to prevent poisoning improved from 34.7% to 71.4%, hot water of 60 °C as a cause of burns from 19.8% to 94.8%.

Additionally, the campaign had significantly improved parental knowledge of the correct way of using modern information technology and telecommunication devices that are already integrated into most children’s daily lives. Therefore, the caregiver’s knowledge of the “healthy” way of dealing with electronic devices, their relation to obesity, and the safe amount of screen time, have improved significantly postcampaign.

Participating parents knowledge about childhood and adolescence safety was assessed using direct question of their self-knowledge-rating using Likert scale of 1 to 10 before attending the educational campaign; their score was 6.8 out of 10, at the

Table 1

Respondents’ characteristics.

	Frequency	Percentage
Sex of the respondent		
Mother	212	68.8
Father	96	31.2
Age		
Below 40	244	79.2
Above 40	64	20.8
Marital status		
Married	274	93.8
Widowed	7	2.4
Divorced	11	3.8
Educational level		
Elementary/others	17	5.5
Intermediate	9	2.9
High school	54	17.5
College or higher	228	74
Nationality		
Saudi	207	67.2
Other nationalities	101	32.8
Employment		
Employed	276	89.6
Unemployed	9	2.9
Not applicable	8	2.6
Retired	15	4.9
Income		
No answer	74	24
<5000 SAR*	23	7.5
5000 to 10,000 SAR	100	32.5
>10,000 SAR	111	36
Number of children, mean (SD)	3.3 (2.2)	
Principal child caregiver		
Mother	278	93.6
Father	3	1
Housemaid	9	2.9
Others	7	2.3

* SAR = Saudi Riyals.

same time their mean score of the predesigned questions to assess objectively their knowledge regarding the multiple domains of childhood and adolescence safety was equal to 36.2%. However, their score assessed by the same predesigned questions was 79.3%, with a significant difference after the campaign educational sessions; *P* value $<.001$.

Their belief in preventing unintentional child injuries and their perceived usefulness of childhood and adolescence safety campaigns increased significantly postcampaign, as shown in Table 3.

5. Discussion

The number of deaths among children younger than 5 years has declined substantially over the past 47 years, and this is evidence that progress is being made in tackling the fundamental causes of childhood mortality.^[8] An example of a successful strategy includes increasing the educational levels of mothers.^[9] Still, unintentional injuries are a worldwide health problem that causes high death rates among childhood and adolescence younger than 18. Furthermore, they are the most common cause of hospital admissions and permanent disabilities.^[10]

The spectrum of unintentional injuries is diverse, including ingestion of toxic materials such as medications or poisoning

Table 2
Proportions of correctly answered knowledge questions before and after the educational campaign.

Knowledge questions	Before session	After session	P
	n (%) Correctly answered	n (%) Correctly answered	
Getting rid of old medications is the best way to prevent child medication poisoning	107 (34.7%)	220 (71.4%)	<.001
Hot water at 60° can cause burns; as such heaters should always be set to 60° or below	61 (19.8%)	292 (94.8%)	<.001
The first life-saving measure for a child with an ingested battery is to take them to the hospital immediately	130 (42.2%)	185 (60.1%)	.228
iPad devices must remain outside the bedroom of a child when asleep	42 (13.6%)	256 (83.1%)	<.001
Children may not be allowed to use electronic devices (eg, iPad, play stations) for more than 1–2 h per day	205 (66.6%)	269 (96.1%)	<.001
Excessive use of social media leads to obesity among children	124 (40%)	256 (83.1%)	<.001
The appropriate size of the school bag for the weight of the child should not exceed 15% of the child weight	162 (52.6%)	257 (83.4%)	<.001
Your child's school back-bag should be placed on their mid-back when carrying them	–	288 (93.5%)	–
When using bags with wheels, it is preferred to choose bags with large wheels	41 (13.3%)	190 (61.7%)	<.001
The appropriate place to install a child seat in the car is the rear-facing back seat	159 (51.6%)	290 (94.2%)	<.001
A child can use the car's regular seat and seatbelt When the waist belt is in the top level for thighs and the shoulder belt at the chest level	–	171 (55.5%)	–
The infant in the car must be placed in the back seat in a rear-facing seat when driving	159 (51.9%)	209 (67.9%)	<.001
A fundamental rescue task for the drowning child is doing CPR immediately	89 (28.9%)	267 (86.7%)	<.001

Table 3
Parental belief in childhood injuries preventability and perceived usefulness of childhood safety campaigns.

	Before session mean (SD)	After session mean (SD)	P
Belief in childhood safety prevention (%)	67.6 (22.9)	76.3 (20.1)	<.001
Perceived usefulness of childhood safety campaigns (1–10 score)	8.8 (1.7)	9.1 (1.2)	<.001
Knowledge score (%)	36.2 (17.7)	79.3 (15.6)	<.001

materials, foreign bodies ingestion or aspiration, falls, traffic accidents, or burns.^[11–13] Many of these injuries could be prevented if parents and caregivers have good literacy about different types of injury and preventative measures and understand the health information and instructions to deal with emergency matters that may result.^[14,15] However, caregivers may not be adherent to safety practices and the recommended injury prevention tasks. Therefore, continued emphasis on these strategies is recommended.^[16,17]

The present study showed that before the implementation of the educational program, most caregivers had insufficient knowledge about the preventative measures of childhood and adolescence safety issues, including the seatbelt, discarding expired medications and batteries, using very hot water, the appropriate way to use electronic devices, using an appropriate car seat, and first aid for a drowning child. In Saudi Arabia, the 5 major causes of deaths secondary to injuries among children and adolescents were motor vehicle accidents, drowning, child maltreatment, fire and weapon, and finally home accident (fall, poisoning, suffocation).^[18] International papers have reported similar unsatisfactory outcomes, emphasizing the need for global and nationwide interventional educational campaigns to correct the current knowledge deficit.^[19,20]

No previous studies in Saudi Arabia evaluated the knowledge and attitude among caregivers regarding childhood and adolescence safety or assessed the impact of educational safety campaigns on the caregivers' overall knowledge and practices. After the intervention of childhood and adolescence safety campaigns, the overall knowledge, attitude, and perceptions of

the usefulness of campaigns and preventability of injuries were significantly increased. Thus, educational campaigns about childhood and adolescence safety measures could result in increasing the knowledge, attitude, and perceptions of caregivers, and in decreasing the rate of injuries, permanent disabilities as well as child death. A study conducted in Brazil observed an increase in mothers' level of knowledge after the educational intervention was applied compared to their basic prior knowledge, confirming the importance of implementing community awareness programs.^[1]

Educational interventions have shown their tremendous benefits in promoting the level of knowledge and awareness among caregivers and limiting unintentional childhood and adolescence injuries. Therefore, we strongly acknowledge the need for combined efforts of all concerned institutions to establish a national awareness program that can reinforce best practices, reduce ineffective measures, and ensure childhood and adolescence's safety. Many countries have inserted these educational campaigns as a powerful program to increase the protection of children and reduce mortality rates.^[10,21,22] Other modalities that could boost parental education are educational videos in parents' native language, that could be posted on suitable social media platforms.^[23] These educational strategies could be particularly useful for parents without a previous medical background.^[24]

Several nations have decreased the rates of injury-related deaths in more than 50% of children through educational safety programs.^[25] As such, health authorities should establish injury prevention programs like road accidents prevention and clear

child restraint laws. One of the attempts in this regard was the Brazilian National Policy for Reduction of Mortality from Accidents and Violence established by the Brazilian Ministry of Health.^[1]

While there has been a decline in global drowning rates, there is still a need for more prevention and research efforts.^[20] Educational training programs for children's first aid are recommended for all subjects to rescue drowning children. While many Saudi cities are located in a desert-environment, drowning could still occur in recreational swimming pools.^[26] The Saudi Red Crescent Authority has a program called Prince Naif first aid program, which targets all population groups, and could be utilized to improve parents' first-aid skills.^[27]

Other strategies to alleviate harm from burns were successful, such as using smoke alarms, safer lamps, and laws on the temperature of hot-water taps, resulting in decreased incidence of disabilities and deaths from burns during childhood.^[28]

The death count from poisoning by chemical and medical substances could be prevented by sufficient parental supervision and safe storage of such substances while discarding hazardous ones.^[15,29] With the coronavirus disease 2019 pandemic, some agencies, like the Consumer Product Safety Commission in the USA, published Home Safe Checklists, so parents and caregivers could check off the safety items in their home environments.^[30]

While our study was the first to explore the impact of a childhood and adolescence safety campaign amongst parents in Saudi Arabia, it still had some limitations. The self-reported practices may not reflect actual behaviors but remain among the best available tools to assess the population's knowledge, attitude, and reported practices. The use of the same reporting tool twice (pre-post campaign changes) may have added carried-on bias to some participants, though the questions were rephrased to minimize this possibility. Future longitudinal studies that follow the participants are warranted to seek whether these changes are maintained over time; and whether these improvements in parental knowledge and attitudes are translated into better childhood and adolescence safety practices. Future research that is conducted in other regions and countries would give more generalization for our described tool of health education.

6. Conclusion

The educational childhood and adolescence safety campaign for caregivers in this study resulted in a significant increase in the overall knowledge and awareness of children's safety and preventable measures. As many children stayed home during the coronavirus disease 2019 pandemic, more educational tools and programs are warranted to promote childhood and adolescence safety practices among parents and caregivers.

Acknowledgments

We would like to thank hodhodata.com for their statistical analysis.

Author contributions

MHT, FAJ, AAE, BA, and RA conceptualized the study, analyzed the data, and wrote the manuscript. FAS, AAH, KAH, TA, AF, AS, GH, RAA, WMAA, and RT contributed to the study design; collected, analyzed, and interpreted data; and edited the

manuscript. AJ, AH interpreted the data and finalized the manuscript. All authors reviewed and approved the final version of the manuscript.

Conceptualization: Mohamad-Hani Temsah, Ayman Al-Eyadhy.

Data curation: Mohamad-Hani Temsah, Ali Alhaboob, Badr Almosned, Raghad Alsebil, Abdulrahman Senjab, Abdulrahman Alarfaj, Fahad Alsohime.

Formal analysis: Mohamad-Hani Temsah, Fadi Aljamaan, Badr Almosned, Amr Jamal, Fahad Alsohime.

Investigation: Mohamad-Hani Temsah, Ali Alhaboob, Badr Almosned, Raghad Alsebil, Abdulrahman Senjab, Fahad Alsohime, Khalid Alhasan.

Methodology: Mohamad-Hani Temsah, Ayman Al-Eyadhy.

Project administration: Ali Alhaboob, Fahad Alsohime.

Resources: Ali Alhaboob.

Validation: Fadi Aljamaan, Talal Aljudi, Mohammed Almazyad, Majed Alabdulhafid, Gamal Hasan, Raghad Asaad Assiri, Wa'ad Massoud Almunassir Alqahtani.

Visualization: Mohamad-Hani Temsah, Fadi Aljamaan, Reem Temsah, Mohammed Almazyad, Raghad Asaad Assiri, Wa'ad Massoud Almunassir Alqahtani, Adi Alherbish, Khalid Alhasan.

Writing – original draft: Mohamad-Hani Temsah, Fadi Aljamaan, Ali Alhaboob, Raghad Alsebil, Reem Temsah, Abdulrahman Senjab, Abdulrahman Alarfaj, Talal Aljudi, Amr Jamal, Alia Habash, Mohammed Almazyad, Majed Alabdulhafid, Gamal Hasan, Khalid Alhasan, Ayman Al-Eyadhy.

Writing – review & editing: Mohamad-Hani Temsah, Fadi Aljamaan, Ali Alhaboob, Badr Almosned, Reem Temsah, Amr Jamal, Alia Habash, Fahad Alsohime, Mohammed Almazyad, Majed Alabdulhafid, Gamal Hasan, Adi Alherbish, Khalid Alhasan, Ayman Al-Eyadhy.

References

- [1] Silva ECS, de Fátima Fernandes MN, Sá MCN, et al. The effect of educational intervention regarding the knowledge of mothers on prevention of accidents in childhood. *Open Nurs J* 2016;10:113.
- [2] WHO. Violence and Injury Prevention. 2015. Available at: www.who.int/violence_injury_prevention/en/. Accessed February 14, 2021.
- [3] Peden M, Oyegbite K, Ozanne-Smith J, et al. World report on child injury prevention. Geneva: World Health Organization; 2008.
- [4] Dershewitz RA, Christophersen ER. Childhood household safety. An overview. *Am J Dis Child* 1984;138:85–8.
- [5] Santagati G, Vezzosi L, Angelillo IF. Unintentional injuries in children up to six years of age and related parental knowledge, attitudes, and behaviors in Italy. *J Pediatr* 2016;177:267.e2–72.e2.
- [6] Etzel RA. Environmental risks in childhood. *Pediatr Ann* 2004;33:431–6.
- [7] Ebel BE, Koepsell TD, Bennett EE, Rivara FP. Use of child booster seats in motor vehicles following a community campaign: a controlled trial. *Jama* 2003;289:879–84.
- [8] Wang H, Abajobir AA, Abate KH, et al. Global, regional, and national under-5 mortality, adult mortality, age-specific mortality, and life expectancy, 1970–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;390:1084–150.
- [9] Fuchs R, Pamuk E, Lutz W. Education or wealth: which matters more for reducing child mortality in developing countries? *Vienna Yearb Popul Res* 2010;8:175–99.
- [10] Soori H, Khodakarim S. Child unintentional injury prevention in Eastern Mediterranean Region. *Int J Crit Illn Inj Sci* 2016;6:33–9.
- [11] Gielen AC, Wilson ME, McDonald EM, et al. Randomized trial of enhanced anticipatory guidance for injury prevention. *Arch Pediatrics Adolesc Med* 2001;155:42–9.
- [12] Al-Zahrany MS, Al-Shuwair FN, Al-Zahrani MA, et al. Self-reported unintentional injuries in families visiting the 'childhood safety campaign' in Saudi Arabia. *Egypt J Hosp Med* 2018;71:2280–6.

- [13] Alhaboob AA. Sociodemographic characteristics and risk factors for childhood poisoning reported by parents at a tertiary care teaching hospital. *Cureus* 2021;13:e13313.
- [14] Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. *Ann Intern Med* 2011;155:97–107.
- [15] Kumar D, Sanders L, Perrin EM, et al. Parental understanding of infant health information: health literacy, numeracy, and the Parental Health Literacy Activities Test (PHLAT). *Acad Pediatrics* 2010;10:309–16.
- [16] Gielen AC, McDonald EM, Wilson ME, et al. Effects of improved access to safety counseling, products, and home visits on parents' safety practices: results of a randomized trial. *Arch Pediatrics Adolesc Med* 2002;156:33–40.
- [17] Kendrick D, Young B, Mason-Jones AJ, et al. Home safety education and provision of safety equipment for injury prevention. *Cochrane Database Syst Rev* 2012;Cd005014doi: 10.1002/14651858.CD005014.pub3.
- [18] Almuneef M, Saleheen H, AlBuhairan F, et al. Child mortality in Saudi Arabia: time for action at all levels. *Int J Pediatr Adolesc Med* 2021;8:165–71.
- [19] İnce T, Yalçın S, Yurdakök K. Parents' attitudes and adherence to unintentional injury prevention measures in Ankara, Turkey. *Balkan Med J* 2017;34:335–42.
- [20] Franklin RC, Peden AE, Hamilton EB, et al. The burden of unintentional drowning: global, regional and national estimates of mortality from the Global Burden of Disease 2017 Study. *Inj Prev* 2020;26(Suppl 1): i83–95.
- [21] Bener A, Hyder AA, Schenk E. Trends in childhood injury mortality in a developing country: United Arab Emirates. *Accid Emerg Nurs* 2007; 15:228–33.
- [22] Greensher J. The injury fact book. *JAMA* 1992;267:3089–90.
- [23] Temsah MH, Al-Eyadhy A, Alsohime F, et al. Effect of lumbar puncture educational video on parental knowledge and self-reported intended practice. *Int J Pediatr Adolesc Med* 2021;8:112–6.
- [24] Temsah M-H, Alsohaime F, Al-Eyadhy A, et al. Hand hygiene perception and handshaking practices among pediatric inpatient caregivers: a cross-sectional study at a teaching hospital in Saudi Arabia. *J Nat Sci Med* 2021;4:130–4.
- [25] Berfenstam R. Sweden's pioneering child accident programme: 40 years later. *Inj Prev* 1995;1:68–9.
- [26] Temsah MH, Alsohaim F, Al-Eyadhy A, et al. Drowning in the desert: family denial of brain death. *Sudan J Paediatr* 2018;18:48–52.
- [27] Saudi Red Crescent Authority. Available at: <https://training.srca.org.sa/>. Accessed February 14, 2021.
- [28] Sminkey L. World report on child injury prevention. *Inj Prev* 2008; 14:69–169.
- [29] MacKay M, Vincenten J, Brussoni M, Towner L. *Child Safety Good Practice Guide: Good Investments in Unintentional Child Injury Prevention and Safety Promotion*. Amsterdam: European Child Safety Alliance, Eurosafe; 2006. 2–30.
- [30] CPSC. COVID-19: Protect those you love by using our safety checklists in and around your home. Available at: <https://www.cpsc.gov/home-safety>. Accessed June 19, 2021.