



Primary synovial sarcoma of the duodenal bulb: a case report and review of the literature

Hanteng Yang^{1#}, Xiaocheng Cheng^{1#}, Wei Zhu², Jingying Xiao¹, Fan Li¹, Yanxian Ren¹, Keshen Wang¹, Zuoyi Jiao¹, Changjiang Luo¹

¹Department of General Surgery, Lanzhou University Second Hospital, Lanzhou, China; ²Department of Pathology, Lanzhou University Second Hospital, Lanzhou, China

[#]These authors contributed equally to this work.

Correspondence to: Changjiang Luo; Zuoyi Jiao. Department of General Surgery and Cuiying Experimental Center, University Second Hospital, No.82 Cuiyingmen, Cheng-Guan District, Lanzhou, China. Email: 157264922@qq.com; jzyldey@163.com.

Abstract: Primary synovial sarcoma of the duodenal bulb is a rare mesenchymal tumor with special morphological features. It usually originates from the major joints or tendon sheaths of the extremities and mostly seen in young population, but rarely found in gastrointestinal tract. In this manuscript, we reported the first case of synovial sarcoma arising between the intestinal wall of the duodenal bulb with a concomitant *SYT/SSX* type of the *t(X;18)* translocation. A 49-year-old male presented to our hospital with a 2-month history of upper abdominal pain along with a 4-day amply jaundice. Tumor marker testing showed only a slight increase of carbohydrate antigen 19-9 (CA19-9). A computed tomography scan of his abdomen showed that indeterminate tissue occupied the duodenal bulb wall, compressed the surrounding tissues, and measured roughly 5.0 cm × 7.7 cm × 8.7 cm. Since the sarcoma grows between the intestinal wall, which cannot be detected by endoscopy, an initial diagnosis of duodenal wall stromal tumor was made at that time. Postoperative Immunohistochemistry results showed that the tumor was positive for the expression of transducin-like enhancer of split 1, B-cell lymphoma 2, and Vimentin. These pathological findings were indicative of the diagnosis of synovial sarcoma, but still did not provide sufficient diagnostic evidence. Finally we confirmed the diagnosis by using fluorescence in situ hybridization (FISH) with detection of the *t(X;18)* (*SYT-SSX*) translocation. No such lesions were found on preoperative examination, so a diagnosis of primary duodenal synovial sarcoma was made. After literature review, we found four reports of duodenal synovial sarcomas, all of which could be detected endoscopically, but there were no results of long-term follow-up. This case is the first reported case of synovial sarcoma arising between the intestinal walls of the duodenal bulb treated twice with ifosfamide and followed up for 13 months without recurrence.

Keywords: Synovial sarcoma; duodenal bulb; immunohistochemistry; fluorescence in situ hybridization

Submitted Feb 20, 2020. Accepted for publication Jul 27, 2020.

doi: 10.21037/tcr-20-1107

View this article at: <http://dx.doi.org/10.21037/tcr-20-1107>

Introduction

Synovial sarcoma is a type of mesenchymal tumor that originates from soft tissue rather than from synovial tissue. This sarcoma originates from soft tissue, has a special predilection for local recurrence and distant metastasis, and it accounts for 5–10% of all soft tissue tumors (1,2). synovial sarcoma usually occurs in young adults around major joints or

tendon sheaths (1) and occasionally in the head and neck (3), lungs (4), heart (5), retroperitoneum (6), prostate (7), within nerves (8), and the kidney (9). Here, we report a 49-year-old Chinese male with primary synovial sarcoma of the duodenal bulb. We also retrieved relevant data on SS in other parts of the digestive system and reviewed the literature. We present the following article in accordance with the CARE reporting checklist (available at <http://>




Figure 1 Computed tomography and magnetic resonance imaging. (A,B) Abdominal computed tomography shows that the tumor was located in the duodenal bulb and had invaded the hilum of the liver; (C,D) abdominal magnetic resonance imaging revealed a diffuse large mass in the antrum of the stomach and the medial part of the duodenal bulb.

dx.doi.org/10.21037/tcr-20-1107).

Case presentation

A 49-year-old male presented to our hospital with a 2-month history of upper abdominal pain along with a 4-day amply jaundice. The patient was previously healthy, normotensive, did not have diabetes, and did not have a history of alcohol intake or smoking. According to the blood test results, he had a total bilirubin level of 43.3 $\mu\text{mol/L}$ and a direct bilirubin level of 40.4 $\mu\text{mol/L}$, a γ -GGT level of 1,798 U/L, an ALP level of 1,454 U/L, and a carbohydrate antigen 19-9 level of 29.16 U/L. A computed tomography (CT) scan of his abdomen revealed that the lump occupied the duodenal bulb wall, compressed the surrounding tissues, and measured roughly 5.0 cm \times 7.7 cm \times 8.7 cm. The arterial phase of an enhanced CT scan showed that the mass was obviously strengthened and that the branch of the pancreaticoduodenal artery was supplied with sufficient blood (*Figure 1A,B*). A Magnetic Resonance Imaging (MRI) scan of his abdomen showed that the duodenum bulb contained a large lump with an approximate size of

9.1 cm \times 5.5 cm \times 7.1 cm; the T1 signal was slightly longer, and the T2 signal was dominant. The arterial phase of an enhanced MRI showed that the mass was obviously strengthened and had an irregular shape (*Figure 1C,D*). On gastroduodenoscopy, the duodenal bulb and descending portion exhibited no abnormalities. Since the tumor was located between the duodenal walls, it could not be examined by endoscopic ultrasound and needle biopsy. We thus considered a diagnosis of duodenal wall stromal tumor.

After a multidisciplinary discussion, the patient was originally diagnosed with a malignant duodenal tumor with obstructive jaundice, and consequently, underwent radical pancreatoduodenectomy. Surgical specimens showed that the duodenal papilla had some edema, and a nodular mass observed below the nipple measured 12 cm \times 8 cm \times 5 cm. The tumor surface was grayish red and gray, and some areas appeared as translucent nodules. The postoperative pathological examination showed that the tumor tissue was fusiform, arranged in a bundle with oval nuclei that were long and fusiform, and contained pink cytoplasm (*Figure 2A-C*). Immunohistochemistry showed that the tumor was positive for the expression of transducin-like enhancer of




Figure 2 Histomorphology of duodenal bulb synovial sarcoma. The tumor tissue is fusiform and bundled. The nuclei are round, long, fusiform, and the cytoplasm is pink [hematoxylin and eosin, 40× (A), 100× (B), 200× (C)].

split 1 (TLE-1), B-cell lymphoma 2 (Bcl-2), Vimentin, and smooth muscle actin (Figure 3A-D), but was negative for CD34 (Figure 3E), Desmin and S-100 protein; 67% of cells were Ki-67-positive (Figure 3F,G). These pathological findings were indicative of the diagnosis of synovial sarcoma, but still did not provide sufficient diagnostic evidence.

Cytogenetic and fluorescence in situ hybridization (FISH) examination were performed on formalin-fixed paraffin-embedded tumor specimens. FISH was used to detect the fusion of the *SSX* gene, which is located on the X chromosome, and the *SS18* gene (*SYT*) located on

chromosome 18; this fusion produces the *SS18* (*SYT*)-*SSX* fusion gene (Figure 4A,B,C) and thus, we detected *SS18* gene translocations (Figure 4A,B,C). Therefore, FISH demonstrated the *t(X;18)* (*SYT-SSX*) gene rearrangement and confirmed the above diagnosis. No such lesions were found on preoperative examination, so a diagnosis of primary duodenal synovial sarcoma was made. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee(s) and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient.

At the time of the reported case, the patient had been treated with ifosfamide twice in the 3 months after surgery. The dose of ifosfamide was 2 g/time for 5 days, and the interval between two chemotherapy was 1 month. After 13 months of follow-up, his condition was stable and remained no signs of clinical or imaging recurrence, and timely follow-up has been conducted. The patient's treatment and follow-up timeline was shown in Figure 5.

Discussion

Synovial sarcoma is a type of mesenchymal tumor that originates in soft tissue rather than from synovial tissue. synovial sarcoma is divided into three tissue types (2). One is the monophasic type, which consists only of spindle-shaped blastomas arranged in bundles, vortices, and herringbone. Another is the biphasic type, which is composed of different proportions of epithelial cells and spindle cells. The other is the poorly differentiated type, which is characterized by darkly stained oval or round cells, as seen in other small round-cell tumors. synovial sarcoma is difficult to diagnose and is easily misdiagnosed, which is closely related to the diagnostic methods used. CT, MRI, and digestive endoscopy are used to diagnose SS of the digestive system, but considering the current state of the technology, it is difficult to make a definitive diagnosis before surgery. Since the diagnosis of synovial sarcoma relies on histology, immunohistochemistry [TLE-1 and Bcl-2 positivity can indicate a diagnosis of synovial sarcoma (10,11)], and cytogenetics, in most cases, the disease is confirmed using the above three methods to examine surgical specimens. Moreover, 90% of SS is associated with the *t(X;18)(p11.2;q11.2)* chromosomal translocation, which results in the fusion of the *SYT* gene on chromosome 18 with the *SSX1* or *SSX2* gene on the X chromosome. This results in the *SYT-SSX* fusion protein under certain




Figure 3 Immunohistochemistry. Transducin-like enhancer of the split 1 (A, 100 \times), B-cell lymphoma 2 (B, 100 \times), Vimentin (C, 100 \times), and smooth muscle actin (D, 200 \times) showed diffuse, strong positive staining. (E,G) The Ki-67 proliferation index was 67% (100 \times). (E, 100 \times) CD34 negativity.

conditions (cytogenetics, fluorescence in situ hybridization, and RT-PCR can detect this fusion). If the translocation can be identified in tumor tissue, then histology can be used to make a diagnosis (12). The molecular test results of *SYT-SSX* were positive in our case (Figure 4A,B,C), and combined histological and immunohistochemical analyses showed multiple characteristics of monophasic synovial sarcoma. Therefore, the diagnosis of synovial sarcoma of the duodenal bulb was established (2,10,11). The differential diagnoses of synovial sarcoma include digestive lymphoma, gastrointestinal stromal tumor (GIST), giant juvenile polyp, sarcoma, and ectopic pancreas. It is worth mentioning that GIST is the most common mesenchymal cell tumor. Sometimes, it is difficult to identify GIST and gastrointestinal synovial sarcoma under the microscope. The diagnosis thus needs to be confirmed by TLE-1 and Bcl-2 positivity by immunohistochemistry or the presence of the t(X;18) chromosomal translocation by cytogenetics

testing.

In this case, confirmation of the diagnosis was not easy. The tumor originated between the intestinal walls of the duodenal bulb, so gastroduodenoscopy failed to detect it, and the location of the tumor was ultimately confirmed by enhanced CT and MRI. Therefore, imaging physician made the diagnosis of duodenal stromal tumor before surgery. After pathological and immunohistochemical examination of the postoperative specimen, the pathologist considered synovial sarcoma, but still could not make a definite diagnosis. FISH examination was not routinely conducted in our institution due to its high cost, however, in order to obtain a definitive diagnosis, we had the patient's consent and performed FISH on the specimen, which helped us make the final diagnosis of duodenal synovial sarcoma. This case may serve as experience for gastrointestinal surgeons. First, the tumor is located between the intestinal walls of the duodenal bulb, which is rare and made biopsy difficult




Figure 4 Detection of the t(X;18) (SYT-SSX) translocation by fluorescence in situ hybridization by a break apart probe demonstrated one fused (yellow arrow) and one separate red and green (red and green arrow) signal indicating the rearrangement of the SYT gene.




Figure 5 The timeline figure of patient treatment and follow-up.

and we cannot make a correct diagnosis in a timely manner. Second, in the previous reports, the location of duodenal synovial sarcoma was all in the duodenal lumen, but in this case, the lesion occurred between the intestinal walls of the duodenal bulb and could not be found endoscopically, and the same report has not been seen in the past. Therefore, the report of this case helps gastroenterologists to increase vigilance when encountering such diseases to avoid misdiagnosis.

Until now, 33 cases of synovial sarcoma of the digestive system have been reported, including 10 cases in the esophagus, 1 case in the esophagogastric junction, 13 cases in the stomach, 1 case in the gastroduodenal junction, 5 cases in the duodenum, 1 case in the jejunum, 1 case in the ileum, and 1 case in the colon. The four cases previously reported can be found under endoscopy. One case relapsed 8 months after surgery, and three cases lost follow-up results. We first reported the case of synovial sarcoma originating between the intestinal wall of the duodenal bulb, and had a follow-up time of 13 months. The clinical situation of previous cases and our case are summarized in *Table 1* (12-33). Symptoms include dysphagia, upper abdominal pain, and bleeding. We can conclude that the median age of patients with synovial sarcoma of the digestive system is 42 years old (range, 14–76 years), and although there is no obvious gender difference, the ratio of males to females is approximately 1:1. The tumor size ranges from 2 to 16 cm. The macroscopic morphology is primarily polypoid. There have been 17 cases of the monophasic type, 12 cases of the biphasic type, and 2 cases of the poorly differentiated subtype. The histological type of 2 cases was not reported in the literature. The monophasic type is common in synovial sarcoma. In the literature, 32 cases were treated with surgery, and 1 report did not describe the treatment. Eight patients received adjuvant chemotherapy after surgery, 3 patients received radiation therapy, and 3 patients received chemotherapy and radiation therapy. The survival of patients in the reported cases ranged from 1 to 224 months.

Surgery is still the primary treatment for synovial sarcoma of the digestive system without tumor metastasis. Chemotherapy and radiation therapy are mainly used for preoperative and postoperative adjuvant therapy. Ifosfamide and doxorubicin are the first-line chemotherapeutic drugs, while pazopanib is a second-line drug. Although pazopanib is the only tyrosine kinase inhibitor approved by the FDA for the treatment of synovial sarcoma, due to the resistance of tumors to pazopanib, its therapeutic effect is not satisfactory. In recent studies, the over-activation of IGF-1

and insulin receptor (IGF1R/InsR) was shown to alter the AKT and ERK pathways in synovial sarcoma cells. AKT and ERK activity was reduced, which in turn reduced the resistance of tumor cells to pazopanib (34). In recent years, radiation therapy has rarely been used in the treatment of synovial sarcoma of the digestive system. Research on targeted therapy has received substantial attention, and recent studies by Isfort *et al.* (35) confirmed that activation of YAP/TAZ signaling is a common pattern in SS and is functionally dependent on the SS18-SSX fusion protein, which interferes with YAP. The interference in signal transduction of TAZ through small molecule inhibitors may provide a new and effective way for the treatment of synovial sarcoma. synovial sarcoma also expresses survivin. The BirM 5 promoter induced by YM 155 can change its activity, and thus, survivin may be a feasible target for the treatment of synovial sarcoma (36). In summary, the above research mechanism requires further clarification, and its clinical application is still in the experimental stage. Whether it can be used for the treatment of digestive tract synovial sarcoma has not been confirmed by relevant literature or clinical data. We should accelerate clinical big data studies, make breakthroughs on the basis of existing research, and strive to use the above research results for the treatment of synovial sarcoma in the digestive tract and elsewhere. At present, there are few reports on duodenal synovial sarcoma, and there are no relevant guidelines for postoperative chemotherapy. We used ifosfamide for chemotherapy in this patient according to the chemotherapy regimen for synovial sarcoma of other tissues. The patient was treated with chemotherapy twice without significant adverse reactions. Up to now, there was no recurrence or metastasis.

The 5- and 10-year survival rates of patients with synovial sarcoma of bone tissue are 76.4% and 60.4%, respectively, which are related to tumor size, tumor grade, chemotherapy, and radiation therapy (37). Since synovial sarcoma of the digestive system is difficult to diagnose, easily misdiagnosed, has a high degree of malignancy, and few cases are seen, there is no relevant literature that reports the long-term survival rate of patients with this disease. After following up for 13 months, our case was no recurrence or metastasis. Bergh *et al.* (38) divided synovial sarcoma patients into a low-risk group (patients age <25 years, tumor size <5 cm, histological classification as monophasic or biphasic) and a high-risk group (patient age <25 years, tumor size >5 cm, histological classification as poorly differentiated). It has also been suggested that

Table 1 Reported cases of synovial sarcoma of the digestive tract

Author, year, reference	Location	Presenting symptoms	Age, years	Gender	Gross features	Size, cm	Histologic type	Translocation	Treatment	Follow-up status and (months)
Palmer <i>et al.</i> 1983, (13)	Esophagus	Dysphagia	75	Female	Polypoid	2.5	Biphasic	-	Surgery + radiotherapy	Died of other cause, 24
Amr <i>et al.</i> 1984, (14)	Esophagus	Dysphagia	25	Male	Polypoid	5	Biphasic	-	Surgery	Alive without evidence of disease, 36
Bloch <i>et al.</i> 1987, (15)	Esophagus	Dysphagia, dyspnea	15	Male	Polypoid	10	Biphasic	-	Surgery + radiotherapy	Alive without evidence of disease, 36
Pulpeiro <i>et al.</i> 1988, (16)	Esophagus	-	24	Male	-	-	Biphasic	-	Surgery	-
Caldwell <i>et al.</i> 1991, (17)	Esophagus	-	29	Female	-	-	-	-	Surgery + chemotherapy + radiotherapy	Alive without evidence of disease, 195
Perch <i>et al.</i> 1991, (18)	Esophagus	-	15	Male	-	-	Biphasic	-	Surgery + radiotherapy	Alive without evidence of disease, 5 to 6years after surgery
Antón-Pacheco <i>et al.</i> 1996, (19)	Esophagus	Dysphagia, weight loss	14	Female	Polypoid	7	Biphasic	-	Surgery + chemotherapy + radiotherapy	Alive without evidence of disease, 30
Habu <i>et al.</i> 1998, (20)	Esophagus	Sensation of something stuck in his throat	20	Male	Polypoid	8	Biphasic	-	Surgery + chemotherapy + radiotherapy	Alive without evidence of disease, 20
Bonavina <i>et al.</i> 1998, (21)	Esophagus	Achalasia	63	Female	Polypoid	-	-	-	-	-
Billings <i>et al.</i> 2000, (22)	Gastroesophageal junction	Incidental finding for pyloric Stenosis	47	Female	Polypoid	5.2	Biphasic	t(X;18)	Surgery	Alive without evidence of disease, 21
Billings <i>et al.</i> 2000, (22)	Stomach	Abdominal pain, nausea, vomiting	55	Female	Spherical intramural	16	Biphasic and poorly differentiated synovial sarcoma	t(X;18)	Surgery	Died of other cause, 6
Chan <i>et al.</i> 2004, (23)	Jejunum	Epigastric pain, vomit, fever	28	Male	Polypoid intramural	15	Monophasic	t(X;18)	Surgery	Died of other cause, 1
Butori <i>et al.</i> 2006, (24)	Esophagus	Dysphagia	72	Female	Polypoid	11	Biphasic	t(X;18)	Surgery + chemotherapy	6
Akhuji <i>et al.</i> 2007, (25)	Stomach	Epigastric pain	42	Male	-	11	Biphasic	t(X;18)	Surgery + chemotherapy	Died of other cause, 24
Parfitt <i>et al.</i> 2007, (26)	Colon	Rectal bleeding	32	Male	Polypoid	2	Monophasic	t(X;18)	Surgery	5

Table 1 (continued)

Table 1 (continued)

Author, year, reference	Location	Presenting symptoms	Age, years	Gender	Gross features	Size, cm	Histologic type	Translocation	Treatment	Follow-up status and (months)
Schreiber-Facklam <i>et al.</i> 2007, (27)	Duodenum	Abdominal pain	39	Female	Polypoid	5	Monophasic	t(X;18)	Surgery + chemotherapy	Recurrence 8 months after surgery
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	67	Female	-	0.8	Monophasic	t(X;18)	Surgery	Alive without evidence of disease, 12
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	49	Male	-	2	Monophasic with a poorly differentiated component	t(X;18)	Surgery	Died of other cause, omental metastasis, 29
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	68	Female	-	2	Monophasic	t(X;18)	Surgery	Alive without evidence of disease, 22
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	29	Male	-	2.8	Monophasic	t(X;18)	Surgery	Alive without evidence of disease, 224
Makhlouf <i>et al.</i> 2008, (28)	Stomach, gastroduodenal junction	-	54	Female	-	3	Monophasic	t(X;18)	Surgery	Recent case
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	58	Female	-	3	Monophasic	t(X;18)	Surgery	Alive without evidence of disease, 21
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	37	Female	-	4	Monophasic	t(X;18)	Surgery	Local recurrence, re-excised. died of other cause, 48
Makhlouf <i>et al.</i> 2008 (28)	Stomach	-	50	Male	-	6	Monophasic	t(X;18)	Surgery + chemotherapy	Alive with recurrence, 6
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	42	Male	Polypoid	8	Biphasic	t(X;18)	Surgery + chemotherapy	Died of other cause, 25
Makhlouf <i>et al.</i> 2008, (28)	Stomach	-	66	Female	Polypoid	15	Monophasic	t(X;18)	Surgery	Lost to follow up
Borens-Fefer <i>et al.</i> 2009, (29)	Duodenum	Pain	14	Male	-	5	Monophasic	t(X;18)	Surgery + chemotherapy	Lost to follow up
Company Campins <i>et al.</i> 2009, (12)	Proximal duodenum	Weight loss Asthenia anorexia, nausea, epigastric pain	69	Female	Spherical intramural	8	Monophasic	t(X;18)	Surgery	Died due to complications, 1
García Ruiz <i>et al.</i> 2010 (30)	Duodenal	Pain	70	Male	Polypoid	9	Biphasic	t(X;18)	Surgery	Lost to follow up

Table 1 (continued)

Table 1 (continued)

Author, year, reference	Location	Presenting symptoms	Age, years	Gender	Gross features	Size, cm	Histologic type	Translocation	Treatment	Follow-up status and (months)
Alsharief et al. 2012, (31)	Ileum	Abdominal pain, distension and heaviness	29	Female	Intramural	8	Monophasic	-	Surgery	Alive without evidence of disease, 6
Wang et al. 2012, (32)	Stomach	Abdominal pain	38	Female	-	7	Monophasic	t(X;18)	Surgery + chemotherapy	Alive without evidence of disease
Sahara et al. 2013, (33)	Stomach	Abdominal pain	22	Male	-	-	Monophasic	t(X;18)	Surgery	Lost to follow up
Present case 2019	Duodenal bulb	Abdominal pain	49	Male	-	12	Monophasic	t(X;18)	Surgery + chemotherapy	Alive without evidence of disease, 13 (continue to follow)

the genotyping and prognostic correlation of SS remains uncertain (39). synovial sarcoma appears in the head and neck, trunk, lung pleura, and peritoneal organs, and areas other than the limbs usually have a worse prognosis. Moreover, tumor size (>5 cm), nerve infiltration, vascular invasion, P53 overexpression, and high expression of Ki-67 increase the risk of SS metastasis and recurrence (40-42). We would like to know if the prognosis of synovial sarcoma of the digestive system is related to the above-mentioned factors. Unfortunately, no clinical data have demonstrated the prognostic factors of digestive system synovial sarcoma. Therefore, in order to solve this problem, we should collect more cases and join more medical institutions to make a breakthrough in the prognosis of synovial sarcoma in digestive system.

In conclusion, primary synovial sarcoma of the digestive system is rare and easily misdiagnosed. We reported the first case of synovial sarcoma arising between the intestinal walls of the duodenal bulb with a concomitant SYT/SSX type of the t(X;18) translocation. At present, there are few reports on the diagnosis and treatment of duodenal synovial sarcoma. We hope to deepen our general perception and understanding of synovial sarcoma through report of this rare sarcoma.

Acknowledgments

We would like to thank all members of the Department of Pathology, Lanzhou University Second Hospital, Lanzhou, Gansu, China, for their expert technical assistance.

Funding: This research was funded by the Lanzhou University Second Hospital “Cuiying Technology Innovation” Program (CY-2017-ZD03).

Footnote

Reporting Checklist: The authors have completed the CARE reporting checklist. Available at <http://dx.doi.org/10.21037/tcr-20-1107>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tcr-20-1107>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are

appropriately investigated and resolved. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee(s) and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Sultan I, Rodriguez-Galindo C, Saab R, et al. Comparing children and adults with synovial sarcoma in the Surveillance, Epidemiology, and End Results program, 1983 to 2005: an analysis of 1268 patients. *Cancer* 2009;115:3537-47.
- Suurmeijer AJH, de Bruijn D, Geurts van Kessel A, et al. *Synovial Sarcoma*. Lyon: IARC Press; 2013.
- Park JK, Ham SY, Hwang JC, et al. Synovial sarcoma of the head and neck: a case of predominantly cystic mass. *AJNR* 2004;25:1103-5.
- Etienne-Mastroianni B, Falchero L, Chalabreysse L, et al. Primary sarcomas of the lung: a clinicopathologic study of 12 cases. *Lung Cancer* 2002;38:283-9.
- Eswaran P, Devadoss P, Narasimhan LS, et al. Synovial sarcoma of the heart: A case report and literature review. *J Cancer Res Ther* 2015;11:659.
- Chatzipantelis P, Kafiri G. Retroperitoneal synovial sarcoma: a clinicopathological study of 6 cases. *J Buon* 2008;13:211-6.
- Jun L, Ke S, Zhaoming W, et al. Primary synovial sarcoma of the prostate: report of 2 cases and literature review. *Int J Surg Pathol* 2008;16:329-34.
- Chu PG, Benhattar J, Weiss LM, et al. Intraneural synovial sarcoma: two cases. *Mod Pathol* 2004;17:258-63.
- Tranesh G, Cortese C, Thiel D, et al. Primary synovial sarcoma of the kidney-A case report and literature review. *Int J Clin Exp Pathol* 2018;11:2241-5.
- Jagdis A, Rubin BP, Tubbs RR, et al. Prospective evaluation of TLE1 as a diagnostic immunohistochemical marker in synovial sarcoma. *Am J Surg Pathol* 2009;33:1743-51.
- Barrott JJ, Zhu JF, Smith-Fry K, et al. The Influential Role of BCL2 Family Members in Synovial Sarcomagenesis. *Mol Cancer Res* 2017;15:1733-40.
- Company Campins MM, Morales R, Dolz C, et al. Primary monophasic synovial sarcoma of the duodenum confirmed by cytogenetic analysis with demonstration of t(X;18): a case report. *J Gastrointest Liver Dis* 2009;18:89-93.
- Palmer BV, Levene A, Shaw HJ. Synovial sarcoma of the pharynx and oesophagus. *J Laryngol Otol* 1983;97:1173-6.
- Amr SS, Shihabi NK, Al Hajj H. Synovial sarcoma of the esophagus. *Am J Otolaryngol* 1984;5:266-9.
- Bloch MJ, Iozzo RV, Edmunds LH Jr, et al. Polypoid synovial sarcoma of the esophagus. *Gastroenterology* 1987;92:229-33.
- Pulpeiro JR, Cruz R, Arenas A, et al. Para-oesophageal synovial sarcoma. *Eur J Radiol* 1988;8:120-1.
- Caldwell CB, Bains MS, Burt M. Unusual malignant neoplasms of the esophagus. Oat cell carcinoma, melanoma, and sarcoma. *J Thorac Cardiovasc Surg* 1991;101:100-7.
- Perch SJ, Soffen EM, Whittington R, et al. Esophageal sarcomas. *J Surg Oncol* 1991;48:194-8.
- Antón-Pacheco J, Cano I, Cuadros J, et al. Synovial sarcoma of the esophagus. *J Pediatr Surg* 1996;31:1703-5.
- Habu S, Okamoto E, Toyosaka A, et al. Synovial sarcoma of the esophagus: report of a case. *Surg Today* 1998;28:401-4.
- Bonavina L, Fociani P, Asnaghi D, et al. Synovial sarcoma of the esophagus simulating achalasia. *Dis Esophagus* 1998;11:268-71.
- Billings SD, Meisner LF, Cummings OW, et al. Synovial sarcoma of the upper digestive tract: a report of two cases with demonstration of the X;18 translocation by fluorescence in situ hybridization. *Mod Pathol* 2000;13:68-76.
- Chan GS, Yuen ST, Chan KW. Synovial sarcoma presenting as a polypoid jejunal mass. *Histopathology* 2004;44:191-3.
- Butori C, Hofman V, Attias R, et al. Diagnosis of primary esophageal synovial sarcoma by demonstration of t(X;18) translocation: a case report. *Virchows Arch* 2006;449:262-7.
- Akhunji S, Musil I, Baisre de Leon A, et al. Synovial sarcoma arising in the gastric wall: case report and literature review. *Cancer Therapy* 2007;5:457-62.
- Parfitt JR, Xu J, Kontozoglou T, et al. Primary

- monophasic synovial sarcoma of the colon. *Histopathology* 2007;50:521-3.
27. Schreiber-Facklam H, Bode-Lesniewska B, Frigerio S, et al. Primary monophasic synovial sarcoma of the duodenum with SYT/SSX2 type of translocation. *Hum Pathol* 2007;38:946-9.
 28. Makhoulouf HR, Ahrens W, Agarwal B, et al. Synovial sarcoma of the stomach: a clinicopathologic, immunohistochemical, and molecular genetic study of 10 cases. *Am J Surg Pathol* 2008;32:275-81.
 29. Borens-Fefer B, Lingier P, Cassart M, et al. Synovial sarcoma of the duodenum in adolescent. *JBR-BTR* 2009;92:142-3.
 30. García Ruiz S, Jiménez Rodríguez RM, Alcaide León P, et al. Synovial two-phase sarcoma in third portion of the duodenum: clinical case and review of the literature. *Rev Esp Enferm Dig* 2010;102:62-3.
 31. Alsharief AN, Fageeh M, Alabdulkarim Y. Monophasic synovial sarcoma presenting as a primary ileal mass: a case report and review of the literature. *J Med Case Rep* 2012;6:83.
 32. Wang CC, Wu MC, Lin MT, et al. Primary gastric synovial sarcoma. *J Formos Med Assoc* 2012;111:516-20.
 33. Sahara S, Otsuki Y, Egawa Y, et al. Primary synovial sarcoma of the stomach--a case report and review of the literature. *Pathol Res Pract* 2013;209:745-50.
 34. Lanzi C, Dal Bo L, Favini E, et al. Overactive IGF1/Insulin Receptors and NRASQ61R Mutation Drive Mechanisms of Resistance to Pazopanib and Define Rational Combination Strategies to Treat Synovial Sarcoma. *Cancers* 2019;11:408.
 35. Isfort I, Cyra M, Elges S, et al. SS18-SSX-Dependent YAP/TAZ Signaling in Synovial Sarcoma. *Clin Cancer Res* 2019;25:3718-31.
 36. Mika A, Luelling SE, Pavek A, et al. Epigenetic Changes at the Birc5 Promoter Induced by YM155 in Synovial Sarcoma. *J Clin Med* 2019;8:408.
 37. Outani H, Nakamura T, Murata H, et al. Localized synovial sarcoma: A single institutional study of 191 patients with a minimum follow-up of 5 years for survivors. *J Surg Oncol* 2019;119:850-5.
 38. Bergh P, Meis-Kindblom JM, Gherlinzoni F, et al. Synovial sarcoma: identification of low and high risk groups. *Cancer* 1999;85:2596-607.
 39. Guillou L, Benhattar J, Bonichon F, et al. Histologic grade, but not SYT-SSX fusion type, is an important prognostic factor in patients with synovial sarcoma: a multicenter, retrospective analysis. *J Clin Oncol* 2004;22:4040-50.
 40. Antonescu CR, Leung DH, Dudas M, et al. Alterations of cell cycle regulators in localized synovial sarcoma: A multifactorial study with prognostic implications. *Am J Pathol* 2000;156:977-83.
 41. Lewis JJ, Antonescu CR, Leung DH, et al. Synovial sarcoma: a multivariate analysis of prognostic factors in 112 patients with primary localized tumors of the extremity. *J Clin Oncol* 2000;18:2087-94.
 42. Trassard M, Le Doussal V, Hacène K, et al. Prognostic factors in localized primary synovial sarcoma: a multicenter study of 128 adult patients. *J Clin Oncol* 2001;19:525-34.

Cite this article as: Yang H, Cheng X, Zhu W, Xiao J, Li F, Ren Y, Wang K, Jiao Z, Luo C. Primary synovial sarcoma of the duodenal bulb: a case report and review of the literature. *Transl Cancer Res* 2020;9(9):5663-5673. doi: 10.21037/tcr-20-1107