

ORIGINAL ARTICLE

Association of a complement receptor 1 gene variant with baseline erythrocyte sedimentation rate levels in patients starting anti-TNF therapy in a UK rheumatoid arthritis cohort: results from the Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate cohort

J Bluett^{1,8}, I Ibrahim^{1,8}, D Plant², KL Hyrich¹, AW Morgan^{3,4}, AG Wilson⁵, JD Isaacs⁶ for BRAGGS⁷ and A Barton^{1,2}

Eligibility for anti-tumour necrosis factor (TNF) therapy in most European countries is restricted to severe, active rheumatoid arthritis (RA). The DAS28 score is a marker of disease severity and incorporates one of two inflammatory markers, erythrocyte sedimentation rate (ESR) or C-reactive protein. We aimed to determine the relation between genetic variants known to affect ESR and levels of ESR in patients with active RA. DNA samples were genotyped for four single-nucleotide polymorphisms (SNPs) rs7527798 (CR1L), rs6691117 (CR1), rs10903129 (TMEM57) and rs1043879 (C1orf63). The association between SNPs and baseline ESR, baseline DAS28-ESR, and change in DAS28-ESR was evaluated. Baseline ESR was significantly associated with CR1 rs6691117 genotype ($P = 0.01$). No correlation was identified between baseline DAS28-ESR or change in DAS28-ESR. In conclusion, genetic variation in the gene encoding CR1 may alter ESR levels but not DAS28-ESR, indicating no adjustment for CR1 genotype is required in the assessment of patients with severe active RA.

The Pharmacogenomics Journal (2014) **14**, 171–175; doi:10.1038/tpj.2013.26; published online 16 July 2013

Keywords: blood sedimentation; immunogenetics; rheumatoid arthritis; tumour necrosis factor-alpha

INTRODUCTION

The development of biologic drugs that block the tumour necrosis factor (TNF) pathway has revolutionized rheumatoid arthritis (RA) treatment and patient prognosis. Anti-TNF drugs reduce joint inflammation, diminish radiological damage and may reduce cardiovascular risk.^{1,2} However, due to increased risk of infection, inefficacy in a subset of patients and the economic impact, predictors of treatment response would be a major clinical advance.³ In the United Kingdom, eligibility for biologics is determined by guidance issued by the National Institute for Health and Clinical Excellence (NICE).⁴ Eligibility to commence and maintain treatment with anti-TNF therapy is determined by the 28 joint-count disease activity score (DAS28).⁵ The DAS28 is an assessment used to measure the level of disease activity in patients with RA and has been validated in several studies.^{6–8} A score of ≥ 5.1 on two separate occasions at least 1 month apart is required before UK patients are eligible for anti-TNF therapy.

DAS28 is a composite index of RA disease activity, which includes number of swollen and tender joints in 28 specified joints, patient global health as measured by a visual analogue

scale and one of two inflammatory markers, erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP).⁶ CRP is an acute-phase protein that is produced by the liver and is very sensitive to short-term changes in inflammation.⁹ ESR is measured by the rate at which red blood cells sediment over 1 h and is reported in mm h^{-1} . In contrast to CRP, raised ESR indicates long-standing, chronic inflammation and is an indirect measure of acute-phase protein levels with a slower response after inflammatory stimulation or resolution.⁹ ESR is increased in a variety of conditions including pregnancy, myeloma, anaemia and is further affected by age and gender, but this is not controlled for in the calculation of the DAS28-ESR.¹⁰ A recent genome-wide association study identified single-nucleotide polymorphisms (SNPs) in several genes, which were associated with ESR levels at genome-wide significance thresholds ($P \leq 1 \times 10^{-8}$) including CR1 rs6691117.¹¹ If this genetic correlation is also observed in RA patients, it could have important clinical implications when assessing eligibility for anti-TNF therapy using the DAS28-ESR. CR1 encodes complement receptor 1 (CD35), a membrane glycoprotein present on erythrocytes and leucocytes that acts as

¹Arthritis Research UK Epidemiology Unit, Manchester Academy of Health Science, University of Manchester, Manchester, UK; ²NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK; ³NIHR Leeds Musculoskeletal Biomedical Research Unit, Chapel Allerton Hospital, The Leeds Teaching Hospitals NHS Trust, Leeds, UK; ⁴Leeds Institute of Molecular Medicine, Wellcome Trust Brenner Building, St. James's University Hospital, Leeds, UK; ⁵Department of Infection and Immunity, University of Sheffield Medical School, Sheffield, UK and ⁶Musculoskeletal Research Group, Newcastle upon Tyne NHS Foundation Trust, Institute of Cellular Medicine, Newcastle University and NIHR Newcastle Biomedical Research Centre, Newcastle-upon-Tyne, UK. Correspondence: Professor A Barton, Arthritis Research UK Epidemiology Unit, Centre for Musculoskeletal Research, Institute of Inflammation and Repair, The University of Manchester, Manchester Academic Health Science Centre, Stopford Building, Oxford Road, Manchester M13 9PT, UK.

E-mail: anne.barton@manchester.ac.uk

⁷Member details are listed in Appendix.

⁸These authors contributed equally to this work.

Received 2 February 2013; revised 23 April 2013; accepted 29 May 2013; published online 16 July 2013

a negative regulator of the complement cascade by increasing clearance of complement opsonized immune complexes, thus preventing immune complex deposition.^{12–14}

The aim of the current study was first, to investigate the importance of known genetic variants that affect ESR and determine whether they significantly influence ESR levels in UK patients with active RA, and secondly, to determine whether the genetic variants correlate with treatment response to anti-TNF medication. We aimed to investigate the rs7527798, rs6691117, rs10903129 and rs1043879 SNPs mapping to the *CR1L*, *CR1*, *TMEM57* and *C1orf63* genes, respectively, which have each been associated with ESR levels, to determine their association with baseline ESR, baseline DAS28-ESR and change in DAS28-ESR in patients with RA before and after 6 months therapy with an anti-TNF drug.

MATERIALS AND METHODS

Subjects

DNA samples from patients included in this study were obtained from the Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate (BRAGGSS). Patients eligible for the BRAGGSS cohort were initially identified through the British Society for Rheumatology Biologics Register (BSRBR). The BSRBR is a prospective observational study of patients with rheumatic diseases newly commenced on anti-TNF biologic therapy, who are followed up every 6 months for a period of at least 5 years.¹⁵ One of the fundamental objectives of the BSRBR is to monitor patient progress, as well as the incidence of long- and short-term side effects. The BRAGGSS cohort was developed for the study of genetic predictors of response to anti-TNF biologic therapy. Consultants at contributing centres across the United Kingdom gave permission to identify their patients from the BSRBR; eligible patients were approached by letter and invited to donate blood samples for DNA extraction when they were due for a routine blood test. Samples were posted to the Arthritis Research UK Epidemiology Unit for processing, storage and analyses. All contributing patients provided informed consent, and the study was approved by a multicenter ethics committee (COREC 04/Q1403/37).

Baseline and 6-month DAS28 values were recorded to allow subsequent analysis. Patients were excluded from this study if they had stopped treatment because of adverse events or reasons other than inefficacy, or after any change in their anti-TNF biologic therapy during the follow-up period.

Genotyping

DNA samples were genotyped using the Sequenom MassArray iPLEX system. In each reaction, 10 ng of DNA was used and the protocol was followed according to the manufacturer's instructions (<http://www.sequenom.com>). For each marker, negative water controls were included for each experiment, and genotype cluster plots were manually reviewed. In addition, SNPs were assessed for deviation from Hardy–Weinberg equilibrium. For purposes of quality control, a 90% sample threshold and 90% genotyping success threshold were used.

Statistics

Baseline levels of ESR in the cohort studied did not follow a normal distribution and were positively skewed; thus, baseline ESR values were log-transformed before analysis. The association between SNPs and ESR was evaluated with linear regression under an additive effect model. Analyses were repeated adjusting for gender and age at baseline. Linear regression models were also used to analyze DAS28-ESR and change in DAS28-ESR over a 6-month period of treatment with anti-TNF therapy.

These analyses were performed using STATA V.11.2 (<http://www.stata.com>). Power calculations were performed using Quanto (version 1.2.3) (<http://hydra.usc.edu/gxe>) under an additive model for a range of marker-allele frequencies.

RESULTS

Clinical response and demographic data were recorded in 2978 patients. In total, 264 stopped their anti-TNF drug for reasons other than inefficacy, 12 had no recorded information regarding a

potential change in their therapy and 146 had either an incomplete baseline DAS28 or 6-month follow-up DAS28.

The rs7527798 (*CR1L*), rs6691117 (*CR1*), rs10903129 (*TMEM57*) and rs1043879 (*C1orf63*) SNP markers were genotyped in 1510 DNA samples. The genotyping success rates for rs7527798, rs6691117, rs1043879 and rs10903129 were 96%. Ninety-six individuals were removed due to genotype success < 90%.

In total, 1223 samples were successfully genotyped for all four SNPs with recorded baseline DAS28 and 6-month follow-up DAS28. Of the samples with genotype information, ESR serum measurements were available for 1188 samples at baseline (pretreatment) and 1195 samples at 6 months. Table 1 describes the demographic and disease characteristics of the BRAGGSS cohort. Genotype frequencies of all four markers conformed to Hardy–Weinberg equilibrium (Table 2). For 1223 individuals for whom change in DAS28-ESR was available, the study had > 80% power (at the 5% significance threshold) to detect a clinically meaningful difference of 0.6 DAS28-ESR units for an allele frequency of 5%.

As expected, following treatment with anti-TNF therapy, we noted a decrease in serum ESR levels and DAS28-ESR within the BRAGGSS cohort. Age was significantly associated with higher baseline ESR, baseline DAS28 and DAS28 6 months after anti-TNF treatment ($P < 0.001$, $= 0.002$ and < 0.001 , respectively). Female sex was significantly associated with higher baseline ESR, baseline DAS28 and DAS28 6 months after anti-TNF treatment ($P = 0.004$, < 0.001 and < 0.001 , respectively).

Two copies of rs6691117 *CR1* GG minor allele were significantly associated with baseline ESR, as shown in Table 2, but not ESR after 6 months of treatment or change of ESR. After correcting for gender and age effects the statistical significance remained ($P = 0.01$). Association remained after correcting for anti-CCP serology. The *CR1* rs6691117 GG was not significantly associated with baseline DAS28-ESR, change in DAS28-ESR or DAS28-ESR after 6 months of treatment ($P = 0.99$, 0.78 , 0.77 respectively). The *CR1L* rs7527798 CC genotype was significantly associated with change in DAS28-ESR following 6 months treatment with anti-TNF therapy ($P = 0.05$) but not change in ESR over the same period ($P = 0.26$). These data are presented in Supplementary Tables S1–5.

DISCUSSION

Recent reports have identified that genetic variants on chromosome 1 affect ESR levels. This has potentially important consequences for RA patients as eligibility for anti-TNF therapy is determined by DAS28, which may incorporate ESR as a

Table 1. Demographic and disease characteristics

Age ^a (years)	56.75 (10.89)
Disease duration ^b (years)	12 (6–20)
Sex, F: <i>n</i> (%)	951 (77.76)
ESR at baseline ^b , mm h ⁻¹	40 (25–66)
Change in ESR ^b , mm h ⁻¹	– 14 (– 3 – 30)
DAS28-ESR at baseline ^a	6.67 (0.97)
Change in DAS28-ESR ^a	– 2.49 (1.52)
Concurrent methotrexate therapy, <i>n</i> (%)	705 (57.65)
Rheumatoid factor positive, <i>n</i> (%)	719 (64.31)
Anti-CCP positive, <i>n</i> (%)	828 (80.78)
Infliximab, <i>n</i> (%)	490 (40.07)
Etanercept, <i>n</i> (%)	493 (40.31)
Adalimumab, <i>n</i> (%)	240 (19.62)

Abbreviations: DAS28, Disease Activity Score 28; ESR, erythrocyte sedimentation rate; F, female; IQR, interquartile range.

^aValues are expressed as mean (s.d.).

^bValues are expressed as median (IQR). All other values are *n* (%). Disease duration was measured in 1180 patients; baseline erythrocyte sedimentation was recorded in 1188 patients; change in ESR was recorded in 1195 patients; and change in DAS28-ESR was available in 1188 patients.

Table 2. Relation of ESR genotypes to baseline ESR levels

SNP	HWE (P)	Genotype	Geometric mean baseline ESR (mm h ⁻¹) (s.d.)	n	β-Coefficient	P-value
CR1L rs7527798	0.30	TT	38.51 (2.00)	598	-0.042	0.21
		TC	36.11 (2.17)	500		
		CC	36.74 (2.05)	90		
CR1 rs6691117	0.46	AA	36.11 (2.13)	750	0.098	0.01
		AG	38.67 (2.00)	393		
		GG	48.21 (1.72)	45		
TMEM57 rs10903129	0.43	GG	35.85 (2.19)	364	0.040	0.18
		AG	37.71 (2.02)	575		
		AA	38.74 (2.05)	249		
C1orf63 rs1043879	0.86	AA	36.26 (2.08)	647	0.049	0.16
		AG	38.67 (2.07)	461		
		GG	38.79 (2.08)	80		

Abbreviations: ESR, erythrocyte sedimentation rate; HWE, Hardy–Weinberg expectation; SNP, single-nucleotide polymorphism. Bold value is a statistically significant result.

biomarker of inflammation. Therefore, as genetic markers affect ESR levels, these markers could influence treatment decisions and could have adverse consequences for those patients who carry the genotypes associated with lower ESR, as they would be less likely to meet eligibility criteria for anti-TNF therapy.

In the current study, two copies of the AA major allele in rs6691117 of the CR1 gene are significantly associated with lower baseline ESR levels prior to anti-TNF therapy in keeping with the direction of effect reported by Kullo *et al.*¹¹ ($P = 7 \times 10^{-12}$). Importantly, however, the genotype did not associate with baseline DAS28-ESR. Although we were unable to adjust for all factors that are associated with ESR, in particular haemoglobin levels, we were able to adjust for age and gender that did not qualitatively affect the results. While this is reassuring, the current cohort had severe active RA, and therefore higher ESR compared with the cohort in Kullo *et al.*¹¹ (46.8 versus 13.1 mm h⁻¹ in the discovery cohort).¹⁶ We cannot exclude that the genetic markers affecting ESR levels may still impact DAS28-ESR in patients with less active inflammatory disease than this severe active RA cohort. If the DAS28 thresholds were to be reduced in the future, the genetic markers may influence the eligibility for anti-TNF therapy. Collection of a cohort of patients with early RA would be required to investigate this possibility. Our primary aim was to investigate if genetic markers affected the eligibility to be started on any anti-TNF; we are unable to investigate a class effect between anti-TNF treatments as the reduced sample size would have limited power. CR1 is a potent inhibitor of complement activation and genetic variation within this gene may affect Rouleaux formation and hence ESR. The rs6691117 is a non-synonymous SNP encoding an isoleucine to valine alteration. This may modify the secondary structure of CR1 affecting its ability to clear complement opsonized immune complexes, thereby increasing ESR.

Carriage of the minor allele CC at rs7527798 (CR1L) was associated with change in DAS28-ESR, but not baseline ESR or change of ESR. This most likely represents a false-positive result as the rs7527798 polymorphism has not, to our knowledge, been associated with swollen joints, tender joints or patient global health.

Failure to detect an association with baseline or 6-month DAS28-ESR may reflect inadequate power to detect modest effects. However, despite not reaching significance thresholds, the direction of effect for rs7527798, rs6691117, rs10903129 and rs1043879 were in keeping with previously reported directions in a previous association study.¹¹

CRP can also be used to calculate DAS28-CRP. A previous study has shown that genetic variants at the CRP locus correlate with the level of CRP.¹⁷ However, a recent study has shown that for patients with severe active RA, CRP and DAS28-CRP are not affected by these genetic markers.¹⁸

CONCLUSIONS

In summary, rs6691117 polymorphism has been shown to influence ESR levels in patients with very active inflammation but not DAS28-ESR. This is reassuring for those patients starting anti-TNF, but further studies with a cohort that includes a wide range of DAS28-ESR levels would be required to ensure rs6691117 does not affect DAS28-ESR and hence eligibility for anti-TNF therapy.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank Arthritis Research UK for their support (grant no. 17552). This report includes independent research supported by the National Institute for Health Research Biomedical Research Unit Funding Scheme. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

REFERENCES

- Peters MJL, van Sijl AM, Voskuyl AE, Sattar N, Smulders YM, Nurmohamed MT. The effects of tumor necrosis factor inhibitors on cardiovascular risk in rheumatoid arthritis. *Curr Pharm Des* 2012; **18**: 1502–1511.
- Malaviya AP, Ostor AJK. Rheumatoid arthritis and the era of biologic therapy. *Inflammopharmacology* 2012; **20**: 59–69.
- Rubbert-Roth A. Assessing the safety of biologic agents in patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2012; **51**(Suppl 5): v38–v47.
- NICE guidelines. *Rheumatoid Arthritis, National clinical guideline for management and treatment in adults*. Royal College of Physicians: London, 2009.
- van der Heijde DM, van't Hof MA, van Riel PL, van Leeuwen MA, van Rijswijk MH, van de Putte LB. Validity of single variables and composite indices for measuring disease activity in rheumatoid arthritis. *Ann Rheum Dis* 1992; **51**: 177–181.
- Prevoo ML, van't Hof MA, Kuper HH, van Leeuwen MA, van de Putte LB, van Riel PL. Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995; **38**: 44–48.
- Welsing PM, van Gestel AM, Swinkels HL, Kiemeny LA, van Riel PL. The relationship between disease activity, joint destruction, and functional capacity over the course of rheumatoid arthritis. *Arthritis Rheum* 2001; **44**: 2009–2017.
- Salaffi F, Peroni M, Ferraccioli GF. Discriminating ability of composite indices for measuring disease activity in rheumatoid arthritis: a comparison of the Chronic Arthritis Systemic Index, Disease Activity Score and Thompson's articular index. *Rheumatology (Oxford)* 2000; **39**: 90–96.
- Kushner I. C-reactive protein in rheumatology. *Arthritis Rheum*. 1991; **34**: 1065–1068.
- Brigden ML. Clinical utility of the erythrocyte sedimentation rate. *Am Fam Physician* 1999; **60**: 1443–1450.

- 11 Kullo IJ, Ding K, Shameer K, McCarty CA, Jarvik GP, Denny JC *et al*. Complement receptor 1 gene variants are associated with erythrocyte sedimentation rate. *Am J Hum Genet* 2011; **89**: 131–138.
- 12 Li J, Wang JP, Ghiran I, Cerny A, Szalai AJ, Briles DE *et al*. Complement receptor 1 expression on mouse erythrocytes mediates clearance of *Streptococcus pneumoniae* by immune adherence. *Infect Immun* 2010; **78**: 3129–3135.
- 13 Swift AJ, Collins TS, Bugelski P, Winkelstein JA. Soluble human complement receptor type 1 inhibits complement-mediated host defense. *Clin Diagn Lab Immunol* 1994; **1**: 585–589.
- 14 Weisman HF, Bartow T, Leppo MK, Marsh Jr. HC, Carson GR, Concino MF *et al*. Soluble human complement receptor type 1: in vivo inhibitor of complement suppressing post-ischemic myocardial inflammation and necrosis. *Science* 1990; **249**: 146–151.
- 15 Hyrich KL, Watson KD, Isenberg DA, Symmons DPM. Register BSRB. The British Society for Rheumatology Biologics Register: 6 years on. *Rheumatology (Oxford)* 2008; **47**: 1441–1443.
- 16 Tukaj S, Kotlarz A, Jozwik A, Smolenska Z, Bryl E, Witkowski JM *et al*. Cytokines of the Th1 and Th2 type in sera of rheumatoid arthritis patients; correlations with anti-Hsp40 immune response and diagnostic markers. *Acta Biochimica Polonica* 2010; **57**: 327–332.
- 17 Suk HJ, Ridker PM, Cook NR, Zee RYL. Relation of polymorphism within the C-reactive protein gene and plasma CRP levels. *Atherosclerosis* 2005; **178**: 139–145.
- 18 Plant D, Ibrahim I, Lunt M, Eyre S, Flynn E, Hyrich KL *et al*. Correlation of C-reactive protein haplotypes with serum C-reactive protein level and response to anti-tumor necrosis factor therapy in UK rheumatoid arthritis patients: results from the Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate cohort. *Am J Hum Genet* 2012; **89**: 131–138.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on the The Pharmacogenomics Journal website (<http://www.nature.com/tpj>)

APPENDIX

Members of the Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate (BRAGGSS)

Prof H Gaston: Cambridge University Hospitals NHS Foundation Trust, Addenbrookes Hospital.

Dr D Mulherin, Dr T Price, Dr T Sheeran, Dr V Chalam, Dr S Baskar: Mid Staffordshire NHS Foundation Trust, Cannock Chase Hospital.

Prof P Emery, Prof A Morgan, Dr M Buch, Dr S Bingham: The Leeds Teaching Hospitals NHS Trust, Chapel Allerton Hospital.

Dr S O'Reilly, Dr L Badcock, Dr M Regan, Dr T Ding, Dr C Deighton, Dr G Summers, Dr N Raj: Derby Hospitals NHS Foundation Trust, Derbyshire Royal Infirmary.

Dr R Stevens: Doncaster and Bassetlaw Hospitals NHS Foundation Trust, Doncaster Royal Infirmary.

Dr N Williams: Peterborough and Stamford Hospitals NHS Foundation Trust, Edith Cavell Hospital.

Prof J Isaacs, Dr P Platt, Dr D Walker, Dr L Kay, Dr B Griffiths, Dr W-F Ng, Dr P Peterson, Dr A Lorenzi, Prof H Foster, Dr M Friswell, Dr B Thompson, Dr M Lee, Dr I Griffiths: The Newcastle upon Tyne Hospitals NHS Trust, Freeman Hospital.

Dr A Hassell, Dr P Dawes, Dr C Dowson, Dr S Kamath, Dr J Packham, Dr M Shadforth, Ann Brownfield: University Hospital of North Staffordshire NHS Trust, Haywood Hospital (Stoke-on-Trent).

Dr R Williams: Hereford Hospitals NHS Trust, Hereford County Hospital.

Dr C Mukhtyar: Norfolk & Norwich University Hospital NHS Trust, Norfolk & Norwich University Hospital.

Dr B Harrison, Dr N Snowden, Dr S Naz: Pennine Acute Hospitals NHS Trust, North Manchester General Hospital.

Dr J Ledingham, Dr R Hull, Dr F McCrae, Dr A Thomas, Dr S Young Min, Dr R Shaban, Dr E Wong: Portsmouth Hospitals NHS Trust, Queen Alexandra Hospital.

Dr C Kelly, Dr C Heycock, Dr J Hamilton, Dr V Saravanan: Gateshead Health NHS Trust, Queen Elizabeth Hospital.

Prof G Wilson, Prof D Bax, Dr L Dunkley, Dr M Akil, Dr R Tattersall, Dr R Kilding, Dr S Till, Dr J Boulton, Dr T Tait: Sheffield Teaching Hospitals NHS Trust, Royal Hallamshire Hospital

Dr M Bukhari, Dr J Halsey, Dr L Ottewell: University Hospital of Morcambe Bay NHS Trust, Royal Lancaster Infirmary.

Prof C Buckley, Dr D Situnayake, Dr D Carruthers, Dr K Grindulis, Dr F Khatack, Dr S Elamanchi, Dr K Raza: Sandwell and West Birmingham Hospitals NHS Trust, Sandwell General/City Hospital.

Dr A Filer, Dr R Jubb: University Hospital Birmingham NHS Foundation Trust, Selly Oak Hospital.

Dr R Abernathy: St Helens and Knowsley Hospitals NHS Trust, St Helens Hospital.

Dr M Plant, Dr S Pathare, Dr F Clarke, Dr S Tuck, Dr J Fordham, Dr A Paul: South Tees Hospitals NHS Trust, The James Cook University Hospital, Middlesbrough.

Dr M Bridges: County Durham and Darlington Acute Hospitals NHS Trust, University Hospital of North Durham.

Dr A Hakim, Whipps Cross University Hospital NHS Trust, Whipps Cross University Hospital.

Dr D O'Reilly, Dr V Rajagopal, Dr S Bhagat: The West Suffolk Hospital NHS Trust, West Suffolk Hospital.

Dr C Edwards: Southampton University Hospital NHS Trust, Southampton General Hospital.

Dr P Prouse, Dr R Moitra, Dr D Shawe: Basingstoke & North Hampshire NHS Foundation Trust, Basingstoke & North Hampshire Hospital.

Dr A Bamji: Queen Mary's Sidcup NHS Trust, Queen Mary's Sidcup (QMS) Hospital.

Dr P Klimiuk: Pennine Acute Hospitals NHS Trust, Royal Oldham Hospital.

Dr A Bowden: Pennine Acute Hospitals NHS Trust, Rochdale Infirmary.

Dr W Mitchell: University Hospitals of Morecambe Bay NHS Trust, Furness Hospital.

Prof I Bruce, Prof A Barton, Dr R Gorodkin, Dr P Ho, Dr K Hyrich, Dr W Dixon: Central Manchester University Hospital NHS Foundation Trust, Manchester Royal Infirmary.

Dr A Rai: Worcestershire Acute Hospitals NHS Trust, Worcestershire Royal Hospital.

Prof G Kitas, Dr N Erb, Dr R Klocke, Dr K Douglas, Dr A Pace, Dr R Sandhu, Dr A Whallett: The Dudley Group of Hospitals NHS Foundation Trust, Russells Hall Hospital.

Dr F Birrell: Northumbria Healthcare NHS Foundation Trust, Wansbeck Hospital.

Dr M Allen, Dr K Chaudhuri: University Hospitals of Coventry and Warwickshire NHS Trust, University Hospital (was Walsgrave Hospital).

Dr C Chattopadhyay: Wrightington, Wigan and Leigh Hospitals NHS Foundation Trust, Wrightington Hospital.

Dr J McHale, Dr A Jones, Dr A Gupta, Dr I Pande, Dr I Gaywood, Dr P Lanyon, Dr P Courtney, Dr M Doherty: Nottingham University Hospitals NHS Trust, Nottingham Hospital.

Dr H Chinoy, Prof T O'Neill, Prof A Herrick, Prof A Jones, Dr R Cooper, Dr R Bucknall: Salford Royal NHS Foundation Trust, Hope Hospital.

Dr C Marguerie, Dr S Rigby, Dr N Dunn: South Warwickshire General Hospital NHS Trust, Warwick Hospital.

Dr S Green, Dr A Al-Ansari, Dr S Webber: Weston Area Health NHS Trust, Weston General Hospital.

Dr N Hopkinson, Dr C Dunne, Dr B Quilty: The Royal Bournemouth & Christchurch Hospitals NHS Foundation Trust, Christchurch Hospital.

Dr B Szebenyi: Northern Lincolnshire and Goole Hospitals NHS Foundation Trust, Diana, Princess of Wales Hospital.

Dr M Green, Dr M Quinn, Dr A Isdale, Dr A Brown, Dr B Saleem: York Hospitals NHS Foundation Trust, York District Hospital.

Dr A Samanta, Dr P Sheldon, Dr W Hassan, Dr J Francis, Dr A Kinder, Dr R Neame, Dr A Moorthy: University Hospitals of Leicester NHS Trust, Leicester Royal Infirmary.

Dr W Al-Allaf: The Royal Wolverhampton Hospitals NHS Trust, New Cross Hospital.

Dr A Taggart: Greenpark Healthcare NHS Trust, Musgrave Park Hospital.

Dr K Fairburn: Chesterfield Royal Hospital NHS Foundation Trust, Chesterfield Royal Hospital.

Dr F McKenna: Trafford Healthcare NHS Trust, Trafford General Hospital.

Dr M Green, Dr A Gough, Dr C Lawson: Harrogate and District NHS Foundation Trust, Harrogate District Hospital.

Dr M Piper, Dr E Korendowych, Dr T Jenkinson, Dr R Sengupta, Dr A Bhalla, Prof N. McHugh, Debbie Bond: Royal National Hospital for Rheumatic Diseases NHS Foundation Trust, Bath Hospital.

Prof R Luqmani, Prof B Bowness, Prof P Wordsworth, Dr J David: Oxford Radcliffe Hospitals NHS Trust, John Radcliffe Hospital, Oxford.

Dr W Smith: Milton Keynes Hospital NHS Foundation Trust, Milton Keynes Hospital.

Dr D Mewar, Dr E Tunn, Dr K Nelson, Dr T Kennedy: Royal Liverpool and Broadgreen University Hospitals NHS Trust, Royal Liverpool Hospital.

Dr J Nixon: Countess of Chester Hospital NHS Foundation Trust, Countess of Chester Hospital.

Prof A Woolf, Dr M Davis, Dr D Hutchinson, Dr A Endean: Royal Cornwall Hospitals NHS Trust, Royal Cornwall Hospital.

Dr D Coady, Dr D Wright, Dr C Morley, Dr G Raftery, Dr C Bracewell, Dr L Kidd: City Hospitals Sunderland NHS Foundation Trust, Royal Sunderland Hospital.

Dr I Abbas, Dr C Filer: Stockport NHS Foundation Trust, Stepping Hill Hospital.

Dr G Kallarackal: Kettering General Hospital NHS Foundation Trust, Kettering General Hospital.