



Assessing the Association Between Nutritional Status, Caries, and Gingivitis in Schoolchildren: A Cross-Sectional Study

Global Pediatric Health
Volume 8: 1–7
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2333794X211001237
journals.sagepub.com/home/gph


Mariane C.F. Barbosa, MS¹, Caio L.B. Reis, DDS², Célia M.C.F. Lopes, PhD³,
Isabela R. Madalena, PhD⁴, Erika C. Küchler, PhD⁴ ,
Flares Baratto-Filho, PhD³, Carmen L.M. Storrer, PhD⁵,
Daniela C. Lima, PhD², and Daniela S.B. Oliveira, PhD²

Abstract

Objective. To evaluate if nutritional status is associated with caries and gingivitis in Brazilian schoolchildren. **Material and methods.** Children of both genders, age ranging from 8 to 11 years old, were included in this study. Caries was diagnosed using ICDAS (International System for Detection and Assessment of Carious Lesions) and gingivitis was diagnosed using the Community Periodontal Index. The nutritional status of each child was defined by BMI Z-score calculation. Data on oral health behavior and dietary habit were collected through parent's questionnaires. Parametric analyzes were performed to compare the groups. The established alpha was 5%. **Results.** The sample consisted of 353 schoolchildren: 16 underweight children, 247 eutrophic children, 64 overweight children, and 26 were obese children. Overweight, Obese and Overweight + Obese children presented less cavitated caries lesion than Eutrophic children ($P < .05$). Gingivitis was not associated with nutritional status ($P > .05$). **Conclusion.** Caries was associated with overweight and obesity in Brazilian schoolchildren.

Keywords

caries, gingivitis, nutritional status, obesity, children

Received October 27, 2020. Received revised February 10, 2021. Accepted for publication February 11, 2021.

Introduction

Childhood obesity, overweight and malnourished are serious public health challenges of this century.¹⁻³ Nutrition is necessary for the normal growth and development of children and teenagers.^{4,5} The association between nutrition/diet and many systemic diseases during childhood,^{6,7} as well as the association with oral health,⁸⁻¹⁰ have been widely investigate by many in researchers.¹¹⁻¹⁷

Caries and gingivitis are oral diseases that commonly affect children worldwide. Both oral diseases share some etiological factors with nutrition status indicators, such as obesity and malnourished. These factors include dietary habits, social economical determinants, and presence of dental biofilm.¹¹⁻¹⁴ The common causative relation between oral health and nutritional status deviations has sparked the interest of many researchers.^{5,15-17}

Several studies worldwide have already shown that caries is associated with nutritional status, such as malnourished, overweight, and obesity.¹⁸⁻²² However, the results of studies are inconclusive, as found in many systematic reviews performed in the past decade.^{14,20,23-25}

¹Federal University of Minas Gerais, Belo Horizonte, Brazil

²Federal University of Alfenas, Alfenas, Brazil

³University of the Region of Joinville, Joinville, Brazil

⁴São Paulo University, Ribeirão Preto, Brazil

⁵São Leopoldo Mandic University, Curitiba, São Paulo, Brazil

Corresponding Author:

Daniela Oliveira, Department of Clinic and Surgery, School of Dentistry, Federal University of Alfenas—UNIFAL, Minas Gerais, Rua Gabriel Monteiro da Silva, 700, Centro, Alfenas, Minas Gerais, 37130-001, Brazil.

Email: barrosodaniela@hotmail.com



Several studies also evaluated the association between nutritional status in the etiology of periodontal diseases, including gingivitis.^{12,26,27} The results found in systematic reviews, suggests a positive association between obesity in children and periodontal disease, such as gingivitis.^{26,28,29}

There is also evidence that underweight individuals might have different caries experience than normal weight ones.²⁰ Underweight children are generally malnourished and lacking vitamin A and D, calcium, and phosphorus, which influenced the morphology, chemical composition, and eruption of their teeth, and thus increased the risk of caries.^{20,23} Thus, this study aimed to evaluate if nutritional status is associated with caries and gingivitis in Brazilian schoolchildren.

Materials and Methods

This is a cross-sectional study that evaluated the association between caries, gingivitis, and nutritional status. The STROBE guideline was followed to conduct this manuscript (Vandenbroucke et al., 2007). The Local Human Ethics Committee (#78568217.7.0000.5142) approved this study. Informed written consent was taking from all legal guardians and assent document from all children.

Participants

This school based-study evaluated school-age-children from 4 public schools located in Alfenas city. Alfenas is a medium-sized city located in the southeast region of Brazil.³⁰ The sample represents the total of schoolchildren, aging from 8 to 11 years, both genders, from these schools and previously was described in Reis et al.³¹ Children with systemic and cognitive disorders were excluded.

Anamnesis and Determination of Oral Clinical Aspects

The caregiver or parent of each child answered the anamnesis regarding the demographic, diet, and dental health habits information.

The physical and dental examination were made in each included child. Intra-examiner training and calibration were conducted before the study (kappa coefficient=0.87). A one trained dentist made the dental exams. The dental evaluation was made in the schools, under natural light, using cotton rollers, gauze, standard mouth mirrors, and ballpoint probe according to the WHO (World Health Organization). The dental exam

included the evaluation of bleeding on probe, dental plaque (visible biofilm), and caries. The Community Periodontal Index was used to evaluate the gingival bleeding (bleeding on probe) according to the WHO.³²

Each child was classified as “yes” or “no” for bleeding on probe. The presence of dental plaque was performed using the Silness and Løe³³ index. The scores ranges from 0 (absence of biofilm) to 3 (biofilm abundant on the surface of the tooth) and 6 sextants were examined.

To diagnosis caries, ICDAS (International System for Detection and Assessment of Carious Lesions) criteria was adopted. The caries lesions were diagnosed during visual examination and presents a scale that ranges from 0 to 6.³⁴ Caries was categorized according to the severity: ICDAS₀ versus ICDAS₁₋₆ (caries); or ICDAS₁₋₂ (non-cavitated caries lesion) versus ICDAS₃₋₆ (cavitated caries lesion) as previously described in Reis et al.³¹

Determination of the Nutritional Status

The anthropometric data (height and weight) were recorded on the day of the dental examination. The heights were evaluated and determined in meters, while the weights were evaluated and determined in kilograms using a weighing machine. Children were wearing light clothes and no shoes. Therefore, the nutritional status was calculated for each child using the Body Mass Index (BMI) *z*-score calculator (<http://zscore.research.chop.edu/index.php>). Height, weight, age, and gender were used as variables in the BMI *z*-score formula and the classifications were performed as recommended by the WHO¹ (Table 1).

Statistical Analysis

GraphPad Prism 5.0 (Graph-Pad, San Diego, CA, EUA) was used for statistical analysis. Caries and gingivitis were analyzed as categorical data. Chi-square and One-way ANOVA with Tukey's post-test were used to compare the groups ($\alpha = 5\%$).

Results

Three hundred fifty-three school-age individuals were included, 48.2% were males and 51.8% were females. The sample consisted of 16 underweight children, 247 eutrophic children, 64 overweight children, and 26 obese children. The characteristics, dietary and dental hygiene aspects according to the status nutritional, gingivitis, and caries are presented in Table 2. Dental plaque

Table 1. Parameters for Classification of Nutritional Status According to the World Health Organization.

Nutritional status	BMI z score
Underweight (malnourished)	<3 percentiles
Eutrophic (normal weight children)	≥3 and ≤ 85th percentiles
Overweight	>85th and ≤ 97th percentiles
Obese	>97th percentiles

Disponible in: <https://www.who.int/childgrowth/standards/technicalreport/en/>.

was associated with overweight, caries and gingivitis ($P < .05$). Use of dental floss was associated with gingivitis ($P = .03$). Caries and gingivitis distribution are presented in the Figure 1.

Table 3 presents caries and gingivitis experience distribution according to nutritional status. Overweight ($P = .04$; $OR = 0.56$; $95\% CI = 0.32-0.99$) and Obese ($P = .05$; $OR = 0.42$; $95\% CI = 0.15-1.0$) children presented less cavitated caries lesion (ICDAS₀₋₂ vs ICDAS₃₋₆) than Eutrophic children. Caries experience also was less in Overweight + Obese children in the 2 cut-offs (ICDAS₀₋₂ vs ICDAS₃₋₆; $P = .01$; $OR = 0.51$; $95\% CI = 0.31-0.87$. ICDAS₀ vs ICDAS₁₋₆; $P = .05$; $OR = 0.62$; $95\% CI = 0.37-1.0$). Nutritional status was associated with caries and gingivitis experience ($P < .05$).

Discussion

Obesity, malnourished, caries, and periodontitis are serious public health problems worldwide. A possible association between nutritional status and caries,²⁸ as well as nutritional status and periodontal diseases^{26,29} have been already suggested. These conditions share similar behavioral habits as etiological conditions.³⁵ The health burden from nutritional changes is the driving factor behind several researches regarding the impact of nutritional status conditions on children's oral health and development.^{36,37} For this reason, we decided to evaluate if variations in the nutritional status are associated with caries and gingivitis in Brazilian schoolchildren.

Our study observed that overweight and obese children had lower caries experience than normal weight children. Our results are in agreement with the finds observed in the previous studies with Brazilians from other regions of the country.³⁸ It is possible that caries is associated with overweight and obesity due to social economic factors. Families with higher income might have more access to sweets and also to dental treatment. Other possible theory suggested for the association between low caries and high BMI may be due to the

Table 2. Distribution of the Characteristics of the Study Subjects among Dental Caries, Gingivitis, and Nutritional Status.

Characteristics	Total of subjects	Eutrophic	Underweight	P*	Overweight	P*	Obese	P*	Health (ICDAS ₀)	Dental caries (ICDAS ₁₋₆)	P	Healthy and non-cavitated caries (ICDAS ₀₋₂)	Cavitated caries lesion (ICDAS ₃₋₆)	P	Non gingivitis	Gingivitis	P
n (%)	353 (100)	247 (69.9)	16 (4.53)		64 (18.1)		26 (7.36)		189 (53.5)	164 (46.5)		203 (57.5)	150 (42.5)		206 (58.3)	147 (41.7)	
Age (mean, SD)	8.9 (0.89)	8.8 (0.89)	9.43 (0.81)	.006	9.03 (0.88)	.06	8.65 (0.89)	.41	8.95 (0.92)	8.79 (0.84)	.87	8.92 (0.81)	8.83 (0.95)	.78	8.91 (0.88)	8.84 (0.90)	.89
Gender (n, %)																	
Male	170 (48.2)	116 (47.0)	8 (50.0)	.81	29 (45.3)	.81	17 (65.4)	.07	92 (48.7)	78 (47.6)	.83	98 (48.3)	72 (48.0)	.81	96 (46.6)	74 (50.3)	.48
Female	183 (51.8)	131 (53.0)	8 (50.0)		35 (64.7)		9 (34.6)		97 (51.3)	86 (52.4)		105 (51.7)	78 (52.0)		110 (53.4)	73 (49.7)	
Biofilm index (mean, SD)	1.79 (0.30)	1.80 (0.28)	1.85 (0.29)	.49	1.72 (0.29)	.04	1.77 (0.28)	.60	1.75 (0.30)	1.83 (0.26)	.007	1.74 (0.30)	1.85 (0.24)	.002	1.73 (0.31)	1.86 (0.23)	.0001
Ingestion of sweets between meals (n, %)																	
Yes	132 (38.5)	6 (33.3)	99 (41.1)	.51	19 (31.1)	.86	9 (37.5)	.78	65 (35.3)	67 (42.1)	.09	69 (35.0)	63 (44.1)	.06	76 (38.4)	56 (38.6)	.95
No	11 (61.5)	12 (66.7)	142 (58.9)		42 (68.9)		15 (62.5)		119 (64.7)	92 (57.9)		128 (65.0)	83 (56.9)		122 (61.6)	89 (61.4)	
Brush teeth before sleep (n, %)																	
Yes	257 (76.9)	172 (74.1)	14 (93.3)	.09	49 (79.0)	.42	22 (88.0)	.12	135 (75.8)	122 (78.2)	.30	147 (76.6)	110 (77.5)	.30	153 (77.7)	104 (75.9)	.86
No	77 (23.1)	60 (25.9)	1 (6.7)		13 (21.0)		3 (12.0)		43 (24.2)	34 (21.8)		45 (23.4)	32 (22.5)		44 (22.3)	33 (24.1)	
How many times brush teeth per day (n, %)																	
1 time	43 (12.6)	33 (13.8)	1 (6.3)	Ref.	8 (12.9)	Ref.	1 (4.0)	Ref.	20 (11.0)	23 (14.4)	Ref.	22 (11.2)	21 (14.4)	Ref.	24 (12.2)	19 (13.1)	Ref.
2 times	136 (39.8)	93 (38.9)	7 (43.7)	.38	25 (37.1)	.81	11 (44.0)	.17	74 (40.7)	62 (38.7)	.36	76 (38.8)	60 (41.1)	.58	76 (38.6)	60 (41.4)	.67
3 or more times	163 (47.7)	113 (47.3)	8 (50.0)	.41	29 (46.8)	.89	13 (52.0)	.17	88 (48.3)	75 (46.9)	.38	98 (50.0)	65 (44.5)	.28	97 (49.2)	66 (45.5)	.66
Use of dental floss (n, %)																	
Yes	218 (63.0)	149 (61.6)	11 (68.8)	.56	40 (64.5)	.66	18 (69.2)	.44	124 (67.0)	94 (58.4)	.09	134 (67.3)	84 (57.2)	.052	136 (67.7)	82 (56.6)	.03
No	128 (37.0)	93 (38.4)	5 (31.2)		22 (35.5)		8 (30.8)		61 (67.0)	67 (41.6)		65 (32.7)	63 (42.8)		63 (32.3)	63 (43.4)	

Abbreviations: SD, standard deviation; Ref, reference (control). Bold forms mean statistical significance difference ($P < .05$).

*In comparison with eutrophic group.




Figure 1. Frequency of children per caries and gingivitis groups. (A) Frequency of ICDAS₀ and ICDAS₁₋₆. (B) Frequency of ICDAS₀₋₂ and ICDAS₃₋₆. (C) Frequency of No Gingivitis and Gingivitis.

Table 3. Association between Nutritional Status and Caries Experience/Gingivitis.

Groups	Eutrophic		Underweight		Overweight		Obese		Overweight + Obese				
	n (%)	n (%)	P-value	OR (95% CI)	n (%)	P-value	OR (95% CI)	n (%)	P-value	OR (95% CI)			
Non-cavitated caries lesion vs Cavitated caries lesion													
ICDAS ₀₋₂	132 (53.4)	8 (50.0)	.78	1.14 (0.41-3.15)	43 (67.2)	.04	0.56 (0.32-0.99)	19 (73.1)	.05	0.42 (0.15-1.0)	62 (68.9)	.01	0.51 (0.31-0.87)
ICDAS ₃₋₆	115 (46.6)	8 (50.0)			21 (32.8)			7 (26.9)			28 (31.1)		
Health vs Caries													
ICDAS ₀	125 (50.6)	8 (50.0)	.96	1.02 (0.37-2.81)	38 (59.4)	.21	0.70 (0.40-1.22)	18 (69.2)	.07	0.45 (0.20-1.11)	56 (62.2)	.05	0.62 (0.37-1.0)
ICDAS ₁₋₆	122 (49.4)	8 (50.0)			26 (40.6)			8 (30.8)			34 (37.8)		
Health vs Gingivitis													
Non gingivitis	141 (57.0)	8 (50.0)	.57	0.7 (0.2-2.0)	37 (57.8)	.91	1.0 (0.5-1.7)	20 (76.9)	.06	0.39 (0.16-0.96)	57 (63.3)	.30	0.77 (0.46-1.27)
Gingivitis	106 (43.0)	8 (50.0)			27 (42.2)			6 (23.1)			33 (36.7)		

All comparisons were performed with Eutrophic group (Eutrophic was the reference for all analysis). Bold forms mean statistical significance difference ($P < .05$). OR 95% CI: odds ratios with 95% confidence intervals.

high consumption of high-fat diets, which are positively associated with obesity, but it is not with caries.³⁹ Besides that, a possible protective effect of fat against caries has been previously suggested in the literature.^{40,41} According to the national data evaluating nutrient consumption, the caloric participation of lipids in the Southeast Brazilian region is higher compared to others Brazilian regions and the national mean.⁴² Although our study did not observe a statistical association between diet and nutritional status, this could be due the limitation of our study that evaluated only the self-reported ingestion of sweets between meals, which is a risk factor for caries.

Another possible theory for this result was suggests that severe untreated caries affects eating ability,^{43,44} however, this would lead to underweight children. This hypothesis is supported by the study of Duijster et al⁴⁵ in Filipino children, which showed that treatment of severely carious lesion in underweight children was associated with significant weight gain. Our study did not observe

statistical association between underweight and caries experience; however, this could be due to the small sample size of underweight children in our population.

Periodontal diseases, including gingivitis, are infectious and inflammatory conditions characterized by a modification in the microbial ecology of biofilms.⁴⁶ Among children and teenagers, the most prevalent periodontal disease is biofilm-induced gingivitis,⁴⁷ which is a precondition for periodontitis, however not all cases of gingivitis will progress to periodontitis.^{29,48} In this study, the nutritional status was not associated with gingivitis. These results contrast with a recent published systematic review and meta-analyzes, in which obesity demonstrated a risk factor for gingivitis among children and adolescents.²⁹ However, the published literature still has no consensus on the relationship between nutritional status and periodontal diseases, including gingivitis in the pediatric population.

One possible limitation of our study is the fact that a full mouth periodontal examination was not performed,

however, the Community Periodontal Index was used as recommended by the World Health Organization. Therefore, it is possible that the presence of gingivitis has been underestimated.

One important aspect this study is that, different from the most previous studies evaluating BMI and caries, that used the sum of decayed, missing and filled teeth (DMFT) criteria for the diagnosis of caries, our study used ICDAS as a diagnostic method. ICDAS is a more elaborate scale and could produce more accurate results, as supported by previous authors.²³

Conclusions

Caries was associated with overweight and obesity, while gingivitis was not associated with nutritional status.

Author Contributions

ECK, DSBO, FBF and CLMS designed the data. MCFB, DCL and DSBO examined and collected the sample. MCFB, CLBR and IRM organized the data. CLBR, CMCFL and ECK analyzed the data. MCFBB, CLBR, IRM, ECK and DSBO interpreted the results and wrote the manuscript. All authors read and approved the final version of the manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) Finance Code 001 and also by Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG).

ORCID iD

Érika Küchler  <https://orcid.org/0000-0001-5351-2526>

Supplemental Material

Supplemental material for this article is available online.

References

- World Health Organization. Multicentre Growth Reference Study Group. WHO Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index- for-Age: Methods and Development. World Health Organization, 2006.
- Macek MD, Mitola DJ. Exploring the association between overweight and dental caries among US children. *Pediatr Dent*. 2006;28:375-380.
- Evangelista SS, Vasconcelos KRF, Xavier TA, et al. Timing of permanent tooth emergence is associated with overweight/obesity in children from the Amazon Region. *Braz Dent J*. 2018;29:465-468 doi:10.1590/0103-6440201802230
- Priyadarshini HR, Hiremath SS, Puranik M. Prevalence of early childhood caries among preschool children of low socioeconomic status in Bangalore city, India. *J Int Soc Prev Community Dent*. 2011;1:27-30. doi:10.4103/2231-0762.86384
- Krishna HNS, Manaswini E, Kumar VY, et al. Association between nutritional status and early childhood caries in Indian children. *J Int Soc Prev Community Dent*. 2017;7:131. doi:10.4103/jispcd.JISPCD_25_17
- Ritchie CS, Joshipura K, Hung HC, Douglass CW. Nutrition as a mediator in the relation between oral and systemic disease: associations between specific measures of adult oral health and nutrition outcomes. *Crit Rev Oral Biol Med*. 2002;13:291-300. doi:10.1177/154411130201300306
- Adeniyi AA, Oyapero OA, Ekekezie OO, Braimoh MO. Dental caries and nutritional status of school children in Lagos, Nigeria—A preliminary survey. *J West Afr Coll Surg*. 2016;6:15-38.
- Kulkarni V, Bhatavadekar NB, Uttamani JR. The effect of nutrition on periodontal disease: a systematic review. *J Calif Dent Assoc*. 2014;42:302-311.
- Gonçalves JDA, Moreira EAM, Rauen MS, Rossi A, Borgatto AF, et al. Associations between caries experience, nutritional status, oral hygiene, and diet in a multi-generational cohort. *Pediatr Dent*. 2016;38:203-211.
- Dimaisip-Nabuab J, Duijster D, Benzian H, et al. Nutritional status, dental caries and tooth eruption in children: a longitudinal study in Cambodia, Indonesia and Lao PDR. *BMC Pediatr*. 2018;18:300. doi:10.1186/s12887-018-1277-6
- Trikaliotis A, Boka V, Kotsanos N, Karagiannis V, Hassapidou M. Short communication: Dmfs and BMI in preschool Greek children. An epidemiological study. *Eur Arch Paediatr Dent*. 2011;12:176-178. doi:10.1007/bf03262802
- Nascimento GG, Seerig LM, Vargas-Ferreira F, Correa FO, Leite FR, Demarco FF. Are obesity and overweight associated with gingivitis occurrence in Brazilian schoolchildren? *J Clin Periodontol*. 2013;40:1072-1078. doi:10.1111/jcpe.12163
- Jepsen S, Blanco J, Buchalla W, et al. Prevention and control of dental caries and periodontal diseases at individual and population level: consensus report of group 3 of joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J Clin Periodontol*. 2017;44:S85-S93. doi:10.1111/jcpe.12687
- Shivakumar S, Srivastava A, Shivakumar GC. Body mass index and dental caries: a systematic review. *Int J Clin Pediatr Dent*. 2018;11:228. doi:10.5005/jp-journals-10005-1516
- Yang F, Zhang Y, Yuan X, et al. Caries experience and its association with weight status among 8-year-old children in Qingdao, China. *J Int Soc Prev Community Dent*. 2015;5:52-58. doi:10.4103/2231-0762.151978

16. Silva RAB, Barreiros D, Oliveira S, da Silva LA, Nelson-Filho P, Küchler EC. Association between Body Mass Index and Caries Experience in Brazilian children and adolescents. *J Dent Child*. 2016;83:146-151.
17. Farsi DJ, Elkhodary HM, Merdad LA, et al. Prevalence of obesity in elementary school children and its association with dental caries. *Saudi Med J*. 2016;37:1387-1394. doi:10.15537/smj.2016.12.15904
18. Shailee F, Sogi GM, Sharma KR. Association between dental caries and body mass index among 12 and 15 years school children in Shimla, Himachal Pradesh. *J Adv Oral Res*. 2013;4:6-10. doi:0.1177/2229411220130102
19. Parkar SM, Chokshi M. Exploring the association between dental caries and body mass index in public school children of Ahmedabad city, Gujarat. *SRM J Res Dent Sci*. 2013;4:101-105. doi:10.4103/0976-433X.121633
20. Chen D, Zhi Q, Zhou Y, Tao Y, Wu L, Lin H. Association between dental caries and BMI in children: a systematic review and meta-analysis. *Caries Res*. 2018;52:230-245. doi:10.1159/000484988
21. Li LW, Wong HM, Gandhi A, McGrath CP. Caries-related risk factors of obesity among 18-year-old adolescents in Hong Kong: a cross-sectional study nested in a cohort study. *BMC Oral Health*. 2018;18:188. doi:10.1186/s12903-018-0657-5
22. Hamasha AA, Alsolaim AA, Alturki HA, Alaskar LA, Alshunaiber RA, Aldebasi WT. The relationship between body mass index and oral health status among Saudi adults: a cross-sectional study. *Community Dent Health*. 2019;36:217-222. doi:10.1922/CDH_4361Hamasha06
23. Hooley M, Skouteris H, Millar L. The relationship between childhood weight, dental caries and eating practices in children aged 4–8 years in Australia, 2004–2008. *Pediatr Obes*. 2012;7:461-470. doi:10.1111/j.2047-6310.2012.00072.x
24. Silva AER, Menezes AMB, Demarco FF, Vargas-Ferreira F, Peres MA. Obesity and dental caries: systematic review. *Rev Saude Publica*. 2013;47:799-812. doi:10.1590/S0034-8910.2013047004608
25. Hayden C, Bowler JO, Chambers S, et al. Obesity and dental caries in children: a systematic review and meta-analysis. *Community Dent Epidemiol*. 2013; 41:289-308. doi:10.1111/cdoe.12014
26. Nascimento GG, Leite FR, Do L, et al. Is weight gain associated with the incidence of periodontitis? A systematic review and meta-analysis. *J Clin Periodontol*. 2015; 42:495-505. doi:10.1111/jcpe.12417
27. Patiño-Marín N, Zavala-Alonso NV, Martínez-Castañón GA, et al. Association between dental hygiene, gingivitis and overweight or the risk of overweight in primary teeth of 4-and 5-year-old preschoolers in México. *Int J Dent Hyg*. 2018;16:411-418. doi:10.1111/idh.12345
28. Li LW, Wong HM, Sun L, Wen YF, McGrath CP. Anthropometric measurements and periodontal diseases in children and adolescents: a systematic review and meta-analysis. *Adv Nutr*. 2015;6:828-841. doi:10.3945/an.115.010017
29. Martens L, De Smet S, Yusof MY, Rajasekharan S. Association between overweight/obesity and periodontal disease in children and adolescents: a systematic review and meta-analysis. *Eur Arch Paediatr Dent*. 2017;18: 69-82. doi:10.1007/s40368-017-0272-1
30. Instituto Brasileiro de Geografia e Estatística. Diretoria de Pesquisas, Coordenação de População e Indicadores Sociais, Estimativas da população residente com data de referência. 2019. Accessed October 3, 2020. <https://cidades.ibge.gov.br/brasil/mg/alfenas/panorama>
31. Reis CLB, Barbosa MCF, Machado BMDS, et al. Genetic polymorphisms in interleukin-6 and interleukin-1-beta were associated with dental caries and gingivitis. *Acta Odontol Scand*. 2020;1:1-7.
32. World Health Organization. *World Health Organization: Oral Health Surveys, Basic Methods*. WHO; 1997.
33. Silness J, Loe H. Periodontal disease in pregnancy II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand*. 1964;22:121-135. doi:10.3109/00016356408993968
34. Jablonski-Momeni A, Stachniss V, Ricketts DN, Heinzl-Gutenbrunner M, Pieper K. Reproducibility and accuracy of the ICDAS-II for detection of occlusal caries in vitro. *Caries Res*. 2008;42:79-87. doi:10.1159/000113160
35. Cinar AB, Murtomaa H. Interrelation between obesity, oral health and life-style factors among Turkish school children. *Clin Oral Investig*. 2011;15:177-184. doi:10.1007/s00784-009-0368-z
36. da Silva RAB, Barreiros D, Oliveira S, da Silva LA, Nelson-Filho P, Küchler EC. Association between body mass index and caries experience in Brazilian children and adolescents. *J Dent Child (Chic)*. 2016; 83:146-151.
37. Fonseca MA. Malnutrition and oral health in children. *Curr Oral Health Rep*. 2017;4:92-96. doi:10.1007/s40496-017-0130-6
38. Lock NC, Susin C, Brusius CD, Maltz M, Alves LS. Obesity and dental caries among South Brazilian schoolchildren: a 2.5-year longitudinal study. *Braz Oral Res*. 2019;33:e56. doi:10.1590/1807-3107bor-2019.vol33.0056
39. Emmett PM, Jones L. Diet, growth, and obesity development throughout childhood in the Avon longitudinal study of parents and children. *Nutr Rev*. 2015;73:175-206. doi:10.1093/nutrit/nuv054
40. Brudevold F, Kashket S, Kent RL Jr. The effect of sucrose and fat in cookies on salivation and oral retention in humans. *J Dent Res*. 1990;69:1278-1282. doi:10.1177/00220345900690061101
41. Giacaman RA, Muñoz-Sandovalm C. Cariogenicity of different commercially available bovine milk types in a biofilm caries model. *Pediatr dent*. 2014;36:1E-6E.
42. Instituto Brasileiro de Geografia e Estatística. Pesquisa de orçamentos familiares 2008-2009: análise do consumo pessoal de alimentos no Brasil. Instituto Brasileiro de Geografia e Estatística; 2011. Accessed April 16, 2020. <https://biblioteca.ibge.gov.br/visualizacao/livros/liv50063.pdf>

43. Sheiham A. 2006. Dental caries affects body weight, growth and quality of life in pre-school children. *Br Dent J.* 2006;201:625-626. doi:10.1038/sj.bdj.4814259
44. Paisi M, Kay E, Bennett C, et al. Body mass index and dental caries in young people: a systematic review. *BMC Pediatr.* 2019;19:122. doi:10.1186/s12887-019-1511-x
45. Duijster D, Sheiham A, Hobdell MH, Itchon G, Monse B. Associations between oral health-related impacts and rate of weight gain after extraction of pulpally involved teeth in underweight preschool Filipino children. *BMC Public Health.* 2013;13:533. doi:10.1186/1471-2458-13-533
46. Suresh S, Mahendra J. Multifactorial relationship of obesity and periodontal disease. *J Clin Diagn Res.* 2014;8:ZE01-ZE03. doi:10.7860/JCDR/2014/7071.4227
47. Lopez R, Fernandez O, Baelum V. Social gradients in periodontal diseases among adolescents. *Community Dent Oral Epidemiol.* 2006;34:184-196. doi:10.1111/j.1600-0528.2006.00271.x
48. Chapple ILC, Bouchard P, Cagetti MG, et al. Interaction of lifestyle, behaviour or systemic diseases with dental caries and periodontal diseases: consensus report of group 2 of the joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J Clin Periodontol.* 2017;44:S39-S51. doi:10.1111/jcpe.12685