

Debate

Open Access

Lysophosphatidic acid, LPA: a bad boy becomes good

Lygia T Budnik*

Address: Institute for Hormone and Fertility Research, University of Hamburg, Centrum for Innovative Medicine, Falkenried 88, D-20251 Hamburg, Germany

Email: Lygia T Budnik* - Budnik@IHF.de

* Corresponding author

Published: 23 April 2003

Received: 17 March 2003

Reproductive Biology and Endocrinology 2003, **1**:37

Accepted: 23 April 2003

This article is available from: <http://www.RBEj.com/content/1/1/37>

© 2003 Budnik; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

The biologically active growth factor LPA has gained broad interest after it was proposed as a possible marker for ovarian cancer [1]. Although much is known about LPA signaling, its physiological and pathological relevance remains unclear. Recent data from Tokumura et al. [2–4] may change our perception of LPA and cause LPA researches to focus on its physiological aspects. LPA functions may also be reflected in the expression of LPA producing enzyme lysophospholipase D (LPLD), which is increased in serum and correlates with the progress of pregnancy [2,3]. Recent exciting data, from the same and other groups, has recognized this enzyme as autotaxin, a cell motility stimulating phosphodiesterase implicated in tumor progression [3,5,6]. At first glance, it may appear contradictory that a factor associated with tumor progression may serve normal physiological functions [7]. To appreciate this better the normal physiology of reproductive systems should be contemplated. Only a few tissues are known to grow under normal physiological conditions and notable among these are the ovary, endometrium, placenta and the growing syncytiotrophoblast. Growth and differentiation of the ovary and endometrium (/placenta) is accompanied by processes which have been likened to inflammation and wound-healing. Unfortunately, these aspects of LPA biology appear to focus the interest of reproductive biologists and gynecologists only, and not the rest of the scientific community. Not only, are reproductive systems part of normal physiological function, but also they provide amenable models for the study of controlled growth and differentiation. Maybe we should accept that the pathological role of LPA could be a reflection of its physiological functions. In this perspective, Autotaxin may presumably take center stage in future drug discovery associated not only with cancer treatment but also with fertility control.




Figure 1
Secretory ecto-phospholipases produce biologically active lipid, lysophosphatidic acid (LPA). LPA should be released into extracellular fluid to signal through its specific G-protein coupled receptor (LP-R) [7]. Secretory ecto-enzymes like Lysophospholipase D (LPLD) or secretory-type Phospholipase A2 (sPLA2) act on phospholipid substrates available in extracellular leaflets of the plasma membrane (and can supply LPA directly to its receptors).

References

1. Fang X, Schummer M, Mao M, Yu S, Tabassam FH, Swaby R, Hasegawa Y, Tanyi JL, LaPushin R, Eder A, Jaffe R, Erickson J and Mills GB **Lysophosphatidic acid is a bioactive mediator in ovarian cancer** *Biochem Biophys Acta* 2002, **1582**:257-264
2. Tokumura A, Kanaya Y, Miyake M, Yamano S, Irahara M and Fukuzawa K **Increased production of bioactive lysophosphatidic acid by**

serum lysophospholipase D in human pregnancy *Biol Reprod* 2002, **67**:1386-1392

3. Tokumura A, Majima E, Kariya Y, Tominaga K, Kogure K, Yasuda K and Fukuzawa K **Identification of human plasma lysophospholipase D, a lysophosphatidic acid-producing enzyme, as autotaxin, a multifunctional phosphodiesterase** *J Biol Chem* 2002, **277**:39436-39442
4. Tokumura A **Physiological and pathophysiological roles of lysophosphatidic acids produced by secretory lysophospholipase D in body fluids** *Biochim Biophys Acta* 2002, **1582**:18-25
5. Umezu-Goto M, Kishi Y, Taira A, Hama K, Dohmae N, Takio K, Yamori T, Mills GB, Inoue K, Aoki J and Arai H **Autotaxin has lysophospholipase D activity leading to tumor cell growth and motility by lysophosphatidic acid production** *J Cell Biol* 2002, **158**:227-233
6. Moolenaar WH **Lysophospholipids in the limelight: autotaxin takes center stage** *J Cell Biol* 2002, **158**:197-199
7. Budnik LT and Mukhopadhyay AK **Lysophosphatidic acid and its role in reproduction** *Biol Reprod* 2002, **66**:859-865

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

