

Aberrant differentiation of *Tsc2*-deficient teratomas associated with activation of the mTORC1-TFE3 pathway

HARUNA KAWANO^{1,2}, YOSHITAKA ITO^{2,3}, FUMIO KANAI⁴, ERI NAKAMURA⁴, NORIHIRO TADA⁴, SETSUO TAKAI⁵, SHIGEO HORIE¹, TOSHIYUKI KOBAYASHI² and OKIO HINO²

Departments of ¹Urology, ²Molecular Pathogenesis and ³Neurosurgery, and ⁴Laboratory of Genome Research, Research Institute for Diseases of Old Age, Juntendo University Graduate School of Medicine, Tokyo 113-8431;

⁵Department of Clinical Radiology, Faculty of Health Sciences, Hiroshima International University, Hiroshima 724-0695, Japan

Received May 24, 2015; Accepted July 21, 2015

DOI: 10.3892/or.2015.4254

Abstract. The model animal of renal cell carcinoma (RCC), the Eker rat, has a germline mutation in the tuberous sclerosis 2 (*Tsc2*) gene. Heterozygous mutants develop RCCs by second hit in the wild-type *Tsc2* allele, whereas homozygous mutants are embryonic lethal. In the present study, a new cell differentiation model was developed to study the mechanism of *Tsc2* mutation-associated pathogenesis by generating *Tsc2*-deficient embryonic stem cells (ESCs) from Eker rats. *Tsc2*^{+/+}, *Tsc2*^{+/-} and *Tsc2*^{-/-} ESCs were all capable of generating three germ layers: mesoderm, ectoderm, and endoderm. Interestingly, epithelial tumor-like abnormal ductal structures were reproducibly observed in *Tsc2*^{-/-} teratomas from different ESC lines. Immunohistochemical analysis revealed that mammalian target of rapamycin complex 1 (mTORC1) signaling was activated in abnormal ducts of *Tsc2*^{-/-} teratomas, on the basis of positive staining for p-S6 and p-4EBP1. In these abnormal ducts, expression levels of epithelial markers (i.e., megalin and cubilin) and the cytoplasmic localization of E-cadherin and β -catenin were similar to those in Eker rat RCCs. Moreover, a transcription factor regulated by mTORC1, named TFE3, was located in the nuclei of abnormal ducts and Eker rat RCCs. As a negative regulator of ESC differentiation, TFE3 may result in tissue-specific differentiation defects related to tumorigenesis in Eker rats and *Tsc2*^{-/-} teratomas. The present study suggests that ESCs derived from Eker rats constitute a novel experimental tool with which to analyze differentiation defects and cell-type specific pathogenesis associated with *Tsc2* deficiency.

Introduction

The Eker rat is a useful animal model with which to study renal cell carcinoma (RCC) (1). Spontaneous tumors develop in the kidney due to a germline mutation in the tuberous sclerosis 2 (*Tsc2*) gene (2,3). Homozygous mutants are embryonic lethal during midgestation (equivalent to mouse E9.5-E13.5) (4). In contrast, heterozygous mutants develop bilateral multicentric renal cell carcinomas within one year after birth (5). The development of multistage renal carcinogenesis can be monitored at the histological level (6). *TSC2* (encoding tuberin) is a tumor-suppressor gene identified as a causative gene of TSC as well as *TSC1* (encoding hamartin) (7-9). These products form a complex that inhibits the mammalian target of rapamycin complex 1 (mTORC1), a serine/threonine protein kinase essential for the regulation of cell growth and proliferation (10). Mammalian cells express two functionally distinct mTOR complexes: mTORC1 and mTORC2. mTORC1 contains mTOR, Raptor and LST8 as primary subunits and is inhibited by rapamycin (11). In TSC-associated tumors, the loss of *TSC1* or *TSC2* induces mTORC1-dependent phosphorylation of p70S6 kinase, ribosomal protein S6 and 4EBP1 (12). On the other hand, mTORC2 contains Rictor, LST8 and Sin1 as primary subunits. This complex functions as a rapamycin-insensitive regulator of the cytoskeleton and cell survival through Akt phosphorylation (13).

Few previous studies have explored the cancer-related cellular mechanisms triggered by the loss of tumor-suppressor genes using embryonic stem cells (ESCs) such as *Apc*-mutated ESCs (14). To establish a new research model for the analysis of renal carcinogenesis, we developed ESCs from Eker rat blastocysts (15). We successfully established *Tsc2*^{+/+}, *Tsc2*^{+/-} and *Tsc2*^{-/-} Eker rat ESCs with pluripotency. Furthermore, in the teratoma formation assay, which is an important process for pluripotent stem cells (PSCs) to prove differentiation ability, epithelial tumor-like abnormal ductal structures resembling Eker rat RCCs were identified in *Tsc2*^{-/-} teratomas. Immunohistochemical analysis revealed the activation of mTORC1 signaling in *Tsc2*^{-/-} teratomas, particularly in abnormal ductal structures, which was suppressed by rapamycin treatment. We suggested that the appearance of these abnormal structures indicates a pathogenic mechanism

Correspondence to: Professor Okio Hino or Dr Toshiyuki Kobayashi, Department of Molecular Pathogenesis, Juntendo University Graduate School of Medicine, Hongo 3-1-3, Bunkyo-ku, Tokyo 113-8431, Japan
E-mail: ohino@juntendo.ac.jp
E-mail: koba396@juntendo.ac.jp

Key words: Eker rat, embryonic stem cells, renal cell carcinoma, teratomas, tuberous sclerosis complex

related to renal tumorigenesis in Eker rats. These teratomas constitute a new research tool for the analysis of tissue-specific tumorigenesis by suppression of mTORC1 signaling.

Materials and methods

Animal studies. All animal experiments were performed according to protocols approved by the Animal Care Committee of Juntendo University of Medicine (Approval no. 250105). All surgeries were performed under isoflurane anesthesia and measures were taken to minimize animal suffering.

Cell culture. Rat ESCs were cultured on mitomycin C-treated mouse embryonic fibroblasts in N2B27-2i medium (1:1 ratio of 2i medium:N2B27 medium; Axon Medchem, Groningen, The Netherlands) containing 1,000 U/ml rat leukemia inhibitory factor (LIF; ESGRO Millipore, Bedford, MA, USA), 3 μ M GSK3 β inhibitor CHIR99021 (Axon), and 1 μ M MEK inhibitor PD0325901 (Axon).

Teratoma formation. A total of 1.5×10^6 *Tsc2*^{+/+}, *Tsc2*^{+/-} or *Tsc2*^{-/-} Eker ESCs suspended in Matrigel were subcutaneously injected into non-obese diabetic/severe combined immunodeficiency (NOD/SCID) mice. After 5 weeks, 1.5 mg/kg body weight of rapamycin (Sigma-Aldrich, St. Louis, MO, USA), or vehicle alone (vehicle control), was intraperitoneally injected every other day, for a total of three injections. Mice were sacrificed the day after the last injection.

Immunohistochemistry. Immunohistochemistry was carried out using a standard method using formalin-fixed, paraffin-embedded tissues. For chromogenic analysis, EnVision+ System HRP-labeled polymer for rabbit or mouse Abs (Dako) were primarily used as secondary antibodies. Alternatively, biotinylated anti-sheep/goat immunoglobulin from donkey (1:100 dilution) and streptavidin-biotinylated horseradish peroxidase complex (1:100) (GE Healthcare Life Sciences, USA) were used for the staining of goat polyclonal antibodies. For immunofluorescence, the following secondary Abs were used: Alexa Fluor 568-conjugated donkey anti-goat Ab (Molecular Probes, Eugene, OR, USA); Alexa Fluor 488-conjugated donkey anti-mouse Ab (Jackson ImmunoResearch Laboratories, West Grove, PA, USA); Alexa Fluor 488-conjugated donkey anti-rabbit Ab (Molecular Probes); Alexa Fluor 488 or 568-conjugated goat anti-mouse or anti-rabbit Abs (Molecular Probes). Nuclei were stained with 6-diamidino-2-phenylindole (DAPI). Fluorescent images were captured and analyzed using an Axioplan 2 microscope (Carl Zeiss, Germany).

Following primary mouse monoclonal antibody (mAb), rabbit polyclonal (pAb), rabbit monoclonal (rAb) and goat polyclonal (gAb) antibodies were used at indicated dilution rates: anti-p-S6K (Thr389) rAb (108D; 1:1,000); anti-p-4EBP1 (Thr37/46) rAb (236B4; 1:100) (both from Cell Signaling Technology, Beverly, MA, USA); anti-LRP2 (megalin) mAb (CD7D5; 1:300; Novus Biologicals, Littleton, CO, USA); anti-cubilin gAb (A-20; 1:100; Santa Cruz Biotechnology, Santa Cruz, CA, USA); anti-Ki67 pAb (1:350; Novus Biologicals); anti-TFE3 pAb (1:500; Atlas Antibodies AB,

Stockholm, Sweden); anti-TFEB pAb (1:100; Bioss USA Antibodies, Woburn, MA, USA); anti- β -catenin pAb (1:1,000; Merck Millipore, Darmstadt, Germany); anti-E-cadherin mAb (1:50; BD Biosciences, San Jose, CA, USA).

Cell counting was performed using at least 4 fields (magnification, x100) of each sample. Ki67 positive-cells were compared between ductal and abnormal ductal cells of *Tsc2*^{+/+} and *Tsc2*^{-/-} teratomas, respectively. For E-cadherin and β -catenin, a cell clearly and continuously stained at the perimeter was defined as positive. Statistical analyses were performed with SAS software version R8.1 (SAS Institute Japan, Ltd., Tokyo, Japan). Values of $p < 0.05$ were considered significant.

Western blotting. For western blotting, teratomas were dissected, snap frozen and stored until use. Protein samples were obtained by lysing teratomas in standard sample buffer (50 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol) for SDS polyacrylamide gel electrophoresis. The following primary antibodies were used at the indicated dilutions: anti-TSC2 pAb (C20; 1:500; Santa Cruz Biotechnology); anti-p-S6K (Thr389) rmAb (108D; 1:1,000; Cell Signaling Technology); anti-S6K pAb (C-18; 1:500; Santa Cruz Biotechnology); anti-p-Akt (Ser473) rmAb (D9E; 1:1,000); anti-Akt pAb (1:1,000); and β -actin pAb (1:1,000) (all from Cell Signaling Technology).

Results

***Tsc2*^{-/-} ESCs forms epithelial tumor-like abnormal ductal structures in teratomas.** In the previous study, we found that *Tsc2*^{+/+}, *Tsc2*^{+/-} or *Tsc2*^{-/-} Eker ESCs could differentiate to all three germ layers and epithelial tumor-like abnormal cells forming ductal structures in *Tsc2*^{-/-} teratomas by renal capsule injection in nude mice (15). To confirm this phenomenon in more mature teratomas, we constructed teratomas by subcutaneous-inject-protocol in NOD/SCID mice. H&E-stained sections revealed that *Tsc2*^{-/-} ESCs had the potential to differentiate into all three germ layers in this system (Fig. 1A).

Tsc2^{-/-} teratomas also contained epithelial tumor-like abnormal cells forming ductal structures (Fig. 1B). They comprised large cells with clear, finely granular and occasionally vacuolated cytoplasm. They ruptured the basement membrane and invaded the surrounding parenchyma. These characteristics are reminiscent of Eker rat renal carcinoma (16).

RCCs of Eker rats originate from renal proximal tubules (17). To explore the cell types of abnormal ductal structures in *Tsc2*^{-/-} teratomas, tissue sections were stained for megalin and cubilin. Expressed in epithelial cells of renal proximal tubules, these receptors mediate the endocytosis of numerous ligands. Although megalin expression is widespread among various epithelial cell types, cubilin expression is restricted to cells in renal proximal tubules, small intestine and yolk sac (18). In tissue sections, non-tumorous renal proximal tubules and Eker rat RCCs stained positive for megalin and cubilin (Fig. 2A). Interestingly, abnormal ductal structures of *Tsc2*^{-/-} teratomas were also positive for megalin and cubilin, suggesting epithelial characteristics (Fig. 2B).

Moreover, tissue sections were stained for Ki67 to check the mitotic activity. Ki67-positive cells were frequently detected in abnormal ductal structures as observed in Eker




Figure 1. Teratomas and abnormal ductal structures derived from rat ESCs. (A) H&E staining of teratomas derived from *Tsc2*^{+/+} (top panels), *Tsc2*^{+/-} (middle panels) and *Tsc2*^{-/-} (lower panels) ESCs. Neural epithelium-, cartilage- and gastrointestinal epithelium-like structures indicated by arrows represent ectodermal, mesodermal and endodermal tissues, respectively. (B) RCC of Eker rat (left panel) and abnormal ductal structures in *Tsc2*^{-/-} teratomas (right panel). Scale bars, 100 μ m.

rat RCCs (Fig. 2C). Compared with ductal parts in *Tsc2*^{+/+} teratomas, abnormal ductal structures in *Tsc2*^{-/-} teratomas showed a significant increase in the number of Ki67-positive cells (Fig. 2D). These observations suggest that cells forming abnormal ductal structures of *Tsc2*^{-/-} teratomas possess an epithelial phenotype. Moreover, these findings suggest that the appearance of abnormal ductal structures reflects abnormal cellular differentiation caused by *Tsc2* deficiency.

Activation of mTORC1 signaling in abnormal ductal structures of *Tsc2*^{-/-} teratomas. The activation state of the mTORC1 pathway in the *Tsc2*^{+/+}, *Tsc2*^{+/-} and *Tsc2*^{-/-} teratomas was investigated by immunostaining of downstream protein targets: phosphorylated S6 (p-S6) and phosphorylated 4EBP1 (p-4EBP1).

Positive staining of p-S6 and p-4EBP1 was broadly observed throughout *Tsc2*^{-/-} teratomas and was considerably more intense in abnormal ductal structures as in Eker rat RCCs (Fig. 2B). These results indicated that mTORC1 signaling is activated in abnormal ducts of *Tsc2*^{-/-} teratomas. The importance of the mTORC1 pathway in the formation of abnormal ductal structures was investigated by treatment with the mTOR inhibitor rapamycin after solid teratomas were

formed. All ESCs maintained differentiation of three germ layers but the development of abnormal ductal structures in *Tsc2*^{-/-} teratomas was suppressed under rapamycin-treated conditions (Fig. 3A).

Upregulation of S6 phosphorylation in *Tsc2*^{-/-} teratomas and its suppression by rapamycin were confirmed by western blot analysis (Fig. 3B). Interestingly, Akt (S473) phosphorylation was suppressed in *Tsc2*^{-/-} teratomas but was significantly reactivated by rapamycin, suggesting that the negative feedback on Akt was conferred by an activated mTORC1 signal. These observations support a relationship between mTORC1 hyperactivation and the development of abnormal ductal structures in *Tsc2*^{-/-} teratomas.

E-cadherin and β -catenin localization are dysregulated in *Tsc2*^{-/-} teratomas. Recent evidence suggests that the localization of E-cadherin is regulated by tuberin via an Akt/mTORC1-dependent signaling pathway and that *Tsc2*^{-/-} epithelial cells display a loss of plasma membrane E-cadherin leading to decreased cell-cell adhesion (19). Regulation of β -catenin by hamartin/tuberin complex was also reported (20,21). Therefore, we examined the expression of both E-cadherin and β -catenin by dual staining with cubilin.




Figure 2. Activation of mTORC1 signaling in abnormal ductal structures of teratomas. Tissue sections of Eker rat RCC (A) and abnormal ductal structures in *Tsc2*^{-/-} teratomas (B) were stained with H&E, anti-p-S6, anti-p-4EBP1, anti-megalin, or anti-cubilin antibodies. (C) RCC of Eker rat and abnormal ductal structures in *Tsc2*^{-/-} teratomas were stained with Ki67. Scale bars, 100 μm. (D) Ki67-positive cells were counted in the ductal parts of *Tsc2*^{+/+} and the abnormal ductal parts of *Tsc2*^{-/-} teratomas. *p<0.05, **p<0.001.

In Eker rat RCCs, both E-cadherin and β-catenin primarily localized to the cytoplasm and were weakly detectable in the plasma membrane (Figs. 4A and 5A). In *Tsc2*^{-/-} teratomas,

abnormal ductal structures exhibited both E-cadherin and β-catenin staining at the plasma membrane, but the intensity was weaker and more heterogeneous compared with




Figure 3. Impact of rapamycin treatment on teratomas. (A) *In vivo* differentiation analysis of teratomas derived from *Tsc2*^{+/+} ESCs without (a) or with (b) rapamycin treatment, from *Tsc2*^{-/-} ESCs without (c) or with (d) rapamycin treatment. Scale bars, 100 μm. (B) Western blot analysis of teratomas for tuberin, Pan-S6, p-S6 (S235/236), Pan-Akt, p-Akt (S473) and β-actin. The two *Tsc2*^{-/-} lanes indicate two independent ESC lines.

that in the normal ducts (Figs. 4B and 5B). Furthermore, rapamycin treatment increased the intensity and homogeneity of plasma membrane E-cadherin and β-catenin in *Tsc2*^{-/-} teratomas, although its effect on β-catenin was relatively weak (Figs. 4B and 5B). Taken together, these observations suggest that the membrane localization of E-cadherin and β-catenin was dysregulated by *Tsc2* deficiency in an mTORC1-dependent manner.

Enhanced nuclear localization of TFE3 in Tsc2-/- teratomas.

A recent study provided evidence regarding the regulation of transcription factor EB (TFEB) and transcription factor E3 (TFE3) by mTORC1 (22,23). Deregulated expression and/or gene rearrangement of these members of the MiTF/TFE transcription factor family have been implicated in the development of RCC (24,25). In fact, TFEB is essential for the expression of genes involved in autophagy and lysosome biogenesis and is negatively regulated by mTORC1 (26) On the other hand, TFE3 was recently identified as a key player in mouse ESCs to maintain their self-renewal state and prevent differentiation (22). Interestingly, the knockdown of *Tsc2* increased the nuclear TFE3 concentration through the mTORC1 pathway. Therefore, the localization of TFEB and




Figure 4. Localization of E-cadherin in Eker rat RCCs and teratomas. (A) Dual staining for E-cadherin (green) and cubilin (red). Nuclei were stained with DAPI (blue). Left panels indicate merged images from three types of staining, and right panels indicate E-cadherin staining alone. (a) RCC. (b) Ductal part of *Tsc2*^{+/+} teratomas. (c) Abnormal ductal structure of *Tsc2*^{-/-} teratomas. (d) Rapamycin-treated *Tsc2*^{-/-} teratomas. Scale bars, 100 μm. (B) Comparison of the number of E-cadherin-positive cells in *Tsc2*^{+/+} teratomas without rapamycin and *Tsc2*^{-/-} teratomas with or without rapamycin. ***p<0.0001.

TFE3 was documented in Eker rat kidneys and teratomas by immunohistochemistry analysis.

In normal kidneys, *Tsc2*^{+/+} and *Tsc2*^{+/+} teratomas, TFE3 was found in the cytoplasm of epithelial cells (Fig. 6). On the other hand, nuclear localization was prominent in Eker rat RCCs, whereas surrounding normal components maintained a cytoplasmic localization (Fig. 6). Likewise, in *Tsc2*^{-/-} teratomas, abnormal ductal structures and the stroma clearly revealed high nuclear accumulation of TFE3. Moreover, rapamycin completely abolished TFE3 accumulation in the




Figure 5. Localization of β -catenin in Eker rat RCCs and teratomas. (A) Dual staining for β -catenin (green) and cubilin (red). Nuclei are stained with DAPI (blue). Left panels indicate merged images from the three types of staining, and right panels indicate β -catenin staining alone. (a) RCC. (b) Ductal part of $Tsc2^{+/+}$ teratomas. (c) Abnormal ductal structure of $Tsc2^{-/-}$ teratomas. (d) Rapamycin-treated $Tsc2^{-/-}$ teratomas. Scale bars, 100 μ m. (B) Comparison of the number of β -catenin-positive cells in $Tsc2^{+/+}$ teratomas without rapamycin and $Tsc2^{-/-}$ teratomas with or without rapamycin. ** $p < 0.001$.

nucleus (Fig. 6). On the other hand, there was no apparent phenotype regarding TFE3 localization (data not shown).

These results suggest that TFE3 activation by mTORC1 signaling is involved in the development of abnormal ductal structures in $Tsc2^{-/-}$ teratomas.

Discussion

Various aspects of tissue specificity and differentiation provide important insights into the mechanisms of tumorigenesis. Differentiation experiments using pluripotent stem cells




Figure 6. Nuclear localizations of TFE3 in Eker rat RCCs and teratomas. Dual staining for TFE3 (green) and cubilin (red). Nuclei are stained with DAPI (blue). Left panels indicate merged images from the three types of staining, and right panels indicate TFE3 staining alone. (A) Normal renal cortex. (B) RCC. (C) Abnormal ductal structure of $Tsc2^{-/-}$ teratomas. (D) Rapamycin-treated $Tsc2^{-/-}$ teratomas. Scale bars, 100 μ m.

(PSCs) constitute valid methods to explore the mechanism of tissue-specific tumorigenesis. In general, tumorigenesis is initiated by the loss of tumor-suppressor gene function according to Knudson's two-hit model (27). Since PSCs experience two hits on the tumor-suppressor gene, they constitute ideal tools to investigate the relationship between differentiation abnormalities and tumor initiation. In humans, it is not easy to establish homozygous mutant PSCs for tumor-suppressor genes. Using reprogramming technology, induced PSCs were generated from a patient heterozygous for BRCA1 mutation (5382insC) (28). These mutant iPSCs exhibited increased protein kinase C θ , but the differentiation capacity was not different between wild-type and mutant iPSCs.

In rodents, several studies have documented homozygous mutant PSCs for tumor-suppressor genes (14,29). For instance, Kielman *et al* established mouse ESCs that were homozygously mutated in the *Apc* gene (14). In these mutant ESCs, β -catenin was upregulated. *Apc*-mutated teratomas revealed severe differentiation defects in neuroectodermal, dorsal mesodermal and endodermal lineages. These data suggest that constitutive activation of the *Apc*/ β -catenin pathway results in differentiation defects in the possibly underlying tumorigenesis in the colon and other self-renewing tissues. Kawamata and Ochiya established *Tp53*-mutated rat iPSCs and rat cell lines of *Tp53* mutant strains (29). Unexpectedly, female, but

not male, homozygous *Tp53* mutant rats exhibited neural tube defects. Concurrently, *Tp53*-null rat ESCs resisted differentiation during the embryoid body (EB) formation assay. Although these studies utilizing PSCs clearly illustrate the importance of tumor-suppressor gene function in differentiation, gross abnormalities observed in embryoid bodies or teratomas failed to capture the detailed tissue specificity.

In the present study, the tissue differentiation of PSCs with a deficiency in tumor-suppressor gene presented a different scenario. In *Tsc2*^{-/-} teratomas, various tissue types were generated such as neuroepithelial tissue, squamous epithelium, mesenchyme (undefined immature connective tissue), smooth muscle, cartilage, bone, adipose, gastrointestinal epithelium and glandular (unspecified) tissues. Therefore, these findings suggest that *Tsc2*^{-/-} ESCs possess the ability to differentiate into all three germ layers. Nevertheless, abnormal ductal structures appeared in these differentiated tissues. In *Tsc2*^{-/-} teratomas, cells of abnormal ducts were positive for cubilin and megalin, suggesting aberrant differentiation of some epithelial components. Although the identity of the cell types remains unknown, some cell type-specific effects of *Tsc2* deficiency may emerge during the differentiation of teratomas. We provide evidence that the enhanced activation of mTORC1 pathways contributes to the development of *Tsc2*^{-/-} abnormal ducts. Since abnormal ducts frequently and reproducibly appeared in *Tsc2*^{-/-} teratomas from different cell lines, the accumulation of specific mutations may not be required for their development. Some epigenetic mechanisms may support differentiation defects associated with *Tsc2* deficiency in teratomas. Such mechanisms may be related to the tissue specificity of tumorigenesis in Eker rats.

E-cadherin plays pivotal roles in epithelial cell behavior, tissue formation and cancer suppression (30). During embryonic development, the expression and function of E-cadherin must be normal for the induction and maintenance of polarized and differentiated epithelia (31). The lethality of E-cadherin knockout mice at an early stage of embryogenesis highlights the significance of E-cadherin in normal development and tissue function (32,33). The epithelial-to-mesenchymal transition and loss of E-cadherin expression are closely related and believed to be involved in tumor initiation as well as metastasis (34). An imbalance in β -catenin signaling often results in disease and deregulated growth related to cancer and metastasis (35). During tumor progression, β -catenin signaling is inappropriately activated by the loss of E-cadherin or mutants in various β -catenin signaling components (21). Eker rat RCC and abnormal ductal structures of *Tsc2*^{-/-} teratomas revealed decreased plasma membrane localization of E-cadherin and β -catenin, which was partially corrected by rapamycin treatment. These data support findings from previous studies, suggesting that *Tsc2* deficiency affects E-cadherin localization through perturbations of transport mechanisms (19). The dysregulation of E-cadherin and β -catenin by mTORC1 hyperactivation may cause a polarity defect during the development of *Tsc2*^{-/-} abnormal ducts and tumorigenesis in Eker rats.

The transcription factor TFE3 has been implicated in renal carcinogenesis (23). However, details of TFE3 function have not been elucidated. In a recent study, TFE3 activation by mTORC1 was determined to be essential for the maintenance of self-renewal state and the capacity to withstand differ-

entiation (22). When mTORC1 is activated, TFE3 moves to the nucleus to promote the transcription of estrogen-related receptor β genes involved in the maintenance of self-renewal and pluripotency (22). In our analysis, TFE3 was detected in the nucleus of both Eker rat RCCs and *Tsc2*^{-/-} teratomas, particularly in abnormal ductal structures.

In conclusion, we established a novel experimental system to analyze the differentiation and cell-type specific defects associated with *Tsc2* deficiency using ESCs derived from Eker rats. Future studies should elucidate how mTORC1 hyperactivation and other mechanisms contribute to the development of abnormal ductal structures in *Tsc2*^{-/-} teratomas. Our system will facilitate the understanding of the pathogenesis caused by *Tsc2* deficiency in Eker rats as well as in human tumor stem cells.

Acknowledgements

The present study was supported in part by the following grants: Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan, the MEXT-supported Program for the Strategic Research Foundation at Private Universities, and Science (Japan), Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science, Japan, and the Grants-in-Aid for Scientific Research from the Ministry of Health, Labour and Welfare (Japan). We wish to acknowledge all members of the Department of Molecular Pathogenesis, Juntendo University Graduate School of Medicine who participated in this study. The authors would like to thank Enago for the English language review. Finally, we would like to thank the Gender Equality Promotion Center, Juntendo University, for their assistance with our experiments.

References

1. Eker E and Mossige J: A Dominant gene for renal adenomas in the rat. *Nature* 189: 858-859, 1961.
2. Kobayashi T, Hirayama Y, Kobayashi E, Kubo Y and Hino O: A germline insertion in the tuberous sclerosis 2 (*Tsc2*) gene gives rise to the Eker rat model of dominantly inherited cancer. *Nat Genet* 9: 70-74, 1995.
3. Yeung RS, Xiao GH, Jin F, Lee WC, Testa JR and Knudson AG: Predisposition to renal carcinoma in the Eker rat is determined by germ-line mutation of the tuberous sclerosis 2 (*TSC2*) gene. *Proc Natl Acad Sci USA* 91: 11413-11416, 1994.
4. Rennebeck G, Kleymenova EV, Anderson R, Yeung RS, Artzt K and Walker CL: Loss of function of the tuberous sclerosis 2 tumor suppressor gene results in embryonic lethality characterized by disrupted neuroepithelial growth and development. *Proc Natl Acad Sci USA* 95: 15629-15634, 1998.
5. Everitt JI, Goldsworthy TL, Wolf DC and Walker CL: Hereditary renal cell carcinoma in the Eker rat: A rodent familial cancer syndrome. *J Urol* 148: 1932-1936, 1992.
6. Hino O, Klein-Szanto AJ, Freed JJ, Testa JR, Brown DQ, Vilensky M, Yeung RS, Hino O, Klein-Szanto AJ, Freed JJ, Testa JR, Brown DQ, Vilensky M, Yeung RS, Tartof KD and Knudson AG: Spontaneous and radiation-induced renal tumors in the Eker rat model of dominantly inherited cancer. *Proc Natl Acad Sci USA* 90: 327-331, 1993.
7. Kandt RS, Haines JL, Smith M, Northrup H, Gardner RJ, Short MP, Dumars K, Kandt RS, Haines JL, Smith M, *et al*: Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease. *Nat Genet* 2: 37-41, 1992.
8. European Chromosome 16 Tuberous Sclerosis Consortium: Identification and characterization of the tuberous sclerosis gene on chromosome 16. *Cell* 75: 1305-1315, 1993.

9. van Slegtenhorst M, de Hoogt R, Hermans C, Nellist M, Janssen B, Verhoef S, Lindhout D, van den Ouweland A, Halley D, Young J, *et al*: Identification of the tuberous sclerosis gene *TSC1* on chromosome 9q34. *Science* 277: 805-808, 1997.
10. Orlova KA and Crino PB: The tuberous sclerosis complex. *Ann NY Acad Sci* 1184: 87-105, 2010.
11. Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, Tempst P and Sabatini DM: mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell* 110: 163-175, 2002.
12. Kenerson HL, Aicher LD, True LD and Yeung RS: Activated mammalian target of rapamycin pathway in the pathogenesis of tuberous sclerosis complex renal tumors. *Cancer Res* 62: 5645-5650, 2002.
13. Sarbassov DD, Ali SM, Kim DH, Guertin DA, Latek RR, Erdjument-Bromage H, Tempst P and Sabatini DM: Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. *Curr Biol* 14: 1296-1302, 2004.
14. Kielman MF, Rindapää M, Gaspar C, van Poppel N, Breukel C, van Leeuwen S, Taketo MM, Roberts S, Smits R and Fodde R: *Apc* modulates embryonic stem-cell differentiation by controlling the dosage of beta-catenin signaling. *Nat Genet* 32: 594-605, 2002.
15. Ito Y, Kawano H, Kanai F, Nakamura E, Tada N, Takai S, Horie S, Arai H, Kobayashi T and Hino O: Establishment of *Tsc2*-deficient rat embryonic stem cells. *Int J Oncol* 46: 1944-1952, 2015.
16. McDorman KS and Wolf DC: Use of the spontaneous *Tsc2* knockout (Eker) rat model of hereditary renal cell carcinoma for the study of renal carcinogens. *Toxicol Pathol* 30: 675-680, 2002.
17. Wolf DC, Whiteley HE and Everitt JI: Preneoplastic and neoplastic lesions of rat hereditary renal cell tumors express markers of proximal and distal nephron. *Vet Pathol* 32: 379-386, 1995.
18. Verroust PJ and Christensen EI: Megalin and cubilin - the story of two multipurpose receptors unfolds. *Nephrol Dial Transplant* 17: 1867-1871, 2002.
19. Barnes EA, Kenerson HL, Jiang X and Yeung RS: Tuberin regulates E-cadherin localization: Implications in epithelial-mesenchymal transition. *Am J Pathol* 177: 1765-1778, 2010.
20. Cadigan KM and Nusse R: Wnt signaling: A common theme in animal development. *Genes Dev* 11: 3286-3305, 1997.
21. Valenta T, Hausmann G and Basler K: The many faces and functions of β -catenin. *EMBO J* 31: 2714-2736, 2012.
22. Betschinger J, Nichols J, Dietmann S, Corrin PD, Paddison PJ and Smith A: Exit from pluripotency is gated by intracellular redistribution of the bHLH transcription factor Tfe3. *Cell* 153: 335-347, 2013.
23. Hong SB, Oh H, Valera VA, Baba M, Schmidt LS and Linehan WM: Inactivation of the *FLCN* tumor suppressor gene induces TFE3 transcriptional activity by increasing its nuclear localization. *PLoS One* 5: e15793, 2010.
24. Huan C, Sashital D, Hailemariam T, Kelly ML and Roman CA: Renal carcinoma-associated transcription factors TFE3 and TFEB are leukemia inhibitory factor-responsive transcription activators of E-cadherin. *Proc Natl Acad Sci USA* 100: 6051-6056, 2003.
25. Davis IJ, Hsi BL, Arroyo JD, Vargas SO, Yeh YA, Motyckova G, Valencia P, Perez-Atayde AR, Argani P, Ladanyi M, *et al*: Cloning of an *Alpha-TFEB* fusion in renal tumors harboring the t(6;11)(p21;q13) chromosome translocation. *Proc Natl Acad Sci USA* 100: 6051-6056, 2003.
26. Martina JA, Chen Y, Gucek M and Puertollano R: MTORC1 functions as a transcriptional regulator of autophagy by preventing nuclear transport of TFEB. *Autophagy* 8: 903-914, 2012.
27. Knudson AG Jr: Mutation and cancer: Statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 68: 820-823, 1971.
28. Soyombo AA, Wu Y, Kolski L, Rios JJ, Rakheja D, Chen A, Kehler J, Hampel H, Coughran A and Ross TS: Analysis of induced pluripotent stem cells from a BRCA1 mutant family. *Stem Cell Rep* 1: 336-349, 2013.
29. Kawamata M and Ochiya T: Two distinct knockout approaches highlight a critical role for p53 in rat development. *Sci Rep* 2: 945, 2012.
30. van Roy F and Berx G: The cell-cell adhesion molecule E-cadherin. *Cell Mol Life Sci* 65: 3756-3788, 2008.
31. Takeichi M: Cadherin cell adhesion receptors as a morphogenetic regulator. *Science* 251: 1451-1455, 1991.
32. Riethmacher D, Brinkmann V and Birchmeier C: A targeted mutation in the mouse E-cadherin gene results in defective preimplantation development. *Proc Natl Acad Sci USA* 92: 855-859, 1995.
33. Larue L, Ohsugi M, Hirchenhain J and Kemler R: E-cadherin null mutant embryos fail to form a trophectoderm epithelium. *Proc Natl Acad Sci USA* 91: 8263-8267, 1994.
34. Tiwari N, Gheldof A, Tatari M and Christofori G: EMT as the ultimate survival mechanism of cancer cells. *Semin Cancer Biol* 22: 194-207, 2012.
35. Roose J and Clevers H: TCF transcription factors: Molecular switches in carcinogenesis. *Biochim Biophys Acta* 1424: M23-M37, 1999.