

MEETING ABSTRACT

Open Access

Truncations in the amino terminus reveal a region key to supporting amphetamine-induced efflux by the human serotonin transporter

Sonja Sucic, Carina Kern, Dragana Vidovic, Subhodeep Sarker, Oliver Kudlacek, Harald H Sitte, Michael Freissmuth*

From 17th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Hungarian Society of Experimental and Clinical Pharmacology (MFT) Innsbruck, Austria. 29-30 September 2011

Background

The serotonin transporter (SERT) terminates neurotransmission via reuptake of serotonin from the synaptic cleft. Upon stimulation with amphetamines, SERT switches into an outward transport mode to rapidly release serotonin. We have previously shown that truncation of the first 64 residues of SERT amino terminus leads to loss of amphetamine-induced efflux [1]. This was comparable to the effects of a single point mutation of a juxtamembrane threonine residue at position 81 [1].

Methods

Truncation mutants of SERT amino terminus were generated by removing 22 (Δ 22-SERT), 32 (Δ 32-SERT) or 42 (Δ 42-SERT) amino terminal residues. In addition, alanine scanning mutagenesis was performed along a segment of amino acid residues 32–42. All mutants were pharmacologically characterised in uptake, binding and efflux studies.

Results

Cellular localisation of the mutants examined by confocal microscopy, revealed no differences compared to the wild-type SERT. Functional analysis showed only modest changes in their substrate uptake properties (no significant changes in the K_m values and a moderate decrease in the V_{max} value of Δ 42-SERT). Similarly, there were no marked alterations in the K_D and B_{max} values of imipramine or in the K_i values of *p*-chloroamphetamine and ibogaine, determined in radiolabelled imipramine

binding assays. However, while amphetamine-induced efflux was unimpaired for Δ 22-SERT and to a slight extent decreased for Δ 32-SERT, it was completely abolished for Δ 42-SERT.

Conclusions

Our results shed new light on the functional role of the amino terminus and point to the segment encompassing residues 32–42 as a region of key importance to supporting amphetamine-induced efflux by SERT.

Acknowledgements

This work was supported by SFB35.

Published: 5 September 2011

Reference

1. Sucic S, Dallinger S, Zdrzil B, Weissensteiner R, Jørgensen TN, Holy M, Kudlacek O, Seidel S, Cha JH, Gether U, Newman AH, Ecker GF, Freissmuth M, Sitte HH: **The N terminus of monoamine transporters is a lever required for the action of amphetamines.** *J Biol Chem* 2010, **285**:10924-10938.

doi:10.1186/1471-2210-11-S2-A20

Cite this article as: Sucic et al.: Truncations in the amino terminus reveal a region key to supporting amphetamine-induced efflux by the human serotonin transporter. *BMC Pharmacology* 2011 **11**(Suppl 2):A20.

* Correspondence: michael.freissmuth@meduniwien.ac.at
Institute of Pharmacology, Center of Physiology and Pharmacology, Medical University of Vienna, 1090 Vienna, Austria