

CORRECTION

Open Access



Correction to: The development and validation of a “5A” severity scale for predicting in-hospital mortality after accidental hypothermia from J-point registry data

Yohei Okada^{1,2*}, Tasuku Matsuyama³, Sachiko Morita⁴, Naoki Ehara⁵, Nobuhiro Miyamae⁶, Takaaki Jo⁷, Yasuyuki Sumida⁸, Nobunaga Okada^{3,9}, Makoto Watanabe³, Masahiro Nozawa¹⁰, Ayumu Tsuruoka¹¹, Yoshihiro Fujimoto¹², Yoshiki Okumura¹³, Tetsuhisa Kitamura¹⁴, Shungo Yamamoto¹⁵, Ryoji Iiduka² and Kaoru Koike¹

Correction to: J Intensive Care
<https://doi.org/10.1186/s40560-019-0384-2>

In the original publication of this article [1], the figure legends of Fig. 2 and Fig. 3 are wrong and should be changed as below:

5A scoring model			Severity	Sum Risk score	Predicted mortality*
Covariates	Score				
A ge	60-69	1	Low risk	0-3	7.1 % [5.8-8.4]
	70-79	2	Mild	4	20.5 % [18.7-22.3]
	≥ 80	3	Moderate	5	35.4 % [33.3-37.6]
ADL	Disturbance	1	Severe	6-9	67.4 % [65.1-69.6]
	A rrest		*Predicted mortality: Mean [95%CI] in the validation cohort		
	Arrest or SBP ≤ 60 mmHg	2			
A cidemia	pH: 7.2-7.35	1			
	<7.2	2			
A lbumin					
	≤ 3 (mg/dl)	1			
Sum score		/9			

Fig. 2 5A scoring model

* Correspondence: yokada-kyf@umin.ac.jp

¹Department of Primary Care and Emergency Medicine, Graduate School of Medicine, Kyoto University, 606-8501, Yoshidakonoe-cho, Sakyo, Kyoto, Japan

²Department of Emergency and Critical Care Medicine, Japanese Red Cross Society, Kyoto Daini Hospital, Kyoto, Japan

Full list of author information is available at the end of the article






Fig. 3 Predicted and observed mortality based on the 5A scoring system

Author details

¹Department of Primary Care and Emergency Medicine, Graduate School of Medicine, Kyoto University, 606-8501, Yoshidakonoe-cho, Sakyo, Kyoto, Japan. ²Department of Emergency and Critical Care Medicine, Japanese Red Cross Society, Kyoto Daini Hospital, Kyoto, Japan. ³Department of Emergency Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan. ⁴Senri Critical Care Medical Center, Saiseikai Senri Hospital, Suita, Japan. ⁵Department of Emergency, Japanese Red Cross Society, Kyoto Daiichi Red Cross Hospital, Kyoto, Japan. ⁶Department of Emergency Medicine, Rakuwa-kai Otowa Hospital, Kyoto, Japan. ⁷Department of Emergency Medicine, Uji-Tokushukai Medical Center, Uji, Japan. ⁸Department of Emergency Medicine, North Medical Center, Kyoto Prefectural University of Medicine, Kyoto, Japan. ⁹Department of Emergency and Critical Care Medicine, National Hospital Organization, Kyoto Medical Center, Kyoto, Japan. ¹⁰Department of Emergency and Critical Care Medicine, Saiseikai Shiga Hospital, Ritto, Japan. ¹¹Department of Emergency and Critical Care Medicine, Kyoto Min-Iren Chuo Hospital, Kyoto, Japan. ¹²Department of Emergency Medicine, Yodogawa Christian Hospital, Osaka, Japan. ¹³Department of Emergency Medicine, Fukuchiyama City Hospital, Fukuchiyama, Japan. ¹⁴Division of Environmental Medicine and Population Sciences, Department of Social and Environmental Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan. ¹⁵Department of Healthcare Epidemiology, School of Public Health in the Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Published online: 11 June 2019

Reference

- Okada Y, et al. The development and validation of a "5A" severity scale for predicting in-hospital mortality after accidental hypothermia from J-point registry data. *J Intensive Care*. 2019;7:27 <https://doi.org/10.1186/s40560-019-0384-2>.