

[ PICTURES IN CLINICAL MEDICINE ]


## Focal Hepatic Steatosis Caused by Metastatic Malignant Insulinoma

Naokazu Muramae, Takashi Kobayashi, Kazunori Otsui and Kazuhiko Sakaguchi


**Key words:** focal hepatic steatosis, malignant insulinoma, chemical shift image, ASVS

(Intern Med 60: 653-654, 2021)


(DOI: 10.2169/internalmedicine.5780-20)




Picture 1.



Picture 2.



Picture 3.



Picture 4.

A 77-year-old woman with a history of distal pancreatectomy for insulinoma 3 years previously presented with recurrent symptoms of hypoglycemia. The results of 72-hr fasting test were marginal. Abdominal echography showed multiple spherical-shaped, high-echoic lesions in the liver

without any findings in the pancreas (Picture 1). An enhanced CT scan revealed non-enhanced multiple heterogeneous low density areas in the liver (Picture 2). T1 weighted MRI showed multiple heterogeneous high intensity lesions

Division of General Internal Medicine, Department of Internal Medicine, Kobe University School of Medicine, Japan

Received: July 4, 2020; Accepted: July 29, 2020; Advance Publication by J-STAGE: September 19, 2020

Correspondence to Dr. Naokazu Muramae, nmuramae@med.kobe-u.ac.jp

(Picture 3). A chemical shift image corresponding to Picture 3 indicated focal hepatic steatosis (Picture 4). Although the target liver biopsy showed large fatty droplets to be deposited in hepatocytes, no tumor cells were found in the specimen. A significant increase in the serum immunoreactive insulin (IRI) level on the celiac and proper hepatic arterial stimulation in arterial stimulation venous sampling (ASVS) were observed, and therefore a diagnosis of liver metastasis of malignant insulinoma was made.

Though typical insulinoma occurs as a hypervascular tumor, focal fatty liver around the small metastatic foci occasionally demonstrate characteristic image findings (1, 2).

**The authors state that they have no Conflict of Interest (COI).**

## References

1. Sohn J, Siegelman E, Osiason A. Unusual patterns of hepatic steatosis caused by the local effect of insulin revealed on chemical shift MR imaging. *AJR Am J Roentgenol* **176**: 471-474, 2001.
2. Fregeville A, Couvelard A, Paradis V, Vilgrain V, Warshauer DM. Metastatic insulinoma and glucagonoma from the pancreas responsible for specific peritumoral patterns of hepatic steatosis secondary to local effects of insulin and glucagon on hepatocytes. *Gastroenterology* **129**: 1150, 1365, 2005.

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

---

© 2021 The Japanese Society of Internal Medicine  
*Intern Med* 60: 653-654, 2021