

## Research Article

# The Acidic Fraction of *Isatidis Radix* Regulates Inflammatory Response in LPS-Stimulated RAW264.7 Macrophages through MAPKs and NF- $\kappa$ B Pathway

Zhenyu Fan <sup>1</sup>, Liangliang Cai <sup>1</sup>, Yage Wang <sup>2</sup>, Qiuyan Zhu <sup>1</sup>, Shengnan Wang <sup>1</sup>, and Bohua Chen <sup>1</sup>

<sup>1</sup>Department of Pharmacy, The Affiliated Hospital of Nantong University, Nantong 226000, China

<sup>2</sup>Department of Microsurgery, The Orthopaedic Hospital of Henan Province, Luoyang 471000, China

Correspondence should be addressed to Bohua Chen; [bhchen1963@163.com](mailto:bhchen1963@163.com)

Received 29 September 2020; Revised 17 February 2021; Accepted 25 February 2021; Published 10 March 2021

Academic Editor: Maria Ferraro

Copyright © 2021 Zhenyu Fan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Isatidis Radix*, the dried root of *Isatidis indigotica* Fort, is a traditional heat-clearing and detoxicating herb, which has the antiviral and anti-inflammatory activity and immune regulation. It has been widely used to treat cold, fever, sore throat, mumps, and tonsillitis in clinics. A previous study demonstrated that the acidic fraction of *Isatidis Radix* (RIAF) had strong anti-inflammatory activity, but the mechanism of action was not well elucidated. Lipopolysaccharide- (LPS-) induced RAW264.7 cells were employed to observe the anti-inflammatory activity of RIAF. The level of interleukin-1 $\beta$  (IL-1 $\beta$ ), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), nitric oxide (NO), prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), and interleukin-6 (IL-6) was determined by enzyme-linked immunosorbent assay kits. Western blot was performed to quantify the expression of extracellular signal-regulated kinase (ERK) 1/2, c-jun NH2-terminal kinase (JNK), p38, inducible NO synthetase (iNOS), cyclooxygenase (COX)-2, and nuclear factor- $\kappa$ B (NF- $\kappa$ B). Immunofluorescence assay and electrophoretic mobility shift assay (EMSA) were used to quantify the translocation and the binding-DNA activity of NF- $\kappa$ B. RIAF could inhibit the secretion of inflammatory cytokines (PGE<sub>2</sub>, IL-6, IL-1 $\beta$ , and NO, other than TNF- $\alpha$ ) in a dose-dependent manner. Further investigation showed that the expression of iNOS and COX-2 induced by LPS were down-regulated by treatment with RIAF. Meanwhile, data from the signal pathway exhibited that RIAF significantly suppressed the phosphorylation of ERK1/2, JNK, and p38 and reduced the translocation of NF- $\kappa$ B from the cytoplasm to nucleus, as well as the binding-DNA activity. The anti-inflammatory mechanism of action of RIAF was to reduce inflammation-associated gene expression (iNOS, COX-2, IL-1 $\beta$ , IL-6) by regulating the phosphorylation of the mitogen-activated protein kinases (MAPK) pathway and intervening the activation of the NF- $\kappa$ B pathway, which partly illustrated the basis of treatment of *Isatidis Radix* on cold, fever, sore throat, mumps, and tonsillitis in clinics.

## 1. Introduction

*Isatidis Radix*, the root of *Isatis indigotica* Fort, has been used to treat colds, fever, sore throat, mumps, and tonsillitis in clinics for thousands of years in China [1, 2]. Inflammatory response is a common process of the diseases associated with bacterial or viral infections on the upper respiratory tract [3–5]. *Isatidis Radix* was also reported to have the anti-inflammatory [1–3], antiviral [6], antibacterial [7], and immune regulatory effects [8], which were attributed to syringic acid, isoquinoline derivatives, and flavonoids

[8–10]. In a previous study, most research efforts focused on screening the active components against inflammation, and the acidic fraction of *Isatidis Radix* (RIAF) showed more strong anti-inflammatory activity than the neutral/basic fraction in lipopolysaccharide- (LPS-) induced RAW264.7 cells. However, the molecular mechanism of RIAF was unclear.

When colds, sore throat, and mumps occur, resident macrophages are activated. Some early cytokines such as interleukin-1 $\beta$  (IL-1 $\beta$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) are upregulated by nuclear factor- $\kappa$ B (NF- $\kappa$ B) [11–13] and/

or the mitogen-activated protein kinase (MAPK) signaling pathway [14], then adjacent cells are stimulated to release various chemokines, and neutrophils are recruited to arrange inflammatory response [15]. Meanwhile, cellular damage leads some key enzyme activation (such as cyclooxygenase-2/COX-2 and inducible NO synthetase/iNOS) to produce high levels of prostaglandins (PGE<sub>2</sub>) and nitric oxide (NO) in the airway epithelium [16]. PGE<sub>2</sub> played a key role in the inflammation-related symptoms of common cold such as sore throat, myalgia, fever, nasal congestion, and rhinorrhoea [17, 18]. NO may contribute to inflammation by not only promoting eosinophil infiltration but also increasing microvascular plasma extravasation in the airways [19]. As a classic model of inflammation, the LPS-induced RAW264.7 macrophages cell was used to evaluate the anti-inflammatory activity and to illustrate the molecular mechanism of natural products or mixture. Therefore, it was employed to uncover the molecular mechanism of the acidic fraction of *Isatidis Radix* (RIAF) against inflammatory response.

## 2. Materials and Methods

**2.1. Chemicals and Reagents.** *Isatidis Radix* was purchased from Anhui Province in China and was identified by Professor Qi-Nan Wu, a botanist of Nanjing University of Chinese Medicine (Nanjing, China). A voucher specimen (NJUCM-20171106) was then deposited at a herbarium at the abovementioned location. RAW264.7 cell lines were provided by the Chinese Academy of Medical Sciences (Beijing, China). The enzyme-linked immunosorbent assay (ELISA) kits for TNF- $\alpha$ , IL-1 $\beta$  and interleukin-6 (IL-6) were obtained from eBioscience (Vienna, Austria). Prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) ELISA kits were purchased from Enzo Life Science (Farmingdale, NY, USA). Antibodies against inducible NO synthetase (iNOS), cyclooxygenase-2 (COX-2), phosphorylated or total p65, extracellular signal-regulated kinase (ERK) 1/2-Thr202/Tyr204, c-jun NH2-terminal kinase (JNK)-Thr183/Tyr185, and p38-Thr180/Tyr182 were obtained from Cell Signaling Technology (Beverly, MA, USA). DyLight 488-conjugated secondary antibody, horse radish peroxidase-conjugated secondary antibodies, and primary antibody against  $\beta$ -actin were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Dulbecco's modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were produced by Kaiji Biotechnology (Nanjing, China) and Sijiqing (Hangzhou, China), respectively. LPS (*Escherichia coli* O55:B5), 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazoliumbromide (MTT), 4', 6-diamidino-2-phenylindole (DAPI), and cell-culture-grade DMSO were from Sigma-Aldrich (St. Louis, MO, USA).

**2.2. Preparation of RIAF.** Dried *Isatidis Radix* (4.0 kg) was decocted twice with distilled water (1 : 10 and then 1 : 8, w/v) for 1 h, and the total filtrates were concentrated to approximately 8 L in a vacuum evaporator at 60°C. 95% ethanol (v/v) was added in the concentrated extract up to 60% (v/v) and kept in a low-temperature bath for more than 12 h. After

that, the supernatant was collected by a round of centrifugation at 3000  $\times$  g for 30 min and evaporated under vacuum to approximately 3.5 L. The solution was acidified to pH 5 with HCl and loaded onto a column packed with cation exchange resins (001  $\times$  7, Cangzhou Bon Adsorber Technology Co., Ltd., Cangzhou, China), and the column was washed with water. The eluents were collected (35 L), alkalinized to pH 9 with NaOH, and loaded onto another column packed with anion exchange resins (201  $\times$  7, Cangzhou Bon Adsorber Technology Co., Ltd., Cangzhou, China). After the column was washed with water and 70% ethanol (pH 2), respectively, the ethanol eluents were evaporated under vacuum. It was subjected to D101 macroporous resin and eluted with a gradient ethanol-water (0, 30%, 50%, 95%). The combined ethanol eluents were dried by using a vacuum-drier at 60°C. The yield was 0.64%, and the purity was 89.52% by acid-base titration with phenolphthalein as the indicator, so the fraction was considered as the acidic fraction of *Isatidis Radix* (RIAF). The chemical characterization of RIAF was performed by HPLC according to the previous study [7], and several compounds such as syringic acid, salicylic acid, benzoic acid, caffeic acid, and 2-amino benzoic acid were identified.

**2.3. Cell Culture and Sample Preparation.** RAW264.7 cell lines were cultured in DMEM containing 10% FBS, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin at 37°C with humidified atmosphere containing 5% CO<sub>2</sub>. The stock solution (5 mg/mL) of RIAF was prepared in deionized water.

**2.4. Cell Viability Assay.** RAW264.7 cells ( $1 \times 10^4$  cells/well) were seeded in a 96-well plate overnight and pretreated by RIAF for 24 h at various concentrations (50, 100, 250, and 500  $\mu$ g/mL). After MTT (5 mg/mL), 20  $\mu$ L was added in each well, and cells were incubated for another 4 h. The supernatant was removed, and 150  $\mu$ L DMSO was added in each well. The optical absorbance at 490 nm was measured using a microplate reader (Molecular Devices, Menlo Park, USA).

**2.5. Measurement of Cytokine Levels.** RAW264.7 cells were plated in a 24-well plate at a density of  $5 \times 10^4$  cells/well for PGE<sub>2</sub> and TNF- $\alpha$ , as well as  $2 \times 10^5$  cells/well and  $1 \times 10^5$  cells/well for IL-1 $\beta$  and IL-6. After 24 h, RAW264.7 cells were preincubated with RIAF (125, 250, and 500  $\mu$ g/mL) for 1 h and then stimulated by LPS (1  $\mu$ g/mL) for 24 h. The levels of PGE<sub>2</sub>, TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 in supernatants were quantified by ELISA kits according to the manufacturer's instruction.

**2.6. Measurement of Nitric Oxide (NO).** RAW264.7 cells ( $2 \times 10^5$  cells/well) were seeded in 96-well plates overnight. Subsequently, RAW264.7 cells were treated with RIAF (125, 250, and 500  $\mu$ g/mL) for 24 h at the presence or absence of LPS (1  $\mu$ g/mL). NO level in the supernatant was detected by the Griess reaction [20]. Briefly, 100  $\mu$ L of the Griess reagent (1% sulfanilamide and 0.1% naphthylendiamine in 2.5% phosphoric acid) was mixed with an equal volume of the

supernatant. The optical densities were measured at 540 nm after incubation in dark for 10 min.

**2.7. Western Blot Analysis.** RAW264.7 cells were pretreated with RIAF (125, 250 and 500  $\mu\text{g}/\text{mL}$ ) for 1 h and stimulated with LPS (1  $\mu\text{g}/\text{mL}$ ) for 20 min. Protein samples were prepared through lysing cells in lysis buffer for 10 min on ice. The total level of proteins was determined by bicinchoninic acid (BCA) assay (Sigma Aldrich, St. Louis, MO, USA). After being separated by 10% SDS-PAGE, the proteins of the sample were moved from within gel onto polyvinylidene difluoride (PVDF) membranes (Millipore, Billerica, MA, USA). After being blocked with 5% nonfat dry milk in PBS buffer for 1 h at room temperature, PVDF membranes were washed five times in PBS containing 0.1% Tween 20 (PBST) for 5 min. The blots were incubated with specific primary antibodies in PBST containing 3% bovine serum albumin (BSA) at 4°C overnight and were, in turn, incubated with corresponding horseradish peroxidase-conjugated secondary antibodies for 1 h at room temperature. The bands were exposed to films with enhanced chemiluminescence detection reagents (Pierce, Rockford, IL, USA).

**2.8. Nuclear Factor- $\kappa\text{B}$  (NF- $\kappa\text{B}$ ) Nuclear Translocation Assay.** RAW264.7 cells ( $1 \times 10^4$  cells/well) were cultured in 96-well plates and pretreated with RIAF (125, 250 and 500  $\mu\text{g}/\text{mL}$ ) for 1 h prior to incubation with LPS for 20 min. Cells were then fixed with a 4% paraformaldehyde for 20 min at 4°C, permeabilized with 0.3% Triton X-100 for 20 min, and blocked with 5% BSA for 1 h at room temperature. Subsequently, cells were incubated with primary anti-p-p65 antibody for 2 h at room temperature, followed by DyLight 488-conjugated secondary antibody. After being washed with PBS, cells were incubated in DAPI solution (10  $\mu\text{g}/\text{mL}$ ) for 10 min in the dark. Images of the live cells were visualized by using a confocal laser scanning microscope (BIO-RAD, Hemel Hempstead, UK).

**2.9. Electrophoretic Mobility Shift Assay (EMSA).** RAW264.7 cells were cultured in 100 mm dishes ( $2 \times 10^6$  cells/dish) and grown until confluent. Cells were preincubated with RIAF (125, 250, and 500  $\mu\text{g}/\text{mL}$ ) for 1 h and stimulated with LPS for 30 min. After being washed with PBS, cells were collected, and nuclear and cytosolic extracts were then prepared according to methods described by Wang et al. [21]. EMSA assay was performed as previously described [22]. In brief, oligonucleotide (5'-AGTT-GAGGGGACTTCCAGGC-3', IDTDNA Technologies; Coralville, IA) was synthesized, which was labeled with biotin for the gel retardation assay. Equal amounts of nuclear extracts (10  $\mu\text{g}$ ) were incubated with oligonucleotide for 20 min in 10  $\mu\text{L}$  buffer (0.1 M Tris, pH 7.5, 0.5 M KCl, 10 mM dithiothreitol/DTT). The NF- $\kappa\text{B}$ -DNA mixture was separated by a 5% nondenaturing polyacrylamide gel and then transferred to nylon membranes. The biotinylated DNA was detected with a LightShift chemiluminescent EMSA kit (Thermo scientific, Rockford, IL, USA).

**2.10. Statistical Analysis.** Results were expressed as the mean  $\pm$  SD. Differences were analyzed by one-way analysis of variance (ANOVA) and Student's *t*-test.  $p < 0.05$  was considered as significant.

### 3. Results

**3.1. Effect of RIAF on the Viability of RAW264.7 Cells.** MTT assay showed that RIAF had no cytotoxicity on RAW 264.7 cells at a range of 50–500  $\mu\text{g}/\text{mL}$  (Table 1). Therefore, the test concentration was no more than 500  $\mu\text{g}/\text{mL}$  in following experiments.

**3.2. Anti-Inflammatory Activity of RIAF on RAW264.7 Cells.** After RAW264.7 cells were stimulated by LPS, the levels of cytokines dramatically increased the production of NO, PGE<sub>2</sub>, IL-1 $\beta$ , IL-6, and TNF- $\alpha$  in the supernatants. Most cytokines were significantly reduced by pretreating with RIAF in a dose-dependent manner (as shown in Figure 1), of which the inhibitory rate was 90.04% (Figure 1(a)) and 91.28% (Figure 1(d)) for NO and IL-6 at high concentration. However, it was also noted that RIAF displayed a slight decline in the production of TNF- $\alpha$  (data not showed). The results suggested that RIAF had a good anti-inflammatory activity.

After being treated with RIAF at 125, 250, and 500  $\mu\text{g}/\text{mL}$  for 1 h, RAW264.7 cells were induced by LPS (1  $\mu\text{g}/\text{mL}$ ) for 24 h, and the supernatant was collected for cytokine measurement with an ELISA kit. Data were expressed as means  $\pm$  SD from three independent experiments.  $^{##} p < 0.01$ , compared with control.  $^* p < 0.05$  and  $^{**} p < 0.01$ , compared with the model.

**3.3. RIAF Inhibited LPS-Induced Expression of iNOS and COX-2.** Compared with the normal group, the expression of iNOS and COX-2 were significantly upregulated by LPS (1  $\mu\text{g}/\text{mL}$ ) in RAW264.7 cells. The level of the two proteins was dramatically downregulated in a dose-dependent manner via pretreating with RIAF (Figure 2). COX-2 and iNOS were reported to play a vital role in the formation of NO and PGE<sub>2</sub>, respectively [20–23]. Thus, the results suggested that the activity of RIAF against NO and PGE<sub>2</sub> was involved in iNOS and COX-2.

Cells were treated with RIAF at various concentrations for 1 h and induced by LPS (1  $\mu\text{g}/\text{mL}$ ) for 20 min. Target proteins were separated by western blot, and the level was quantified by Image Tool 3.0. A representative of three independent experiments is shown in Figure 2.  $^{##} p < 0.01$ , compared with control.  $^* p < 0.05$  and  $^{**} p < 0.01$ , compared with the model.

**3.4. Effects of RIAF on LPS-Induced Phosphorylations of MAPKs.** MAPKs are well recognized as an important signal pathway in LPS-induced macrophages activation, which is involved in the expression of specific gene association with inflammatory mediators such as IL-1 $\beta$ , IL-6, and PGE<sub>2</sub> [14]. Meanwhile, Figure 3 also showed, significantly, the strong

phosphorylation of MAPKs (ERK1/2-Thr202/Tyr204, JNK-Thr183/Tyr185, and p38-Thr180/Tyr182) in RAW264.7 cells after exposure to LPS for 20 min compared with the normal group. When RAW264.7 cells were pretreated by RIAF, the level of phosphorylation of the three proteins was obviously reduced in a concentration-dependent manner, and the amount of nonphosphorylated MAPKs was unaffected. The results indicated that RIAF could inhibit the inflammatory response through blocking the phosphorylation of extracellular ERK1/2, JNK, and p38 in LPS-induced RAW264.7 cells.

Cells were incubated with RIAF at various concentrations (125, 250, and 500  $\mu\text{g}/\text{mL}$ ) for 1 h and then stimulated by LPS (1  $\mu\text{g}/\text{mL}$ ) for 20 min. The levels of phosphorylated or total ERK1/2-Thr202/Tyr204, JNK-Thr183/Tyr185, and p38-Thr180/Tyr182 were determined by western blot, which were then quantified by Image Tool 3.0. A representative of three independent experiments is shown in Figure 3. p-ERK1/2, p-JNK, and p-p38: phosphorylation of ERK1/2-Thr202/Tyr204, JNK-Thr183/Tyr185, and p38-Thr180/Tyr182; T-ERK1/2, T-JNK, and T-p38: total ERK1/2, JNK, and p38.  $##p < 0.01$ , compared with control.  $*p < 0.05$  and  $**p < 0.01$ , compared with the model.

**3.5. Effect of RIAF on Phosphorylation of NF- $\kappa$ B in LPS-Induced RAW264.7 Cells.** As another key regulator involving inflammatory response in macrophage cells [24], NF- $\kappa$ B was also investigated to further reveal the anti-inflammatory mechanisms of RIAF. After RAW264.7 was stimulated by LPS, the phosphorylation of the NF- $\kappa$ B p65 subunit was evidently increased. RIAF significantly inhibited the phosphorylation of p65-ser-536 in a dose-dependent manner (see Figure 4). This was why RIAF could reduce the production of IL-1 $\beta$  and IL-6 and the expression of iNOS and COX-2 in LPS-induced RAW264.7 cells.

Cells were incubated with RIAF at various concentrations (125, 250, and 500  $\mu\text{g}/\text{mL}$ ) for 1 h and then stimulated by LPS (1  $\mu\text{g}/\text{mL}$ ) for 20 min. The levels of p-p65, total p65, and  $\beta$ -actin were analyzed by western blot. p-p65: phosphorylation of p65; T-p65: total p65. The content of proteins was quantified as gray values by Image Tool 3.0. A representative of three independent experiments is shown in Figure 4.  $##p < 0.01$ , compared with control.  $*p < 0.05$  and  $**p < 0.01$ , compared with the model.

**3.6. Effect of RIAF on Translocation of NF- $\kappa$ B in LPS-Induced RAW264.7 Cells.** NF- $\kappa$ B nuclear translocation has been reported to be required for NF- $\kappa$ B-dependent gene transcription following LPS stimulation. As shown in Figure 5, compared with the control group, p-p65 was clearly translocated into the nucleus from the cytoplasm of RAW264.7 cells with LPS. However, the level of the phosphorylated p65 in the nucleus was reduced by RIAF after pretreatment for 1 h followed by stimulation with LPS for 20 min, which indicated that RIAF could inhibit LPS-induced NF- $\kappa$ B nuclear translocation in RAW264.7 cells under the concentration of 500  $\mu\text{g}/\text{mL}$ .

After being pretreated with RIAF for 1 h, cells were stimulated by LPS (1  $\mu\text{g}/\text{mL}$ ) for 20 min. Nuclei were stained by DAPI (blue); p-p65 was labeled by DyLight 488-labeled immunostaining (green).

**3.7. Effect of RIAF on LPS-Stimulated NF- $\kappa$ B-DNA Binding Activity.** The NF- $\kappa$ B pathway plays a crucial role in mediating the transcription of a number of genes involved in inflammation on LPS-induced macrophage activation. NF- $\kappa$ B-DNA binding is a directly initial trigger [25]. Figure 6 shows that LPS (1  $\mu\text{g}/\text{mL}$ , 30 min) treatment caused an evident increase in the DNA binding activity of NF- $\kappa$ B, while NF- $\kappa$ B DNA binding complexes were concentration-dependently reduced in RAW264.7 cells pretreated with RIAF.

After being pretreated with RIAF at various concentrations, RAW264.7 cells were exposed to LPS (1  $\mu\text{g}/\text{mL}$ ) for another 20 min. Nuclear extracts were prepared through a Pierce nuclear extraction kit and separated by a 5% polyacrylamide gel with a biotin-labeled oligonucleotide containing the NF- $\kappa$ B consensus sequence. Binding competition assays were run with a 100-fold amount of unlabeled NF- $\kappa$ B oligonucleotide as the competitor (cold NF- $\kappa$ B). Figure 6 is a representative of three independent experiments.

## 4. Discussion

At present, nonsteroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen and celecoxib are used to alleviate common symptoms associated with airway infection or cold (fever, sore throat, and mumps) [18]. Nevertheless, more and more debates have been increasingly reported over the appropriate use of those medications with the serious side effects of NSAIDs [26–28]. Many exterior-releasing and heat-clearing herbs were traditionally used to treat acute respiratory tract infection or complication in practice and showed a good anti-inflammatory activity. For example, Isatidis Radix, a heat-clearing and detoxicating herb, was reported to treat cold, fever, sore throat, and so on and had no obvious side effects [1–3], of which some compounds (epigoitrin, clemastanin B, indigoticoside A, syringic acid, caffeic acid, and salicylic acid) were also reported to inhibit the production of NO and PGE<sub>2</sub>. Most of the compounds had weak acid property due to the phenolic hydroxyl group [2]. It was consistent with the result that RIAF could inhibit inflammatory response.

Inflammation is a defensive response to foreign harmful stimuli, which is initiated by pathogen-associated molecular patterns (PAMPs) of microbial pathogens (such as LPS) in the process of cold, fever, sore throat, and mumps [29] and led to overproduction of cytokines such as NO, PGE<sub>2</sub>, TNF- $\alpha$ , IL-1 $\beta$ , and IL-6. MAPKs and NF- $\kappa$ B were well known as two key signaling pathways involved in the abovementioned process [30, 31]. Among the cytokines, PGE<sub>2</sub> was involved in sore throat via stimulating sensory nerves of pharynx and nasopharynx and increasing capillary permeability [15]. TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and PGE<sub>2</sub> were well regarded as endogenous pyrogens [17, 32]. *In vitro* assay confirmed that RIAF could significantly inhibit the abovementioned

TABLE 1: Effect of RIAF on the viability of RAW264.7 cells.

Concentration ( $\mu\text{g/mL}$ )	Cell viability (%)
—	$100 \pm 0.00$
500	$103.97 \pm 3.71$
250	$110.18 \pm 5.20$
100	$113.59 \pm 3.14$
50	$112.22 \pm 1.08$




FIGURE 1: Effects of RIAF on inflammatory cytokines in LPS-induced RAW264.7 cells.




FIGURE 2: Effects of RIAF on the expression of COX-2 and iNOS in LPS-induced RAW264.7 cells.




FIGURE 3: Effect of RIAF on MAPKs in LPS-induced RAW264.7 cells.




FIGURE 4: Effect of RIAF on phosphorylation of NF-κB in LPS-induced RAW264.7 cells.

cytokines except for TNF- $\alpha$ , which implied that the therapeutic effect of RIAF was contributed to the anti-inflammatory activity.

iNOS and COX-2 were two rate-limiting enzymes in the NO or PGE<sub>2</sub> synthesis pathway, and the expression was regulated by the MAPKs and NF- $\kappa$ B pathway in LPS-induced inflammatory responses [23, 33, 34]. Western blot showed that RIAF could downregulate the expression of the two proteins, which indicated that the mechanism of RIAF against NO and PGE<sub>2</sub> could be partially attributed to reducing the level of the two inducible enzymes on attack by foreign stimulants.

After the macrophage was stimulated by LPS, LPS could induce the phosphorylation of MAPKs or NF- $\kappa$ B through toll-like receptor 4 and initiate the expression of cytokines (such as IL-1, IL-6, PGE<sub>2</sub>, and TNF) [35–37]. Among MAPKs, ERK1/2, JNK, and p38 had a central role in the regulation of cytokines [38]. p50-p65 dimers, an NF- $\kappa$ B subgroup, played an important role in regulating cytokine production by residues the phosphorylation of the Ser<sup>536</sup> residue on p65 and degradation of inhibitor kappa B ( $\kappa$ B) [39]. The results revealed that RIAF could significantly suppress the phosphorylation of the MAPK and NF- $\kappa$ B pathway. Nuclear translocation assay and EMSA further




FIGURE 5: Effect of RIAF on the nuclear translocation of p-p65 protein.

confirmed that RIAF could decrease the translocation of NF- $\kappa$ B from the cytoplasm to nucleus and block binding of p65 to target genes. Therefore, it was reasonable that RIAF could inhibit inflammatory cytokines in LPS-induced RAW264.6 cells through regulating the phosphorylation of the MAPKs pathway and intervening the activation of the NF- $\kappa$ B pathway. Meanwhile, a previous study [3] also showed that methanolic extracts of *Isatidis Radix* could significantly inhibit the release of inflammatory mediators (NO and PGE<sub>2</sub>) and proinflammatory cytokines (IL-6 and TNF- $\alpha$ ) in LPS-stimulated RAW264.7 macrophages through inhibition of NF- $\kappa$ B signaling. These results indicated that RIAF was the functional fraction to regulate the anti-inflammatory effect of *Isatidis Radix*. Yet, it should also note that RIAF was not able to inhibit TNF- $\alpha$ .

The reason for the selective inhibition of RIAF on NO, PGE<sub>2</sub>, IL-1 $\beta$ , and IL-6, but not TNF- $\alpha$ , was still unclear. Traditional Chinese medicine (TCM) is characterized by

multiple components, weak effects, and synergistic effects. The overall effect of *Isatidis Radix* on the inflammatory response was not the result of the action of a single or several compounds, but the interaction of multiple compounds. Compared with *Isatidis Radix*, RIAF had some differences in the ratio and kind of the active compounds, which may influence the effect of RIAF on TNF- $\alpha$ . Indeed, further investigation is required to confirm this assumption.

In conclusion, our observations suggested that RIAF displayed a strong anti-inflammatory activity by suppressing MAPKs and NF- $\kappa$ B activation to reduce inflammation-associated gene expression (iNOS, COX-2, IL-1 $\beta$ , and IL-6) (Figure 7). This implied that RIAF, at least in part, was the basis of the therapeutic efficacy of *Isatidis Radix* in cold, fever, sore throat, mumps, and tonsillitis. However, it is necessary to further investigate the activity and mechanism of RIAF *in vivo*. Meanwhile, compounds of RIAF against inflammatory responses should be identified in the future.

LPS (1 μg/ml)	Cold	-	-	+	+	+	+
RIAF (μg/ml)	NF-κB	-	-	-	125	250	500




FIGURE 6: Effect of RIAF on LPS-stimulated NF-κB-DNA complex formation in RAW264.7 cells.




FIGURE 7: Putative RIAF inhibition pathway in macrophage-mediated inflammatory signaling.



## Data Availability

The underlying data supporting the results in the manuscript are included within the article.

## Conflicts of Interest

All the authors declare no conflicts of interest.

## Authors' Contributions

Bohua Chen conceived the study. Yage Wang prepared the acidic fraction of *Isatidis Radix*. Zhenyu Fan, Qiuyan Zhu, and Shengnan Wang carried out cell culture, measurement, and translocation or electrophoretic motility shift assay. Zhenyu Fan and Liangliang Cai interpreted the results and wrote the manuscript.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (Grant No. 81900528). The authors are grateful to Chengwei Liu, Xiang Li, and Weihua Wang for excellent technical assistance throughout the project.

## References

- [1] Y. Ma, N. Li, C. Liu et al., "Comparative studies on anti-inflammatory and analgesic activities of different fractions from *Isatidis Radix*," *Hinese Traditional and Herbal Drugs*, vol. 45, no. 17, pp. 2517–2521, 2014.
- [2] P. Xiao, H. Huang, J. Chen, and X. Li, "In vitro antioxidant and anti-inflammatory activities of *Radix isatidis* extract and bioaccessibility of six bioactive compounds after simulated gastro-intestinal digestion," *Journal of Ethnopharmacology*, vol. 157, pp. 55–61, 2014.
- [3] E. K. Shin, D. H. Kim, H. Lim, H.-K. Shin, and J.-K. Kim, "The anti-inflammatory effects of a methanolic extract from *Radix isatidis* in murine macrophages and mice," *Inflammation*, vol. 33, no. 2, pp. 110–118, 2010.
- [4] C. Michel, S. Salvatico, H. Belkhelda, L. Haddioui, and C. Roques, "Activity of drill lozenges on the main microorganisms responsible for upper respiratory tract infections," *European Annals of Otorhinolaryngology, Head and Neck Diseases*, vol. 130, no. 4, pp. 189–193, 2013.
- [5] D. Sidell and N. L. Shapiro, "Acute tonsillitis," *Infectious Disorders-Drug Targets*, vol. 12, no. 4, pp. 271–276, 2012.
- [6] Z. W. Chen, L. W. Wu, S. T. Liu, C. P. Cai, P. F. Rao, and L. J. Ke, "Mechanism study of anti-influenza effects of *Radix isatidis* water extract by red blood cells capillary electrophoresis," *Zhongguo Zhong Yao Za Zhi*, vol. 31, no. 20, pp. 1715–1719, 2006.
- [7] W. Kong, Y. Zhao, L. Shan, X. Xiao, and W. Guo, "Thermochemical studies on the quantity–antibacterial effect relationship of four organic acids from *Radix isatidis* on *Escherichia coli* growth," *Biological & Pharmaceutical Bulletin*, vol. 31, no. 7, pp. 1301–1305, 2008.
- [8] Y.-L. Zhao, J.-B. Wang, L.-M. Shan, C. Jin, L. Ma, and X.-H. Xiao, "Effect of *Radix isatidis* polysaccharides on immunological function and expression of immune related cytokines in mice," *Chinese Journal of Integrative Medicine*, vol. 14, no. 3, pp. 207–211, 2008.
- [9] L.-W. He, H.-Q. Liu, Y.-Q. Chen, J.-Y. Yang, T.-L. Wang, and W. Li, "Total synthesis and anti-viral activities of an extract of *Radix isatidis*," *Molecules*, vol. 19, no. 12, pp. 20906–20912, 2014.
- [10] Y. Liu, J. Fang, T. Lei, W. Wang, and A. Lin, "Anti-endotoxic effects of syringic acid of *Radix isatidis*," *Journal of Huazhong University of Science and Technology. Medical sciences=Hua zhong ke ji da xue xue bao. Yi xue Ying De wen ban=Huazhong keji daxue xuebao. Yixue Yingdewen ban*, vol. 23, no. 2, pp. 206–208, 2003.
- [11] S. Akira, "Toll-like receptor signaling," *Journal of Biological Chemistry*, vol. 278, no. 40, pp. 38105–38108, 2003.
- [12] I. Iontcheva, S. Amar, K. H. Zawawi, A. Kantarci, and T. E. Van Dyke, "Role for moesin in lipopolysaccharide-stimulated signal transduction," *Infection and Immunity*, vol. 72, no. 4, pp. 2312–2320, 2004.
- [13] N. R. Aggarwal, L. S. King, and F. R. D'Alessio, "Diverse macrophage populations mediate acute lung inflammation and resolution," *American Journal of Physiology-Lung Cellular and Molecular Physiology*, vol. 306, no. 8, pp. L709–L725, 2014.
- [14] Y. Luo, M. Liu, Y. Dai et al., "Norisoboldine inhibits the production of pro-inflammatory cytokines in lipopolysaccharide-stimulated RAW 264.7 cells by down-regulating the activation of MAPKs but not NF- $\kappa$ B," *Inflammation*, vol. 33, no. 6, pp. 389–397, 2010.
- [15] S. Bathala and R. Eccles, "A review on the mechanism of sore throat in tonsillitis," *The Journal of Laryngology & Otology*, vol. 127, no. 3, pp. 227–232, 2013.
- [16] R. Eccles, "Mechanisms of symptoms of common cold and flu," in *Common Cold*, R. Eccles and O. Weber, Eds., Birkhäuser, Basel, Switzerland, pp. 23–45, 2009.
- [17] E. Pecchi, M. Dallaporta, A. Jean, S. Thirion, and J.-D. Troadec, "Prostaglandins and sickness behavior: old story, new insights," *Physiology & Behavior*, vol. 97, no. 3–4, pp. 279–292, 2009.
- [18] R. Eccles, "Understanding the symptoms of the common cold and influenza," *The Lancet Infectious Diseases*, vol. 5, no. 11, pp. 718–725, 2005.
- [19] B. J. Nevin and K. J. Broadley, "Nitric oxide in respiratory diseases," *Pharmacology & Therapeutics*, vol. 95, no. 3, pp. 259–293, 2002.
- [20] G. Chen, K.-K. Li, C.-H. Fung et al., "Er-Miao-San, a traditional herbal formula containing *Rhizoma Atractylodis* and *Cortex Phellodendri* inhibits inflammatory mediators in LPS-stimulated RAW264.7 macrophages through inhibition of NF- $\kappa$ B pathway and MAPKs activation," *Journal of Ethnopharmacology*, vol. 154, no. 3, pp. 711–718, 2014.
- [21] X. Wang, T. Kikuchi, and Y. Rikihisa, "Proteomic identification of a novel *Anaplasma phagocytophilum* DNA binding protein that regulates a putative transcription factor," *Journal of Bacteriology*, vol. 189, no. 13, pp. 4880–4886, 2007.
- [22] A. Duvoix, S. Delhalle, R. Blasius et al., "Effect of chemopreventive agents on glutathione S-transferase P1-1 gene expression mechanisms via activating protein 1 and nuclear factor kappaB inhibition," *Biochemical Pharmacology*, vol. 68, no. 6, pp. 1101–1111, 2004.
- [23] Z. Zhao, X. Tang, X. Zhao et al., "Tylvalosin exhibits anti-inflammatory property and attenuates acute lung injury in different models possibly through suppression of NF- $\kappa$ B activation," *Biochemical Pharmacology*, vol. 90, no. 1, pp. 73–87, 2014.
- [24] Y.-M. Ma, X.-Z. Zhang, Z.-Z. Su et al., "Insight into the molecular mechanism of a herbal injection by integrating

- network pharmacology and in vitro,” *Journal of Ethnopharmacology*, vol. 173, pp. 91–99, 2015.
- [25] T. Zhang, L. Sun, R. Liu et al., “A novel naturally occurring salicylic acid analogue acts as an anti-inflammatory agent by inhibiting nuclear factor-kappaB activity in RAW264.7 macrophages,” *Molecular Pharmaceutics*, vol. 9, no. 3, pp. 671–677, 2012.
- [26] W. Thurlow, “Acetaminophen, aspirin, and renal failure,” *The New England Journal of Medicine*, vol. 346, no. 20, pp. 1588–1589, 2002.
- [27] S. Bancos, M. P. Bernard, D. J. Topham, and R. P. Phipps, “Ibuprofen and other widely used non-steroidal anti-inflammatory drugs inhibit antibody production in human cells,” *Cellular Immunology*, vol. 258, no. 1, pp. 18–28, 2009.
- [28] X. Gao, M. Guo, L. Peng et al., “UPLC Q-TOF/MS-based metabolic profiling of urine reveals the novel antipyretic mechanisms of qingkailing injection in a rat model of yeast-induced pyrexia,” *Evidence Based Complementary and Alternative Medicine*, vol. 2013, Article ID 864747, 8 pages, 2013.
- [29] C. C. Corriveau and R. L. Danner, “Endotoxin as a therapeutic target in septic shock,” *Infectious Agents Disease*, vol. 2, no. 1, pp. 35–43, 1993.
- [30] “The coming era of human phenotyping,” *Nature Biotechnology*, vol. 33, no. 6, p. 567, 2015.
- [31] J. Choi, S.-H. Kim, and S. Kim, “Suppressive effects of PG201, an antiarthritic botanical formulation, on lipopolysaccharide-induced inflammatory mediators in RAW264.7 cells,” *Experimental Biology and Medicine*, vol. 237, no. 5, pp. 499–508, 2012.
- [32] L. M. Harden, I. du Plessis, S. Poole, and H. P. Laburn, “Interleukin (IL)-6 and IL-1 $\beta$  act synergistically within the brain to induce sickness behavior and fever in rats,” *Brain, Behavior, and Immunity*, vol. 22, no. 6, pp. 838–849, 2008.
- [33] R. J. Choi, E. M. Shin, H. A. Jung, J. S. Choi, and Y. S. Kim, “Inhibitory effects of kaurenoic acid from *Aralia continentalis* on LPS-induced inflammatory response in RAW264.7 macrophages,” *Phytomedicine*, vol. 18, no. 8-9, pp. 677–682, 2011.
- [34] M. K. O’Banion, “Prostaglandin E2 synthases in neurologic homeostasis and disease,” *Prostaglandins & Other Lipid Mediators*, vol. 91, no. 3-4, pp. 113–117, 2010.
- [35] M. Fujihara, M. Muroi, K.-I. Tanamoto, T. Suzuki, H. Azuma, and H. Ikeda, “Molecular mechanisms of macrophage activation and deactivation by lipopolysaccharide: roles of the receptor complex,” *Pharmacology & Therapeutics*, vol. 100, no. 2, pp. 171–194, 2003.
- [36] T. Hanada and A. Yoshimura, “Regulation of cytokine signaling and inflammation,” *Cytokine and Growth Factor Reviews*, vol. 13, no. 4-5, pp. 413–421, 2002.
- [37] H. W. Jung, R. Mahesh, J. H. Park, Y. C. Boo, K. M. Park, and Y.-K. Park, “Bisabolangelone isolated from *Ostericum koreanum* inhibits the production of inflammatory mediators by down-regulation of NF- $\kappa$ B and ERK MAP kinase activity in LPS-stimulated RAW264.7 cells,” *International Immunopharmacology*, vol. 10, no. 2, pp. 155–162, 2010.
- [38] S.-H. Yang, A. D. Sharrocks, and A. J. Whitmarsh, “MAP kinase signalling cascades and transcriptional regulation,” *Gene*, vol. 513, no. 1, pp. 1–13, 2013.
- [39] P. Li, J. Chen, J. Wang et al., “Systems pharmacology strategies for drug discovery and combination with applications to cardiovascular diseases,” *Journal of Ethnopharmacology*, vol. 151, no. 1, pp. 93–107, 2014.