

CLINICAL CONCEPTS

Emergency Medical Services

Rationale and development of a prehospital goal-directed bundle of care to prevent rearrest after return of spontaneous circulation

David G. Dillon MD, PhD¹ | Juan Carlos C. Montoy MD, PhD² | Nichole Bosson MD, MPH^{3,4,5,6} | Jake Toy DO, MS^{3,4,5,6} | Senai Kidane MD^{7,8} | Dustin W. Ballard MD, MBE^{1,8,9} | Marianne Gausche-Hill MD^{3,4,5,6} | Joelle Donofrio-Odmann DO¹⁰ | Shira A. Schlesinger MD, MPH^{11,12,13} | Katherine Staats MD^{14,15} | Clayton Kazan MD, MS¹⁶ | Brian Morr BS, MICP¹⁷ | Kristin Thompson RN¹³ | Kevin Mackey MD^{8,17} | John Brown MD, MPA¹⁸ | James J. Menegazzi PhD¹⁹ | the California Resuscitation Outcomes Consortium

¹Department of Emergency Medicine, University of California, Davis, California, USA

²Department of Emergency Medicine, University of California, San Francisco, California, USA

³Los Angeles County EMS Agency, Santa Fe Springs, California, USA

⁴Lundquist Institute for Biomedical Innovation, Harbor-UCLA Medical Center, Torrance, California, USA

⁵Department of Emergency Medicine, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

⁶Department of Emergency Medicine, Harbor-UCLA Medical Center, Torrance, California, USA

⁷Contra Costa County Emergency Medical Services, Martinez, California, USA

⁸The Permanente Medical Group, Oakland, California, USA

⁹Division of Research, Kaiser Permanente Northern California, Oakland, California, USA

¹⁰Departments of Emergency Medicine and Pediatrics, University of California, San Diego, California, USA

¹¹Department of Emergency Medicine, David Geffen School of Medicine, Los Angeles, California, USA

¹²Harbor-UCLA Medical Center, Los Angeles, California, USA

¹³Newport Beach Fire Department, Newport Beach, California, USA

¹⁴Imperial County Emergency Medical Services, Imperial, USA

¹⁵Department of Emergency Medicine, University of California, San Diego, California, USA

¹⁶Los Angeles County Fire Department, Los Angeles, California, USA

¹⁷Sacramento City Fire Department, Sacramento, California, USA

¹⁸San Francisco Emergency Medical Services Agency, San Francisco Department of Public Health, San Francisco, California, USA

¹⁹Department of Emergency Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA

Supervising Editor: Juan March, MD

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Author(s). *Journal of the American College of Emergency Physicians Open* published by Wiley Periodicals LLC on behalf of American College of Emergency Physicians.

Correspondence

Juan Carlos C. Montoy, 1001 Potrero Ave, Rm 6A01, San Francisco, CA 94110, USA.
Email: juancarlos.montoy@ucsf.edu

Funding information

National Heart, Lung, and Blood Institute,
Grant/Award Numbers: 1R01HL117979,
K38HL165363

Abstract

In patients with out-of-hospital cardiac arrest (OHCA) who attain return of spontaneous circulation (ROSC), rearrest while in the prehospital setting represents a significant barrier to survival. To date, there are limited data to guide prehospital emergency medical services (EMS) management immediately following successful resuscitation resulting in ROSC and prior to handoff in the emergency department. Post-ROSC care encompasses a multifaceted approach including hemodynamic optimization, airway management, oxygenation, and ventilation. We sought to develop an evidenced-based, goal-directed bundle of care targeting specified vital parameters in the immediate post-ROSC period, with the goal of decreasing the incidence of rearrest and improving survival outcomes. Here, we describe the rationale and development of this goal-directed bundle of care, which will be adopted by several EMS agencies within California. We convened a group of EMS experts, including EMS Medical Directors, quality improvement officers, data managers, educators, EMS clinicians, emergency medicine clinicians, and resuscitation researchers to develop a goal-directed bundle of care to be applied in the field during the period immediately following ROSC. This care bundle includes guidance for prehospital personnel on recognition of impending rearrest, hemodynamic optimization, ventilatory strategies, airway management, and diagnosis of underlying causes prior to the initiation of transport.

KEYWORDS

hemodynamic management, out of hospital cardiac arrest, oxygenation, post resuscitation care, prehospital emergency care, ventilation

1 | INTRODUCTION

Emergency medical services (EMS) management of patients with out-of-hospital cardiac arrest (OHCA) immediately following return of spontaneous circulation (ROSC) is a crucial and under-studied component of early cardiac arrest care. While reported rates of field ROSC are near 40%, survival and neurologically intact survival rates after OHCA remain low at approximately 9% and 7%, respectively.¹ Rearrest in the prehospital setting represents a significant barrier to survival, with up to 40% of patients who achieve ROSC experiencing a rearrest event prior to hospital arrival.²⁻⁶ Prior work has shown that rearrest is independently associated with decreased survival and reduced rates of survival with good neurologic status.³⁻⁶ Thus, optimization of recognition of impending rearrest, hemodynamics, oxygenation, and ventilation in the immediate post-ROSC period is critical to reduce the risk of rearrest and improve chances of survival.^{5,6}

Current EMS protocols for post-ROSC care are heterogeneous, with the most common recommended interventions being to obtain a 12-lead electrocardiogram and administer vasopressors; few protocols offer guidance on the timing of post-ROSC stabilization and transport initiation.⁷ The American Heart Association (AHA) 2020 Post-Resuscitation Care guidelines recommend prompt initiation of hemodynamic and ventilatory support to mitigate the systemic impacts of ischemia-reperfusion injury, as hypotension, hypoxia, and hyperven-

tilation in the post-ROSC phase have all been found to be independently associated with worse outcomes after OHCA.⁸⁻¹³ However, timing of these interventions is not specified and there is no direction on whether these should be performed prior to transport; similarly, clearly defined vital sign targets have not been established.¹⁴ Multiple recent studies attempting to evaluate optimal blood pressure and oxygenation targets have yielded inconsistent results.^{10,13,15-18} Given that most patients who rearrest do so within the first few minutes after ROSC, it is likely that interventions to prevent rearrest would be most effective if implemented by EMS in the immediate post-ROSC period and not delayed until ambulance transport or hospital arrival.⁴

We contend that instituting a goal-directed bundle of care in the immediate post-ROSC period, prior to transport, will decrease the incidence of rearrest and thereby improve patient survival outcomes. The recommendations of this bundle of care are limited to the clinical treatment thresholds and protocols outlined in this manuscript, which focus on the post-arrest period. EMS providers will follow their local EMS agency's protocols for cardiac arrest care, and equipment choices are left to the discretion of provider agencies. The California Resuscitation Outcomes Consortium (CAL-ROC) investigators aim to conduct a multicenter clinical trial in EMS systems across California to evaluate the impact of a prehospital goal-directed care bundle for post-ROSC treatment to be initiated on scene. The objective of this paper is to describe the rationale for, and development of, this goal-directed bundle of care

Clinical Research Perspective

- Among patients with out-of-hospital cardiac arrest who attain return of spontaneous circulation (ROSC), rearrest while in the prehospital setting represents a significant barrier to survival. We hypothesized that targeted interventions in the immediate post-ROSC period should be performed in the field to further stabilize patients and prevent rearrest prior to hospital arrival.
- We convened a diverse group of experts including emergency medical services (EMS) Medical Directors, emergency medicine clinicians, and resuscitation researchers to develop a goal-directed bundle of care to be applied in the field during the period immediately following ROSC, with the goal of preventing rearrest prior to hospital arrival.
- This care bundle, based on national and international guidelines, may be considered for adoption by EMS agencies and includes guidance for prehospital personnel on recognition of impending rearrest, hemodynamic optimization, ventilatory strategies, airway management, and diagnosis of underlying causes prior to the initiation of transport.

and to offer it as a protocol that can be adopted by other EMS agencies seeking to implement the most up-to-date evidence.

2 | RATIONALE

Hypotension, hypoxia, and hyperventilation have been strongly associated with decreased survival and worsened neurologic outcomes in patients with OHCA.³⁻⁶ Rearrest is also well established to result in decreased survival and worsened neurologic outcomes.³⁻⁶ Mechanistically, we believe that the prevention of hypotension, hypoxia, and hyperventilation will result in reduced rates of rearrest, and will in turn result in increased survival and increased survival with good neurologic outcomes. Based on current best practices and recommendations, we developed a post-ROSC goal-directed bundle of care that directly addresses each of these physiologic insults, presented in Figure 1 and supported by the evidence below. The recommendations made by the AHA/American College of Cardiology (ACC), European Resuscitation Council and European Society of Intensive Care Medicine, and the International Liaison Committee on Resuscitation (ILCOR) provide the foundation for the development of this bundle of care and are summarized in Table 1. There are many other important and active areas of ongoing research in cardiac arrest management, including the use of cardiopulmonary resuscitation (CPR) assistance devices, targeted temperature management in the field, and post-arrest management after a transfer of care in the ED, among many other areas of active investigation. This bundle does not directly support or reject any of these

interventions. This bundle applies to adults (≥ 18 year old) with OHCA, as the management of pediatric cardiac arrest strategies, including treatment thresholds, differ for pediatric as opposed to adult patients. The evidence for the bundle components is further described in the following sections.

3 | METHODS

The goal of this endeavor was to develop and present a standardized OHCA bundle of care within the CAL-ROC consortium based on currently published national and international guidelines and expert opinion. We used an iterative method during once-weekly CAL-ROC video conference calls (from December 2022 through February 2023) to develop the Stabilize on Scene (SOS)-bundle. Stakeholders from EMS agencies and experts in the pre-hospital treatment of OHCA self-identified from leadership positions within the CAL-ROC consortium, including EMS Medical Directors, quality improvement officers, data managers, educators, EMS clinicians, emergency medicine clinicians, and resuscitation researchers. The same group of stakeholders and experts were present and involved throughout the process of developing this bundle of care. Initial meetings were open-ended and targeted toward identifying potential components of the SOS-bundle that should be standardized. For each component that was identified, the available evidence was then gathered and discussed in working groups until a consensus was reached for the specific action, target, triggers, and interventions to be included in the bundle. These recommendations were then brought back to the larger group and edited in an iterative process until the bundle as a whole was finalized via unanimous support of all authors (Figure 1). Ultimately, there were four components of the bundle: hemodynamic optimization, ventilation and oxygenation, airway management, and ST-elevation myocardial infarction (STEMI) identification. The study was deemed not human subjects research per each institution's Human Research Protection Program.

3.1 | Circulation: Hemodynamic optimization

Hypotension in the post-ROSC period is common and is associated with poor neurologic outcomes and higher mortality.^{6,8,10,11,19-22} Nearly 40% of patients with ROSC experience rearrest, and post-ROSC hypotension is associated with a higher risk of rearrest and decreased survival.^{2,4} Hypotension may also be harmful in a dose-dependent mechanism, as longer periods of hypotension and lower blood pressures are both associated with higher mortality.⁸

The association between poor perfusion and worse outcomes among patients with ROSC is well established, and blood pressure management is recommended across multiple guidelines.²³⁻²⁶ While most of these data are from observational studies, there is emerging evidence that the avoidance or prompt treatment of post-ROSC hypotension may be beneficial.^{22,27} Interventions to address hypotension may help reduce the risk of rearrest or otherwise improve patient outcomes.

CAL-ROC SOS BUNDLE			
CIRCULATION		AIRWAY	
HEMODYNAMIC OPTIMIZATION		AIRWAY MANAGEMENT	
GOAL = AVOID HYPOTENSION		GOAL = SECURE AIRWAY	
<u>Actions</u>	<u>Trigger</u>	<u>Actions</u>	<u>Trigger</u>
Assess BP q3min	All		If BVM ineffective or after
Administer NS Bolus 500mL	All	Place Advanced Airway*	HD optimized and prior to transport
Prepare push-dose epinephrine	All	Confirm airway with EtCO ₂	All
Administer push-dose epinephrine	SBP<120mmHg	BREATHING	
Repeat NS Bolus 500mL	Persistent SBP<120mmHg	VENTILATION AND OXYGENATION	
Repeat push-dose epinephrine	Persistent SBP<120mmHg	GOAL = AVOID HYPER/HYPOVENTILATION	
STEMI IDENTIFICATION		<u>Actions</u>	
GOAL = IDENTIFY STEMI & DYSRHYTHMIAS		<u>Trigger</u>	
<u>Actions</u>	<u>Trigger</u>	Initiate ventilations at 10 BPM	All
Continuous cardiac rhythm monitoring	Focused assessment q3mins or if clinical deterioration	Continuous EtCO ₂ and SpO ₂ monitoring	Focused assessment q3mins or if clinical deterioration
Obtain and transmit 12-lead ECG	All	Target EtCO ₂ 35-45mmHg	All
		Target SpO ₂ 94-98%	All

FIGURE 1 The California Resuscitation Outcomes Consortium (CAL-ROC) stabilize on scene (SOS) post-return of spontaneous circulation (post-ROSC) goal-directed bundle of care. Summary of the interventions and events or physiologic thresholds to trigger each intervention.

TABLE 1 Society and California Resuscitation Outcomes Consortium (CAL-ROC) guidelines for post-return of spontaneous circulation (post-ROSC) goals of care.

	American College of Cardiology/American Heart Association	European Resuscitation Council and European Society of Intensive Care Medicine	International Liaison Committee on Resuscitation	CAL-ROC SOS-bundle
Airway	Early placement of endotracheal tube	Intubate unless immediate return of cerebral function ^a Supraglottic airway if no experienced intubator	Insert advanced airway	Intubate or supraglottic airway per local protocol
Oxygenation	100% FiO ₂ then titrate to 92%–98% when able	100% FiO ₂ until able to measure; then SpO ₂ 94%–98%	100% oxygen until reliable measurement of oxygen level is possible, then avoid hypoxia and hyperoxia	100% FiO ₂ ; when able to reliably measure, titrate FiO ₂ to SpO ₂ 94%–98%
Ventilation	PaCO ₂ 35–45 mm Hg	PaCO ₂ 35–45 mmHg	Normocapnia, possible mild hypercapnia ^b	EtCO ₂ 35–45 mmHg
Blood pressure	SBP > 90 mmHg MAP > 65 mmHg	MAP ≥ 65 mmHg Fluids, noradrenaline, and/or dobutamine depending on individual patient need for intravascular volume, vasoconstriction or inotropy	SBP > 100 mmHg	SBP ≥ 120 mmHg

Abbreviations: FiO₂, fraction of inspired oxygen; MAP, mean arterial pressure; PaCO₂, partial pressure of carbon dioxide; SBP, systolic blood pressure; SpO₂, peripheral oxygen saturation; SOS, Stabilize on Scene.

^aPatients who have had a brief period of cardiac arrest and an immediate return of normal cerebral function and are breathing normally may not require tracheal intubation but should be given oxygen via a facemask if their arterial blood oxygen saturation is less than 94%.

^bPer ILCOR, "These variations in methodology and in definitions of target ranges prohibit the task force from being able to recommend specific numbers."




FIGURE 2 Rates of rearrest and survival in out-of-hospital cardiac arrest by lowest recorded systolic blood pressure. Blue dotted line correlates with a systolic blood pressure (SBP) of 90 mmHg. Red dotted line correlates with a SBP of 120 mmHg. Adapted with permission from Smida et al.⁸

There remains some uncertainty over which blood pressure thresholds to target, as there are mixed results for different blood pressure targets.^{15,25,28,29} The most recent AHA guidelines recommend maintaining systolic blood pressure (SBP) above 90 mmHg and a mean arterial pressure (MAP) above 65 mmHg.¹⁴ Since hypotension is often defined as a SBP < 90 mmHg, this is frequently selected as an arbitrary binary threshold in outcome studies, which inform clinical guidelines, but the optimal threshold remains unknown. To date, randomized trials exploring higher blood pressure targets have not consistently demonstrated significant changes in survival, though higher blood pressure has been associated with better outcomes.^{15–18,30,31} One observational study of prehospital data suggested an improved prognosis with an SBP target between 100 and 130 mmHg¹⁰; a separate study demonstrated that maintaining SBP at 120 mmHg or above compared to a SBP below 90 mmHg as a trigger for action resulted in a 39% relative decrease in the incidence of rearrest and a 3.3-fold increase in survival (Figure 2).⁸

These findings are analogous to the evolving evidence for the prehospital care of patients with traumatic brain injury. The most recent guidelines for prehospital management of traumatic brain injury, which previously recommended a threshold of 90 mmHg, have increased the target SBP to ≥ 110 mmHg and new research suggests this threshold should be even higher.³² Patients resuscitated from cardiac arrest have the additional complication that a precipitous drop in blood pressure frequently occurs soon after ROSC as exogenous adrenaline is metabolized.²² Initiating treatment at a threshold of 90 mmHg will most often result in a period of marked hypotension prior to the patient's response. Therefore, a higher target seems appropriate.

Given the balance of evidence demonstrating potential benefit without data to suggest increased harm, and given the finding that rearrest is often preceded by a dropping blood pressure, the SOS-bundle uses an SBP cutoff of 120 mmHg for the initiation of vasopressor support. The SOS-bundle will standardize the approach to preventing and

addressing hypotension through a combination of intravenous fluids and early vasopressor administration. To prevent hypotension, all post-ROSC patients will be given a 500 mL bolus of crystalloid unless they have contraindications (eg., heart failure with signs of volume overload). Early detection will be facilitated by regular and frequent blood pressure checks (every 3 min), and early treatment will be facilitated by the routine (i.e., regardless of post-ROSC blood pressure) preparation of push-dose epinephrine (adrenaline) per local EMS agency protocol for all patients, in parallel with the fluid bolus. Patients with SBP < 120 mmHg after the initiation of the fluid bolus will be treated with push-dose epinephrine, with repeat doses titrated to maintain SBP. If the pressure is below the SBP threshold after the initial fluid bolus is complete, a second 500 mL bolus will be given, with a maximum total volume of 1000 mL. Push-dose epinephrine will be continued as needed every 1–5 min as needed until handoff of care in the ED. We note that EMS agencies implementing this bundle may choose to modify specific factors such as the dose and frequency (or even the SBP trigger or vasopressor selection) or may use a vasopressor infusion to achieve the same goals. We maintain that early targeted hemodynamic support is critical to preventing rearrest and improving outcomes and should be included in the post-ROSC bundle of care in some form.

3.2 | Circulation: STEMI identification

Cardiac arrest may be caused by coronary occlusion that would benefit from urgent percutaneous coronary intervention (PCI).^{33,34} EMS providers in the United States routinely use 12-lead electrocardiograms (ECGs) to identify patients with ST-segment elevation myocardial infarction in order to direct immediate field management and selection of the appropriate PCI-capable destination facility. In addition to those patients exhibiting STEMI on their ECGs, nearly one-third of patients that achieve ROSC have coronary lesions

that would benefit from PCI even when not exhibiting a definitive STEMI.³⁵

Post-ROSC patients with ECGs exhibiting ischemia benefit from direct transport to hospitals with emergency PCI capability.³⁶ In addition to obtaining the 12-lead ECG and paramedic interpretation, the ECG should be transmitted to an appropriate receiving facility EM physician. This has been shown to increase diagnostic accuracy and diminish delays to initiation of interventional treatment and reduction in infarct size, which may result in improved functional recovery upon discharge of post-cardiac arrest patients.^{37,38}

The optimal time to obtain the ECG during post-ROSC care is still under investigation.³⁶ Some research suggests that waiting 8 min after obtaining ROSC before performing a 12-lead ECG will more accurately reflect a true STEMI event due to the possibility of false-positive STEMI patterns that are due to cardiac arrest ischemia rather than coronary artery occlusion.³⁹ Furthermore, there are data that suggest repeated ECGs and rhythm checks are more likely to pick up STEMIs in chest pain patients than a single ECG (at least 8% in one study), suggesting that serial monitoring may be more useful than a single diagnostic snapshot.⁴⁰

Based on these data, we conclude that obtaining a 12-lead ECG and transmitting it to the appropriate receiving facility emergency medicine physician is an important part of the SOS-bundle. Acknowledging the lack of definitive data for timing of the ECG during post-ROSC care, the SOS-bundle recommends that the ECG is obtained after ROSC while still on scene. In addition to obtaining a 12-lead ECG, patients will be placed on continuous cardiac monitor to monitor for dysrhythmias.⁴⁰ Ideally patients post-ROSC would be transported to a PCI-capable facility based on local resources and policies.

3.3 | Airway: Airway management

Effective airway management to support oxygenation and ventilation in patients with OHCA is critical to post-ROSC stabilization and survival.⁴¹⁻⁴⁴ How this is accomplished, including by what technique and device, remains controversial.⁴⁴⁻⁴⁸ Classic teaching regarded intubation as the gold standard for cardiac arrest; however, recent studies and guidelines emphasize the importance of provider experience over specific modalities for best outcomes.⁴⁹ When comparing bag-valve-mask (BVM) use, supraglottic insertion, or endotracheal intervention, studies show conflicting data for which approach is most successful, and this difference is generally linked to provider experience, training, and preference rather than to the intervention itself.^{50,49}

Importantly, relying on BVM alone presents significant challenges to maintaining effective oxygenation, which is made even more difficult during patient movement and ambulance transport.⁵¹ Further, an advanced airway facilitates more consistent and accurate measures of EtCO₂ to apply the ventilation target. Thus, the SOS-bundle recommends placement of a supraglottic device or endotracheal tube prior to transport, according to local guidelines and EMS provider experience. Additionally, the SOS-bundle follows current best practices, stating

that patients should be preoxygenated and every effort made to allow for best first pass success, with concurrent use of passive oxygenation with nasal cannula when possible.⁵² Advanced airway placement will be confirmed by EtCO₂ and continuous capnography monitoring will take place to ensure early detection of displacement.

3.4 | Breathing: Oxygenation

Hypoxia is both a symptom of and a cause of cardiac arrest; if not reversed quickly it contributes to decreased survival and poor neurological outcomes.⁵³⁻⁵⁵ Because of this clear association, guidelines recommend 100% oxygen for post-ROSC patients until SpO₂ can be measured, at which time oxygen delivery should be titrated down to avoid the incidence of hyperoxia and its associated harms.^{5,50,56,57}

Current guidelines for post-resuscitation care advocate the targeting of normoxia for post-ROSC oxygenation.⁵⁶ Normoxia range is defined as 94%–98% as per the European Resuscitation Council (ERC) and 92%–98% as per the AHA.^{24,25,56} Evidence in favor of these ranges is still evolving. The recently published Reduction of Oxygen After Cardiac Arrest (EXACT) trial is the first trial conducted in the prehospital setting comparing the standard oxygen saturation target of 98%–100% measured by colorimetric pulse oximetry to a treatment strategy targeting 90%–94%.¹³ This study showed a higher likelihood of survival for patients with the higher oxygen goal. Importantly, use of a lower pulse oximetry target has potential to perpetuate disparities in survival from OHCA due to overestimation of arterial oxygenation in dark complexion patients because pulse oximetry can read falsely high in patients with darker skin.⁵⁸

While further studies are warranted, the current evidence supports titrating oxygen in post-ROSC patients in the prehospital setting to avoid both hypoxia and hyperoxia.^{54,55,50} Acknowledging a lack of definitive evidence on target SpO₂ range, the SOS-bundle adopts the ILCOR range of 94%–98%, rather than the lower AHA range, because this threshold could (i) help buffer against brief periods of hypoxia that may be more common when using 92% as a lower threshold and (ii) protect against hypoxia in patients with darker skin with falsely high pulse oximetry measurements. In patients with unreliable SpO₂ measures or other deteriorating vital signs, a 100% FiO₂ will be administered, with titration per the SOS-bundle protocol if monitoring is adequate or patient clinical condition subsequently improves.

3.5 | Breathing: Ventilation

Hyperventilation has been recognized as an important cause of poor outcomes in cardiac arrest patients for almost two decades.^{59,60} The potential harms of hyperventilation include overinflation of the lungs resulting in barotrauma, air trapping, increased intra-thoracic pressure, decreased central venous return, and decreased cardiac output, as well as risk of coronary artery constriction, cerebral artery constriction, and gastric insufflation.⁶⁰⁻⁶⁴ These complications can be caused by either excessive volume and/or excessive rate of assisted ventilations.

TABLE 2 Mitigation of hyperventilation with control of ventilation rate and volume.

Intervention	Helps control rate	Helps control tidal volume delivered
Strategy		
Education and training (i.e., “squeeze, release, release”; simulation-based training with feedback)	X	X
Synchronous compression to ventilation ratio	X	
Guiding ventilation to compression count	X	
Devices		
Auditory and visual prompt devices (e.g., metronomes and rate timing lights)	X	
Using smaller resuscitator volume bag-mask devices (e.g., pediatric vs. adult bag device)		X
Capnography	X	
Thoracic impedance waveform	X	
Ventilation feedback flowmeters	X	X
Mechanical ventilation	X	X

Source: Adapted from tab. 1 (Carlson et al.^{61,p.57}).

This harm can occur in the initial cardiac arrest management and/or in post-arrest care.

Until recently, the importance of adult cardiac arrest ventilation and its potential benefits and risks have not been emphasized in training of EMS personnel.^{60,61} In practice, assisted ventilation during resuscitation—both intra-arrest and post-ROSC—often exceeds these parameters.^{24,60,62,65–67} Current strategies to avoid intra-arrest hyperventilation emphasize using synchronous compression-to-ventilation ratios.^{14,26,68} Several supplemental approaches to optimizing ventilation have also been evaluated, with provider education and training strategies having been shown to assist in appropriate patient ventilation (Table 2).⁶¹ For example, CPR metronome training has shown benefit in avoiding over-ventilation, while simulation training shows promise in minimizing deviance from AHA guidelines and training EMS clinicians to state “squeeze” just until chest rise is initiated and then state “release, release” to allow time for exhalation may control both rate and volume.^{69–71} Equipment-based strategies to limit excessive tidal volumes have also been examined, including volume-restricted bags (e.g., using pediatric-sized BVMs in adult patients) and BVM flowmeters.^{61,72} While conceptually promising, evidence of benefit of such devices at this time is mostly limited to simulation settings.⁷² Relatedly, guidelines do not offer instruction on post-ROSC ventilation mechanism, and manual or mechanical ventilation will be used according to local practice.

As with intra-arrest care, training and equipment-based strategies to minimize hyperventilation in the post-ROSC period are conceptually

appealing but not well studied in field settings. Real-time monitoring of ventilation quality in cardiac arrest resuscitation is usually assessed using end-tidal carbon dioxide (EtCO₂) measurement, with a standard post-ROSC target of normocapnia (EtCO₂ between 35 and 45 mmHg).¹⁴ EtCO₂ is also frequently used to evaluate correct advanced airway.¹⁴ Despite the common use of EtCO₂, it is not universally available and, when available, several challenges in using EtCO₂ to guide ventilation strategies remain.^{61,65} Varied etiologies of cardiac arrest, resuscitation duration, use of vasoactive medications and sodium bicarbonate, and oscillations caused by chest compressions may result in changes in the pattern and values of EtCO₂, which make any single value or waveform difficult to determine the quality of the ventilation.⁷³ Newer strategies to mitigate these factors include use of waveform algorithms to better detect the presence of ventilation; however, these programs are not currently widely used.^{74,75} Despite its shortcomings, EtCO₂ is still used as the reference standard for ventilation monitoring in the field during cardiac arrest and post-ROSC care.

Of note, several trials of hypercapnia during the post-ROSC phase of cardiac arrest (PCO₂ 50–55 mmHg) have showed some promise in adult patients admitted to the intensive care unit.^{76,77} However, in a systematic review by Holmberg et al., of seven trials and 36 observational studies evaluating oxygenation and ventilation targets after cardiac arrest, point estimates from these trials favored normocapnia, and a more recent randomized trial found no benefit to mild hypercapnia.^{77,41,42}

The SOS-bundle incorporates current post-ROSC guidelines of starting with a minimum of 10 ventilations per minute and titrating the rate to a target EtCO₂ of normocapnia (35–45 mmHg). While acknowledging ongoing research regarding the accuracy of EtCO₂ in providing real-time feedback on adequate ventilation during post-ROSC care and evolving evidence regarding EtCO₂ targets, we have chosen to include normocapnic EtCO₂ targets in the SOS-bundle as it is the current reference standard method for assessing patient response to ventilation in the field.

4 | CONCLUSION

The goal-directed bundle of care presented here was developed by iterative consensus by the CAL-ROC, with the intended therapeutic goals of preventing hypotension, hypoxia, and hyperventilation in patients with ROSC following out-of-hospital cardiac arrest, and the intended outcome goals of reducing rearrest and improving survival and good neurologic outcomes. Although post-arrest management of these parameters is an evolving field, with ongoing evaluations of association and causality, our protocol incorporates the most up-to-date evidence and guidelines and is refined by expert opinion. While details regarding optimal management remain to be determined, our focus herein is to establish early goal-directed management to stabilize patients and prevent further physiologic insults. We believe that this bundle should be initiated in the immediate post-ROSC period and prior to transport in order to maximize the chances of

preventing decompensation and rearrest. Our future work will implement this bundle across multiple sites to measure feasibility of implementation, success in bundle targets, and—most importantly—improvements in patient outcomes.

AUTHOR CONTRIBUTIONS

Conceptualization: Nichole Bosson, Joelle Donofrio-Odmann, Marianne Gausche-Hill, Juan Carlos C. Montoy, and James J. Menegazzi. **Investigation:** All authors. **Methodology:** Nichole Bosson, Joelle Donofrio-Odmann, Marianne Gausche-Hill, Juan Carlos C. Montoy, and James J. Menegazzi. **Roles/writing—original draft:** all authors. **Writing—review and editing:** All authors. All authors contributed to the conception and design of the study, drafted portions of the article and provided it critical revisions for important intellectual content, give final approval of the version to be submitted, and agree to be accountable for all aspects of the work.

ACKNOWLEDGMENTS

We would like to thank the following CAL-ROC members for their support of this project: Joanny All, Peter Benson, Christian Bey, Lily Choi, Rosie Christ, Chris Coyne, Leslie Duke, Anthony Fabio, Catherine Farrokhi, Erica Garcia, Hernando Garzon, Gagandeep Grewal, Antonio Gumucio, Loreen Gutierrez, Caroline Jack, Chris Kahn, Kristi Koenig, Lisa Madrid, Tammi McConnell, Kenneth Miller, Daniel Nishijima, Sara Rasnake, John Rose, David Salcido, Daniel Shepherd, Pranav Shetty, Joshua Smith, Carl Schultz, and Reza Vaezazizi. The authors declare that all methods are available within the article; all supporting data are available in the cited references.

CONFLICT OF INTEREST STATEMENT

James J. Menegazzi was supported in part by a grant from the National Heart, Lung, and Blood Institute, National Institutes of Health (NHLBI 1R01HL117979-01) and David G Dillon was supported in this work by NHLBI grant K38HL165363. They have three patents issued and has been loaned equipment for use in his lab. These are described in more detail in his conflicts of interest form. The other authors declare no conflicts of interest.

REFERENCES

1. CARES. 2021 Annual report. Accessed December 9, 2022. https://mycares.net/sitepages/uploads/2022/2021_flipbook/index.html?page=1
2. Lerner EB, O'Connell M, Pirralo RG. Rearrest after prehospital resuscitation. *Prehosp Emerg Care*. 2011;15(1):50-54. doi:10.3109/10903127.2010.519820
3. Salcido DD, Schmicker RH, Kime N, et al. Effects of intra-resuscitation antiarrhythmic administration on rearrest occurrence and intra-resuscitation ECG characteristics in the ROC ALPS trial. *Resuscitation*. 2018;129:6-12. doi:10.1016/j.resuscitation.2018.05.028
4. Salcido DD, Stephenson AM, Condlie JP, Callaway CW, Menegazzi JJ. Incidence of rearrest after return of spontaneous circulation in out-of-hospital cardiac arrest. *Prehosp Emerg Care*. 2010;14(4):413-418. doi:10.3109/10903127.2010.497902
5. Salcido DD, Schmicker RH, Buick JE, et al. Compression-to-ventilation ratio and incidence of rearrest—a secondary analysis of the ROC CCC trial. *Resuscitation*. 2017;115:68-74. doi:10.1016/j.resuscitation.2017.04.007
6. Salcido DD, Sundermann ML, Koller AC, Menegazzi JJ. Incidence and outcomes of rearrest following out-of-hospital cardiac arrest. *Resuscitation*. 2015;86:19-24. doi:10.1016/j.resuscitation.2014.10.011
7. Spigner MF, Benoit JL, Menegazzi JJ, McMullan JT. Prehospital protocols for post-return of spontaneous circulation are highly variable. *Prehosp Emerg Care*. 2021;25(2):191-195. doi:10.1080/10903127.2020.1754979
8. Smida T, Menegazzi JJ, Crowe RP, Weiss LS, Salcido DD. Association of prehospital hypotension depth and dose with survival following out-of-hospital cardiac arrest. *Resuscitation*. 2022;180:99-107. doi:10.1016/j.resuscitation.2022.09.018
9. Smida T, Menegazzi JJ, Crowe RP, Bardes J, Scheidler JF, Salcido DD. Association of prehospital post-resuscitation peripheral oxygen saturation with survival following out-of-hospital cardiac arrest. *Resuscitation*. 2022;181:28-36. doi:10.1016/j.resuscitation.2022.10.011
10. Javaudin F, Desce N, Le Bastard Q, et al. Impact of pre-hospital vital parameters on the neurological outcome of out-of-hospital cardiac arrest: results from the French National Cardiac Arrest Registry. *Resuscitation*. 2018;133:5-11. doi:10.1016/j.resuscitation.2018.09.016
11. Lacocque J, Siegel L, Sporer KA. Prehospital, post-ROSC blood pressure and associated neurologic outcome. *Am J Emerg Med*. 2021;49:195-199. doi:10.1016/j.ajem.2021.05.073
12. Schneider AG, Eastwood GM, Bellomo R, et al. Arterial carbon dioxide tension and outcome in patients admitted to the intensive care unit after cardiac arrest. *Resuscitation*. 2013;84(7):927-934. doi:10.1016/j.resuscitation.2013.02.014
13. Bernard SA, Bray JE, Smith K, et al. Effect of lower vs higher oxygen saturation targets on survival to hospital discharge among patients resuscitated after out-of-hospital cardiac arrest: the EXACT randomized clinical trial. *JAMA*. 2022;328(18):1818-1826. doi:10.1001/jama.2022.17701
14. Panchal AR, Bartos JA, Cabanas JG, et al. Part 3: adult basic and advanced life support: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2020;142(16_suppl_2):S366-S468. doi:10.1161/CIR.0000000000000916
15. Kjaergaard J, Moller JE, Schmidt H, et al. Blood-pressure targets in comatose survivors of cardiac arrest. *N Engl J Med*. 2022;387(16):1456-1466. doi:10.1056/NEJMoa2208687
16. Jakkula P, Pettila V, Skrifvars MB, et al. Targeting low-normal or high-normal mean arterial pressure after cardiac arrest and resuscitation: a randomised pilot trial. *Intensive Care Med*. 2018;44(12):2091-2101. doi:10.1007/s00134-018-5446-8
17. Grand J, Meyer AS, Kjaergaard J, et al. A randomised double-blind pilot trial comparing a mean arterial pressure target of 65 mm Hg versus 72 mm Hg after out-of-hospital cardiac arrest. *Eur Heart J Acute Cardiovasc Care*. 2020;9(4_suppl):S100-S109. doi:10.1177/2048872619900095
18. Bray JE, Hein C, Smith K, et al. Oxygen titration after resuscitation from out-of-hospital cardiac arrest: a multi-centre, randomised controlled pilot study (the EXACT pilot trial). *Resuscitation*. 2018;128:211-215. doi:10.1016/j.resuscitation.2018.04.019
19. Chiu YK, Lui CT, Tsui KL. Impact of hypotension after return of spontaneous circulation on survival in patients of out-of-hospital cardiac arrest. *Am J Emerg Med*. 2018;36(1):79-83. doi:10.1016/j.ajem.2017.07.019
20. Kilgannon JH, Roberts BW, Reihl LR, et al. Early arterial hypotension is common in the post-cardiac arrest syndrome and associated with increased in-hospital mortality. *Resuscitation*. 2008;79(3):410-416. doi:10.1016/j.resuscitation.2008.07.019
21. Laurikkala J, Wilkman E, Pettila V, et al. Mean arterial pressure and vasopressor load after out-of-hospital cardiac arrest: associations

- with one-year neurologic outcome. *Resuscitation*. 2016;105:116-122. doi:[10.1016/j.resuscitation.2016.05.026](https://doi.org/10.1016/j.resuscitation.2016.05.026)
22. Trzeciak S, Jones AE, Kilgannon JH, et al. Significance of arterial hypotension after resuscitation from cardiac arrest. *Crit Care Med*. 2009;37(11):2895-2903; quiz 2904. doi:[10.1097/ccm.0b013e3181b01d8c](https://doi.org/10.1097/ccm.0b013e3181b01d8c)
 23. Berg KM, Soar J, Andersen LW, et al. Adult advanced life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation*. 2020;142(16_suppl_1):S92-S139. doi:[10.1161/CIR.0000000000000893](https://doi.org/10.1161/CIR.0000000000000893)
 24. Nolan JP, Sandroni C, Bottiger BW, et al. European Resuscitation Council and European Society of Intensive Care Medicine guidelines 2021: post-resuscitation care. *Intensive Care Med*. 2021;47(4):369-421. doi:[10.1007/s00134-021-06368-4](https://doi.org/10.1007/s00134-021-06368-4)
 25. Soar J, Nolan JP, Bottiger BW, et al. European Resuscitation Council guidelines for resuscitation 2015: section 3. Adult advanced life support. *Resuscitation*. 2015;95:100-147. doi:[10.1016/j.resuscitation.2015.07.016](https://doi.org/10.1016/j.resuscitation.2015.07.016)
 26. Wyckoff MH, Singletary EM, Soar J, et al. 2021 International consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; neonatal life support; education, implementation, and teams; first aid task forces; and the COVID-19 Working Group. *Resuscitation*. 2021;169:229-311. doi:[10.1016/j.resuscitation.2021.10.040](https://doi.org/10.1016/j.resuscitation.2021.10.040)
 27. Toy J, Tolles J, Bosson N, et al. Association between a post-resuscitation care bundle and the odds of field rearrest after successful resuscitation from out-of-hospital cardiac arrest: a pre/post study. *Prehosp Emerg Care*. 2023;1-9. doi:[10.1080/10903127.2023.2172633](https://doi.org/10.1080/10903127.2023.2172633)
 28. Ameloot K, De Deyne C, Eertmans W, et al. Early goal-directed haemodynamic optimization of cerebral oxygenation in comatose survivors after cardiac arrest: the neuroprotect post-cardiac arrest trial. *Eur Heart J*. 2019;40(22):1804-1814. doi:[10.1093/eurheartj/ehz120](https://doi.org/10.1093/eurheartj/ehz120)
 29. Jakkula P, Reinikainen M, Hastbacka J, et al. Targeting two different levels of both arterial carbon dioxide and arterial oxygen after cardiac arrest and resuscitation: a randomised pilot trial. *Intensive Care Med*. 2018;44(12):2112-2121. doi:[10.1007/s00134-018-5453-9](https://doi.org/10.1007/s00134-018-5453-9)
 30. Bray JE, Bernard S, Cantwell K, Stephenson M, Smith K, Committee VS. The association between systolic blood pressure on arrival at hospital and outcome in adults surviving from out-of-hospital cardiac arrests of presumed cardiac aetiology. *Resuscitation*. 2014;85(4):509-515. doi:[10.1016/j.resuscitation.2013.12.005](https://doi.org/10.1016/j.resuscitation.2013.12.005)
 31. Russo JJ, Di Santo P, Simard T, et al. Optimal mean arterial pressure in comatose survivors of out-of-hospital cardiac arrest: an analysis of area below blood pressure thresholds. *Resuscitation*. 2018;128:175-180. doi:[10.1016/j.resuscitation.2018.04.028](https://doi.org/10.1016/j.resuscitation.2018.04.028)
 32. Spaite DW, Hu C, Bobrow BJ, et al. Optimal out-of-hospital blood pressure in major traumatic brain injury: a challenge to the current understanding of hypotension. *Ann Emerg Med*. 2022;80(1):46-59. doi:[10.1016/j.annemergmed.2022.01.045](https://doi.org/10.1016/j.annemergmed.2022.01.045)
 33. Spaulding CM, Joly LM, Rosenberg A, et al. Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med*. 1997;336(23):1629-1633. doi:[10.1056/NEJM199706053362302](https://doi.org/10.1056/NEJM199706053362302)
 34. Jentzer JC, Clements CM, Wright RS, White RD, Jaffe AS. Improving survival from cardiac arrest: a review of contemporary practice and challenges. *Ann Emerg Med*. 2016;68(6):678-689. doi:[10.1016/j.annemergmed.2016.05.022](https://doi.org/10.1016/j.annemergmed.2016.05.022)
 35. Aufderheide TP, Engel TW, Saleh HO, et al. Change in out-of-hospital 12-lead ECG diagnostic classification following resuscitation from cardiac arrest. *Resuscitation*. 2021;169:45-52. doi:[10.1016/j.resuscitation.2021.10.012](https://doi.org/10.1016/j.resuscitation.2021.10.012)
 36. Alrawashdeh A, Nehme Z, Williams B, et al. Impact of emergency medical service delays on time to reperfusion and mortality in STEMI. *Open Heart*. 2021;8(1):e001654. doi:[10.1136/openhrt-2021-001654](https://doi.org/10.1136/openhrt-2021-001654)
 37. Tanguay A, Lebon J, Brassard E, Hebert D, Begin F. Diagnostic accuracy of prehospital electrocardiograms interpreted remotely by emergency physicians in myocardial infarction patients. *Am J Emerg Med*. 2019;37(7):1242-1247. doi:[10.1016/j.ajem.2018.09.012](https://doi.org/10.1016/j.ajem.2018.09.012)
 38. Kobayashi A, Misumida N, Aoi S, et al. STEMI notification by EMS predicts shorter door-to-balloon time and smaller infarct size. *Am J Emerg Med*. 2016;34(8):1610-1613. doi:[10.1016/j.ajem.2016.06.022](https://doi.org/10.1016/j.ajem.2016.06.022)
 39. Millin MG, Comer AC, Nable JV, et al. Patients without ST elevation after return of spontaneous circulation may benefit from emergent percutaneous intervention: a systematic review and meta-analysis. *Resuscitation*. 2016;108:54-60. doi:[10.1016/j.resuscitation.2016.09.004](https://doi.org/10.1016/j.resuscitation.2016.09.004)
 40. Tanguay A, Lebon J, Lau L, Hebert D, Begin F. Detection of STEMI using prehospital serial 12-lead electrocardiograms. *Prehosp Emerg Care*. 2018;22(4):419-426. doi:[10.1080/10903127.2017.1399185](https://doi.org/10.1080/10903127.2017.1399185)
 41. Eastwood G, Nichol AD, Hodgson C, et al. Mild hypercapnia or normocapnia after out-of-hospital cardiac arrest. *N Engl J Med*. 2023. doi:[10.1056/NEJMoa2214552](https://doi.org/10.1056/NEJMoa2214552)
 42. Grunau B, Kime N, Leroux B, et al. Association of intra-arrest transport vs continued on-scene resuscitation with survival to hospital discharge among patients with out-of-hospital cardiac arrest. *JAMA*. 2020;324(11):1058-1067. doi:[10.1001/jama.2020.14185](https://doi.org/10.1001/jama.2020.14185)
 43. O'Neill JF, Deakin CD. Do we hyperventilate cardiac arrest patients? *Resuscitation*. 2007;73(1):82-85. doi:[10.1016/j.resuscitation.2006.09.012](https://doi.org/10.1016/j.resuscitation.2006.09.012)
 44. Newell C, Grier S, Soar J. Airway and ventilation management during cardiopulmonary resuscitation and after successful resuscitation. *Crit Care*. 2018;22(1):190. doi:[10.1186/s13054-018-2121-y](https://doi.org/10.1186/s13054-018-2121-y)
 45. Jabre P, Penaloza A, Pinero D, et al. Effect of bag-mask ventilation vs endotracheal intubation during cardiopulmonary resuscitation on neurological outcome after out-of-hospital cardiorespiratory arrest: a randomized clinical trial. *JAMA*. 2018;319(8):779-787. doi:[10.1001/jama.2018.0156](https://doi.org/10.1001/jama.2018.0156)
 46. Wang HE, Schmicker RH, Daya MR, et al. Effect of a strategy of initial laryngeal tube insertion vs endotracheal intubation on 72-hour survival in adults with out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA*. 2018;320(8):769-778. doi:[10.1001/jama.2018.7044](https://doi.org/10.1001/jama.2018.7044)
 47. Bengier JR, Kirby K, Black S, et al. Effect of a strategy of a supraglottic airway device vs tracheal intubation during out-of-hospital cardiac arrest on functional outcome: the AIRWAYS-2 randomized clinical trial. *JAMA*. 2018;320(8):779-791. doi:[10.1001/jama.2018.11597](https://doi.org/10.1001/jama.2018.11597)
 48. Wnent J, Tjelmeland I, Lefering R, et al. To ventilate or not to ventilate during bystander CPR—a EuReCa TWO analysis. *Resuscitation*. 2021;166:101-109. doi:[10.1016/j.resuscitation.2021.06.006](https://doi.org/10.1016/j.resuscitation.2021.06.006)
 49. Dorsett M, Panchal AR, Stephens C, et al. Prehospital airway management training and education: an NAEMSP position statement and resource document. *Prehosp Emerg Care*. 2022;26(sup1):3-13. doi:[10.1080/10903127.2021.1977877](https://doi.org/10.1080/10903127.2021.1977877)
 50. Vincent JL, Taccone FS, He X. Harmful effects of hyperoxia in postcardiac arrest, sepsis, traumatic brain injury, or stroke: the importance of individualized oxygen therapy in critically ill patients. *Can Respir J*. 2017;2017:2834956. doi:[10.1155/2017/2834956](https://doi.org/10.1155/2017/2834956)
 51. Lyng JW, Guyette FX, Levy M, Bosson N. Prehospital manual ventilation: an NAEMSP position statement and resource document. *Prehosp Emerg Care*. 2022;26(sup1):23-31. doi:[10.1080/10903127.2021.1981506](https://doi.org/10.1080/10903127.2021.1981506)
 52. Gleason JM, Christian BR, Barton ED. Nasal cannula apneic oxygenation prevents desaturation during endotracheal intubation: an integrative literature review. *West J Emerg Med*. 2018;19(2):403-411. doi:[10.5811/westjem.2017.12.34699](https://doi.org/10.5811/westjem.2017.12.34699)
 53. Thomas M, Voss S, Bengier J, Kirby K, Nolan JP. Cluster randomised comparison of the effectiveness of 100% oxygen versus titrated oxygen in patients with a sustained return of spontaneous circulation following out of hospital cardiac arrest: a feasibility study. PROXY: post

- ROSC OXYgenation study. *BMC Emerg Med.* 2019;19(1):16. doi:[10.1186/s12873-018-0214-1](https://doi.org/10.1186/s12873-018-0214-1)
54. Thomas A, van Diepen S, Beekman R, et al. Oxygen supplementation and hyperoxia in critically ill cardiac patients: from pathophysiology to clinical practice. *JACC Adv.* 2022;1(3):100065. doi:[10.1016/j.jacadv.2022.100065](https://doi.org/10.1016/j.jacadv.2022.100065)
 55. Kilgannon JH, Jones AE, Shapiro NI, et al. Association between arterial hyperoxia following resuscitation from cardiac arrest and in-hospital mortality. *JAMA.* 2010;303(21):2165-2171. doi:[10.1001/jama.2010.707](https://doi.org/10.1001/jama.2010.707)
 56. Merchant RM, Topjian AA, Panchal AR, et al. Part 1: executive summary: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation.* 2020;142(16_suppl_2):S337-S357. doi:[10.1161/CIR.0000000000000918](https://doi.org/10.1161/CIR.0000000000000918)
 57. Berg KM, Cheng A, Panchal AR, et al. Part 7: systems of care: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation.* 2020;142(16_suppl_2):S580-S604. doi:[10.1161/CIR.0000000000000899](https://doi.org/10.1161/CIR.0000000000000899)
 58. Sjoding MW, Dickson RP, Iwashyna TJ, Gay SE, Valley TS. Racial bias in pulse oximetry measurement. *N Engl J Med.* 2020;383(25):2477-2478. doi:[10.1056/NEJMc2029240](https://doi.org/10.1056/NEJMc2029240)
 59. Pitts S, Kellermann AL. Hyperventilation during cardiac arrest. *Lancet.* 2004;364(9431):313-315. doi:[10.1016/S0140-6736\(04\)16740-8](https://doi.org/10.1016/S0140-6736(04)16740-8)
 60. Aufderheide TP, Lurie KG. Death by hyperventilation: a common and life-threatening problem during cardiopulmonary resuscitation. *Crit Care Med.* 2004;32(9 Suppl):S345-S351. doi:[10.1097/01.ccm.0000134335.46859.09](https://doi.org/10.1097/01.ccm.0000134335.46859.09)
 61. Carlson JN, Colella MR, Daya MR, et al. Prehospital cardiac arrest airway management: an NAEMSP position statement and resource document. *Prehosp Emerg Care.* 2022;26(sup1):54-63. doi:[10.1080/10903127.2021.1971349](https://doi.org/10.1080/10903127.2021.1971349)
 62. Vissers G, Duchatelet C, Huybrechts SA, Wouters K, Hachimi-Idrissi S, Monsieurs KG. The effect of ventilation rate on outcome in adults receiving cardiopulmonary resuscitation. *Resuscitation.* 2019;138:243-249. doi:[10.1016/j.resuscitation.2019.03.037](https://doi.org/10.1016/j.resuscitation.2019.03.037)
 63. Aufderheide TP, Sigurdsson G, Pirralo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation.* 2004;109(16):1960-1965. doi:[10.1161/01.CIR.0000126594.79136.61](https://doi.org/10.1161/01.CIR.0000126594.79136.61)
 64. Fritz C, Jaeger D, Luo Y, et al. Impact of different ventilation strategies on gas exchanges and circulation during prolonged mechanical cardiopulmonary resuscitation in a porcine model. *Shock.* 2022;58(2):119-127. doi:[10.1097/SHK.0000000000001880](https://doi.org/10.1097/SHK.0000000000001880)
 65. Neth MR, Idris A, McMullan J, Benoit JL, Daya MR. A review of ventilation in adult out-of-hospital cardiac arrest. *J Am Coll Emerg Physicians Open.* 2020;1(3):190-201. doi:[10.1002/emp2.12065](https://doi.org/10.1002/emp2.12065)
 66. Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during in-hospital cardiac arrest. *JAMA.* 2005;293(3):305-310. doi:[10.1001/jama.293.3.305](https://doi.org/10.1001/jama.293.3.305)
 67. Maertens VL, De Smedt LE, Lemoyne S, et al. Patients with cardiac arrest are ventilated two times faster than guidelines recommend: an observational prehospital study using tracheal pressure measurement. *Resuscitation.* 2013;84(7):921-926. doi:[10.1016/j.resuscitation.2012.11.015](https://doi.org/10.1016/j.resuscitation.2012.11.015)
 68. Hanisch JR, Counts CR, Latimer AJ, Rea TD, Yin L, Sayre MR. Causes of chest compression interruptions during out-of-hospital cardiac arrest resuscitation. *J Am Heart Assoc.* 2020;9(6):e015599. doi:[10.1161/JAHA.119.015599](https://doi.org/10.1161/JAHA.119.015599)
 69. Na JU, Han SK, Choi PC, Shin DH. Effect of metronome rates on the quality of bag-mask ventilation during metronome-guided 30:2 cardiopulmonary resuscitation: a randomized simulation study. *World J Emerg Med.* 2017;8(2):136-140. doi:[10.5847/wjem.j.1920-8642.2017.02.010](https://doi.org/10.5847/wjem.j.1920-8642.2017.02.010)
 70. Wayne DB, Didwania A, Feinglass J, Fudala MJ, Barsuk JH, McGaghie WC. Simulation-based education improves quality of care during cardiac arrest team responses at an academic teaching hospital: a case-control study. *Chest.* 2008;133(1):56-61. doi:[10.1378/chest.07-0131](https://doi.org/10.1378/chest.07-0131)
 71. Gausche M, Lewis RJ, Stratton SJ, et al. Effect of out-of-hospital pediatric endotracheal intubation on survival and neurological outcome: a controlled clinical trial. *JAMA.* 2000;283(6):783-790. doi:[10.1001/jama.283.6.783](https://doi.org/10.1001/jama.283.6.783)
 72. Kim JW, Park SO, Lee KR, Hong DY, Baek KJ. Efficacy of Amflow(R), a real-time-portable feedback device for delivering appropriate ventilation in critically ill patients: a randomised, controlled, cross-over simulation study. *Emerg Med Int.* 2020;2020:5296519. doi:[10.1155/2020/5296519](https://doi.org/10.1155/2020/5296519)
 73. Aramendi E, Elola A, Alonso E, et al. Feasibility of the capnogram to monitor ventilation rate during cardiopulmonary resuscitation. *Resuscitation.* 2017;110:162-168. doi:[10.1016/j.resuscitation.2016.08.033](https://doi.org/10.1016/j.resuscitation.2016.08.033)
 74. Gutierrez JJ, Leturiondo M, Ruiz de Gauna S, et al. Enhancing ventilation detection during cardiopulmonary resuscitation by filtering chest compression artifact from the capnography waveform. *PLoS One.* 2018;13(8):e0201565. doi:[10.1371/journal.pone.0201565](https://doi.org/10.1371/journal.pone.0201565)
 75. Ruiz de Gauna S, Leturiondo M, Gutierrez JJ, et al. Enhancement of capnogram waveform in the presence of chest compression artefact during cardiopulmonary resuscitation. *Resuscitation.* 2018;133:53-58. doi:[10.1016/j.resuscitation.2018.09.024](https://doi.org/10.1016/j.resuscitation.2018.09.024)
 76. Eastwood GM, Schneider AG, Suzuki S, et al. Targeted therapeutic mild hypercapnia after cardiac arrest: a phase II multi-centre randomised controlled trial (the CCC trial). *Resuscitation.* 2016;104:83-90. doi:[10.1016/j.resuscitation.2016.03.023](https://doi.org/10.1016/j.resuscitation.2016.03.023)
 77. Holmberg MJ, Nicholson T, Nolan JP, et al. Oxygenation and ventilation targets after cardiac arrest: a systematic review and meta-analysis. *Resuscitation.* 2020;152:107-115. doi:[10.1016/j.resuscitation.2020.04.031](https://doi.org/10.1016/j.resuscitation.2020.04.031)

How to cite this article: Dillon DG, Montoy JCC, Bosson N, et al. Rationale and development of a prehospital goal-directed bundle of care to prevent rearrest after return of spontaneous circulation. *JACEP Open.* 2024;5:e13321. <https://doi.org/10.1002/emp2.13321>