

Genetic disruption of dopamine β -hydroxylase dysregulates innate responses to predator odor in mice

Joyce Liu¹, Daniel J. Lustberg¹, Abigail Galvez, L. Cameron Liles, Katharine E. McCann, David Weinshenker*

Department of Human Genetics, Emory University School of Medicine, Atlanta, GA, USA

ARTICLE INFO

Handling editor: R Lawrence Reagan

Keywords:

Toxoplasma gondii
Norepinephrine
Dopamine
Dopamine β -hydroxylase
Predator

ABSTRACT

In rodents, exposure to predator odors such as cat urine acts as a severe stressor that engages innate defensive behaviors critical for survival in the wild. The neurotransmitters norepinephrine (NE) and dopamine (DA) modulate anxiety and predator odor responses, and we have shown previously that dopamine β -hydroxylase knockout (*Dbh* $-/-$), which reduces NE and increases DA in mouse noradrenergic neurons, disrupts innate behaviors in response to mild stressors such as novelty. We examined the consequences of *Dbh* knockout on responses to predator odor (bobcat urine) and compared them to *Dbh*-competent littermate controls. Over the first 10 min of predator odor exposure, controls exhibited robust defensive burying behavior, whereas *Dbh* $-/-$ mice showed high levels of grooming. Defensive burying was potently suppressed in controls by drugs that reduce NE transmission, while excessive grooming in *Dbh* $-/-$ mice was blocked by DA receptor antagonism. In response to a cotton square scented with a novel “neutral” odor (lavender), most control mice shredded the material, built a nest, and fell asleep within 90 min. *Dbh* $-/-$ mice failed to shred the lavender-scented nestlet, but still fell asleep. In contrast, controls sustained high levels of arousal throughout the predator odor test and did not build nests, while *Dbh* $-/-$ mice were asleep by the 90-min time point, often in shredded bobcat urine-soaked nesting material. Compared with controls exposed to predator odor, *Dbh* $-/-$ mice demonstrated decreased c-fos induction in the anterior cingulate cortex, lateral septum, periaqueductal gray, and bed nucleus of the stria terminalis, but increased c-fos in the locus coeruleus and medial amygdala. These data indicate that relative ratios of central NE and DA signaling coordinate the type and valence of responses to predator odor.

1. Introduction

1.1. Innate fear responses

Responding to external stimuli with appropriate behaviors (e.g., approach or avoidance) is crucial for survival of most animals. In rodents, detection of cat odorants prompts adaptive defensive and escape behaviors that reduce risk of predation. It is important to understand the neurobiological underpinnings of species-typical and maladaptive responses to threats because dysregulated behavioral responses to stressful stimuli represent many hallmark features of neuropsychiatric disorders, including posttraumatic stress disorder (PTSD), anxiety, and substance use disorder (Dias et al., 2013).

1.2. Norepinephrine modulates anxiety-like traits

The catecholamine neuromodulators norepinephrine (NE) and dopamine (DA) govern emotional and motivated behavior. NE is critical for arousal, attention, orienting to salient stimuli, and organization of stress responses (Poe et al., 2020; Sara 2009), while DA mediates reward and approach (Krach et al., 2010). The locus coeruleus (LC) is the main source of NE throughout the brain and projects to regions implicated in responses to potential threats (Lustberg et al. 2020a, 2020b; Poe et al., 2020; Schwarz and Luo 2015). Although NE is the dominant neurotransmitter produced by the LC, it can also release DA under certain conditions (Beas et al., 2018; Kempadoo et al., 2016; Petter et al., 2023; Takeuchi et al., 2016).

The LC is activated by novelty and predator odor, and NE transmission facilitates anxiety-like behavior, particularly in paradigms that

* Corresponding author.

E-mail address: dweinsh@emory.edu (D. Weinshenker).

¹ The first two authors contributed equally to this work.

involve stress and novelty (Curtis et al., 2012; Day et al., 2004; Hayley et al., 2001; Lustberg et al., 2020b). NE has been previously implicated in predator odor responses as well. Exposure to trimethylthiazoline (TMT), a component of fox odor, promotes freezing behavior in rats, which is attenuated by pharmacological blockade of NE transmission in the bed nucleus of the stria terminalis (BNST) (Fendt et al., 2005), and systemic or intra-dorsal premammillary nucleus β -adrenergic receptor antagonism abates defensive responses to cat odor (De Monte et al., 2008). Likewise, CRISPR-induced disruption of NE synthesis in the mouse LC blocks the ability of rat odor to trigger arousal and wakefulness (Yamaguchi et al., 2018).

1.3. Norepinephrine and dopamine control distinct stress-induced behaviors

We have previously used dopamine β -hydroxylase knockout (*Dbh* $-/-$) mice that lack NE and have excessive DA in the LC and other noradrenergic cell groups to investigate the relative contribution of these catecholamines to stress responses. The anxiogenic effects of novelty stress are abolished in *Dbh* $-/-$ mice, and they also fail to exhibit species-typical repetitive burying and nestlet shredding behaviors in novel environments (Lustberg et al. 2020a, 2020b). Pharmacological restoration of NE levels (without normalizing DA levels) rescues these phenotypes, while pharmacological blockade of NE transmission recapitulates them in normal mice. These findings indicate that the lack of NE is primarily responsible for the aberrant responses.

However, we recently reported that both NE and DA contribute to novel odorant stress (Lustberg et al., 2022). In response to non-threatening but novel olfactory stimuli (e.g. plant-derived essential oils presented in a novel environment), normal mice escalate digging, burrowing, and burying. In contrast, *Dbh* $-/-$ mice do not engage in these behaviors, and instead groom excessively in novel environments regardless of the presence of various “neutral” odors. Importantly, acute suppression of NE transmission blocks novel odor-induced digging in control mice, while DA receptor antagonists attenuate grooming in *Dbh* $-/-$ mice, indicating that the NE:DA ratio in noradrenergic neurons modulates the nature of these behavioral responses.

1.4. Study objectives and summary

While our previous work has provided insight concerning the roles of NE and DA in behavioral and brain responses to mild stressors such as novel environments and novel (but non-threatening) odors, the contribution of these catecholamines to severe innate threats has not been fully identified. Because loss of DBH uniquely decreases NE while simultaneously increasing DA, we speculated that *Dbh* $-/-$ mice would lack species-typical defensive responses to predator odor and potentially display paradoxical appetitive/approach behaviors towards it. To test this, we exposed *Dbh* $-/-$ and control mice to neutral odors or bobcat urine applied to cotton nestlets in a novel environment and measured defensive burying, grooming, nestlet shredding, and arousal. We then used a pharmacological approach to dissect the relative contributions of individual NE and DA receptor subtypes. We also assessed the effects of *Dbh* knockout on molecular signatures of neuronal activation in brain regions that receive innervation from the LC and are implicated in behavioral responses to predator odor.

2. Materials and methods

2.1. Animals

A total of 40 *Dbh* $-/-$ mice, maintained on a mixed 129/SvEv and C57BL/6 J background as previously described (Thomas et al. 1995, 1998), were used in this study. *Dbh* $-/-$ males were bred to *Dbh* $+/-$ females, and pregnant *Dbh* $+/-$ dams were administered drinking water containing the β -adrenergic receptor (AR) agonist isoproterenol

and the α 1AR agonist phenylephrine (20 μ g/ml each; Sigma-Aldrich) with vitamin C (2 mg/ml) from E9.5–E14.5, and the synthetic NE precursor L-3,4-dihydroxyphenylserine (DOPS; 2 mg/ml; Lundbeck, Deerfield, IL) + vitamin C (2 mg/ml) from E14.5-parturition to prevent embryonic lethality resulting from complete *Dbh* deficiency. *Dbh* $-/-$ mice are easily distinguished from their NE-competent littermates by their visible delayed growth and bilateral ptosis phenotypes. A total of 28 *Dbh* $+/-$ littermates were used as controls because their behavior and catecholamine levels are indistinguishable from wild-type (*Dbh* $+/+$) mice (Bourdelat-Parks et al., 2005; Szot et al., 1999; Thomas et al., 1998). Adult mice 4–10 months of age were used for all experiments.

Because no sex differences in stress-induced digging or grooming have been reported in past literature (Dixit et al., 2020; Londei et al., 1998, Smolinsky et al., 2009) or were observed in published experiments from our lab (Lustberg et al., 2022) and pilot experiments from this study, male and female mice from the same *Dbh* genotype were evenly distributed across drug treatment groups, and data were pooled between sexes. All animal procedures and protocols were approved by the Emory University Animal Care and Use Committee, in accordance with the National Institutes of Health guidelines for the care and use of laboratory animals. Mice were maintained on a 12 h/12 h light/dark cycle (7:00/19:00 h) with access to food and water ad libitum, except during behavioral testing. Behavioral testing was conducted under standard lighting conditions during the light cycle (14:00–17:00) in the same room where the mice were housed to minimize the stress of cage transport on test days.

2.2. Behavioral analysis

Mice were group-housed in static cages in same-sex groups of 2–5. In the first set of experiments, individual mice were removed from their home cages and placed into new standard mouse cages (13" \times 7" \times 6") containing only clean bedding substrate and a cotton nestlet square (2" \times 2", ~3 g) pre-soaked with either 1 ml of deionized water (odorless control) or predator odor (bobcat urine; Maine Outdoor Solutions, LLC, Hermon, Maine). Each mouse received both treatments in a counter-balanced order spaced 2 d apart. A clear plexiglass cover was pressed flush over the cages to prevent vaporization or dispersal of the odorants over the course of testing, while still allowing filming and behavioral scoring. The mice in the odorant-treated cages were filmed with a front-facing, mounted digital video camera for 10 min for assessment of defensive (digging/burying) and non-defensive (grooming) behaviors (De Boer and Koolhaas 2003; Kalueff and Tuohimaa 2005; Kemble and Bolwahn 1997; Londei et al., 1998; Lustberg et al., 2020a; Pond et al., 2021, Smolinsky et al., 2009). Time spent in defensive burying and grooming was manually scored with digital stopwatches by a trained observer blind to genotype and treatment. Defensive burying involved vigorous displacement of the bedding toward the odorant stimulus, often resulting in the burying of the odorant-soaked nestlet. Defensive burying sometimes presented as intense burrowing away from the stimulus, which resembled “swimming” or tunneling through the bedding substrate (Bondi et al., 2007; De Boer and Koolhaas 2003; Fucich and Morilak 2018; Lustberg et al., 2020a; Treit et al., 1981). Burying is considered aversion-related because it resembles responses exhibited after exposure to the shock probe test or predator odor (Garbe and Rawleigh, 1993, Hwa et al., 2019; McGregor et al., 2002; Sluyter et al., 1996). Grooming was defined as repetitive paw movements oriented around the whiskers and face, as well as licking or scratching of the body and tail (De Boer and Koolhaas 2003; Kalueff and Tuohimaa 2005, Smolinsky et al., 2009). Some reports suggest that grooming may be a self-soothing behavior that increases under stressful conditions, while others propose that it is a “rest behavior” that indicates low levels of stress (Guild and Dunn 1982; Kemble and Bolwahn 1997, Kemble and Gordon, 1995). For experiments with D1 receptor antagonist (see below), locomotor activity was measured as distanced travelled (m) using ANY-maze software (version 6.0, Stoelting Co, Wood Dale, IL,

USA).

The second set of experiments was identical to the first set of experiments, except that mice were pretreated with vehicle, adrenergic, or dopaminergic drugs (see below) prior to predator odor exposure. All drugs were administered 30 min prior to testing except nepicastat, which was administered 2 h prior to testing. Each mouse received both a vehicle and drug pretreatment in a counterbalanced order spaced 2 d apart. Each mouse received no more than two different drugs, and trials with the same mice were spaced at least two weeks apart.

The third set of experiments were identical to the first set of experiments except that (1) the cotton nestlet square pre-soaked with either 1 ml of a novel “neutral” odor (lavender oil, 1:100 dilution in deionized water; Anjou, Fremont, CA) or bobcat urine, and (2) mice were left undisturbed for 90 min, after which a trained experimenter returned to record whether the mouse was asleep and whether the nestlet was fully (100 %) shredded. Sleep was assessed behaviorally using EEG-validated scoring criteria, as we have described (Porter-Stransky et al., 2019).

2.3. Drugs

The following drugs were intraperitoneally administered to dissect the relative contributions of NE and DA receptor signaling to behavior: nepicastat (100 mg/kg, DBH inhibitor) (Synosia Therapeutics, Basel, Switzerland); prazosin (0.5 mg/kg, α 1AR antagonist) (Sigma-Aldrich, St. Louis, MO); propranolol (5 mg/kg, β AR antagonist) (Sigma-Aldrich); atipamezole (1 mg/kg, α 2AR antagonist) (Sigma-Aldrich), flupenthixol (0.25 mg/kg, nonspecific DA receptor antagonist) (Sigma-Aldrich); SCH-23390 (0.03 mg/kg, D1 receptor antagonist) (Sigma-Aldrich); L-741,626 (10 mg/kg, D2 receptor antagonist) (Sigma-Aldrich). Doses selected were based on previous studies and pilot experiments to control for confounding effects such as motor impairment (Lustberg et al. 2020a, 2022; Mitchell et al., 2008; Pina and Cunningham 2014; Yan et al., 2016). Because the 0.03 mg/kg dose of SCH-23390 suppressed general motor activity, we also tested 0.01 and 0.006 mg/kg doses. All drugs were dissolved in bacteriostatic saline except prazosin and L-741,626. Prazosin was first dissolved in 1.5 % DMSO and 1.5 % Cremophor EL before being added to saline, and L-741,626 was first dissolved in 10 % ethanol and 1.5 % Cremophor EL before being added to saline. The corresponding vehicle that each drug was dissolved in was used as a control.

2.4. *c-fos* immunohistochemistry (IHC)

Dbh +/- and *Dbh* -/- mice were exposed to predator odor for 90 min as described above, then euthanized with an overdose of sodium pentobarbital (Fatal Plus, 150 mg/kg, i.p.; Med-Vet International, Mettawa, IL) for transcardial perfusion with cold 4% paraformaldehyde in 0.01 M PBS. After extraction, brains were post-fixed for 24 h in 4 % paraformaldehyde at 4°C, and then transferred to cryoprotectant 30% sucrose/PBS solution for 72 h at 4°C. Brains were embedded in OCT medium (Tissue-Tek; Sakura, Torrance, CA) and serially sectioned by cryostat (Leica) into 20- μ m coronal slices. Brain sections were stored in 0.01 M PBS (0.02 % sodium azide) at 4 °C before IHC.

For IHC, brain sections were blocked for 1 h at room temperature in 5% normal goat serum (NGS; Vector Laboratories, Burlingame, CA) diluted in 0.01 M PBS/0.1 % Triton-X permeabilization buffer. Sections were then incubated for 48 h at 4°C in NGS blocking/permeabilization buffer, including primary antibodies raised against *c-fos* (rabbit anti-*c-fos*, Millipore, Danvers, MA, ABE457; 1:5000) and the NE transporter (NET; chicken anti-NET, #260006, Synaptic System, Goettingen, Germany; 1:3000). After washing in 0.01 M PBS, sections were incubated for 2 h in blocking/permeabilization buffer with goat anti-rabbit AlexaFluor 488 and goat anti-chicken AlexaFluor 568 (Invitrogen, Carlsbad, CA; 1:500). After washing, the sections were mounted onto Superfrost Plus slides and coverslipped with Fluoromount-G plus DAPI (Southern Biotech, Birmingham, AL).

2.5. Statistical analysis

The effects of the water vs predator odor on defensive burying and grooming in *Dbh* +/- and *Dbh* -/- mice were compared using repeated measures 2-way ANOVA (genotype x odorant), with post hoc Tukey tests for multiple comparisons where appropriate. The effects of drugs vs vehicle on time spent engaged in repetitive behaviors or locomotor activity were assessed using repeated measures 1-way ANOVA with post hoc Dunnett's multiple comparison tests (SCH-23390) or paired sample t-tests (all other drugs). Contingency data of nestlet shredding and sleeping between genotypes were analyzed using Fisher's exact tests. Although the current study was not sufficiently powered to detect sex differences, preliminary analysis of the data revealed roughly equivalent distributions and means between males and females (data not shown), similar to what we previously reported for behavioral and neurochemical responses to novel odors (Lustberg et al., 2022). Thus, male and female data were pooled within each genotype and treatment group.

3. Results

3.1. *Dbh* -/- mice exhibit decreased defensive burying and increased grooming in response to predator odor

We first assessed the responses of *Dbh* -/- and NE-competent *Dbh* +/- control mice to predator odor and water by measuring defensive burying (Fig. 1A, Supplemental movie 1) and grooming (Fig. 1B, Supplemental movie 1). Control animals displayed modest levels of grooming during water exposure and no defensive burying. Predator odor dramatically increased defensive burying and total digging but had




Fig. 1. Effects of DBH knockout on behavioral responses to predator odor. *Dbh* +/- (n = 7) and *Dbh* -/- (n = 7) mice were exposed to water or bobcat urine for 10 min, and the amount of time engaged in defensive burying (A) and grooming (B) were assessed. Shown are mean +/- SEM. ***p < 0.001, ****p < 0.0001.

no impact on grooming in control mice. By contrast, *Dbh* $-/-$ mice displayed high levels of grooming in response to water and were behaviorally indifferent to predator odor, with no emergence of defensive burying and no change in grooming.

For defensive burying, there were main effects of genotype [$F(1,24) = 15.96, p = 0.0005$], odorant [$F(1,24) = 15.96, p = 0.0005$], and odorant \times genotype interaction [$F(1,24) = 15.96, p = 0.0005$]. Post hoc analysis revealed that *Dbh* $+/-$ mice showed significantly increased defensive burying ($p < 0.0001$) in the predator odor condition compared with water, and they also engaged in significantly more defensive burying ($p < 0.0001$) than the *Dbh* $-/-$ mice in the predator odor condition. For grooming, there was a main effect of genotype [$F(1,24) = 18.82, p = 0.002$], but no main effect of odorant [$F(1,24) = 0.7293, p = 0.40$] or odorant \times genotype interaction [$F(1,24) = 0.57, p = 0.46$].

3.2. Adrenergic antagonists suppress predator odor-induced digging behavior in control animals

Our data from *Dbh* $-/-$ mice suggested that NE is required for defensive burying in response to predator odor. To determine whether this is true in NE-competent mice as well as which receptors are involved, we next assessed the effects of specific AR antagonists on the predator odor responses of *Dbh* $+/-$ mice. The compounds tested included prazosin (0.5 mg/kg, α 1AR antagonist), atipamezole (1 mg/kg, α 2AR antagonist), and propranolol (5 mg/kg, β AR antagonist). Compared with their respective vehicles, each compound tested significantly reduced burying (prazosin: $t(7) = 2.46, p = 0.044$; atipamezole: $t(6) = 3.38, p = 0.015$; propranolol: $t(7) = 2.85, p = 0.025$) (Fig. 2A–C), indicating that α 1AR, α 2AR, and β AR activation all contribute to this predator odor-induced defensive behavior. Prazosin was the only compound that attenuated grooming in control mice (Fig. 2D–F) (prazosin: $t(7) = 3.03, p = 0.019$; atipamezole: $t(6) = 0.74, p = 0.49$; propranolol: $t(7) = 0.85, p = 0.43$). Overall, these results




Fig. 2. Effects of adrenergic drugs on behavioral responses to predator odor in control mice. *Dbh* $+/-$ ($n = 8$) mice were pretreated with vehicle, the α 1AR antagonist prazosin (0.5 mg/kg), the α 2AR antagonist atipamezole (1 mg/kg), or the β AR antagonist propranolol (5 mg/kg) and exposed to bobcat urine 30 min later for 10 min. Shown is the mean \pm SEM time engaged in defensive burying (A–C) and grooming (D–F). * $p < 0.05$.

indicated that defensive burying was the predator odor-induced behavior most sensitive to blocking NE transmission.

3.3. Pharmacological DBH inhibition reduces defensive burying and increases grooming following exposure to predator odor in normal mice

Because AR antagonists recapitulated the decrease in digging behaviors but not the increase in grooming observed in the *Dbh*^{-/-} mice, we reasoned that increased grooming was mediated by ectopic DA produced by LC neurons in the knockouts. To test this idea, we used the selective DBH inhibitor nepicastat, which unlike AR antagonists increases DA in addition to decreasing NE (Devoto et al., 2015; Ohkubo et al., 2019; Schroeder et al., 2010; Stanley et al., 1997). Nepicastat administration to *Dbh*^{+/-} control mice phenocopied both aberrant responses of *Dbh*^{-/-} mice to predator odor [defensive burying: $t(9) = 3.29$, $p = 0.009$; grooming: $t(9) = 2.60$, $p = 0.029$] (Fig. 3), suggesting that increasing DA contributes to abnormal *Dbh*^{-/-} mouse grooming.

3.4. DA receptor antagonists attenuate excessive grooming in *Dbh*^{-/-} mice in the presence of predator odor

To directly test whether excessive DA transmission was required for grooming in *Dbh*^{-/-} mice, we assessed the effects of DA receptor antagonists on predator odor response in the knockouts. The drugs tested included flupenthixol (0.25 mg/kg, nonselective DA receptor

antagonist), L-741,626 (10 mg/kg, D2 receptor antagonist), and SCH-23390 (0.03 mg/kg, D1 receptor antagonist). Flupenthixol [$t(12) = 2.76$, $p = 0.0037$] (Fig. 4A) significantly reduced grooming in *Dbh*^{-/-} mice in the presence of predator odor, while L-741,626 had no effect [$t(7) = 0.75$, $p = 0.48$] (Fig. 4B). SCH-23390 completely eliminated grooming, but we also noted that mice administered SCH-23390 ceased nearly all movement. SCH-23390-treated mice were capable of moving and did so when gently prodded, but failed to ambulate volitionally. In an attempt to separate the effects of D1 blockade on grooming and general motor activity, we also tested two lower doses (0.01 and 0.006 mg/kg), and the results are shown in Fig. 4C and D. There was a main effect of dose on both grooming [$F(1.701, 11.90) = 40.42$, $p < 0.0001$], and locomotor activity [$F(1.226, 8.579) = 15.59$, $p = 0.0028$]. Post hoc analysis revealed that the 0.03 dose fully suppressed grooming [$q(7) = 12.66$, $p < 0.0001$] and locomotor activity [$q(7) = 8.48$, $p = 0.002$]. Similarly, there was almost no grooming following administration of the 0.01 mg/kg dose [$q(7) = 8.83$, $p = 0.0001$], while locomotor activity returned to ~50 % of saline control [$q(7) = 7.06$, $p = 0.005$]. Finally, the 0.006 mg/kg dose induced a borderline significant reduction of grooming [$q(7) = 2.81$, $p = 0.06$] with no effect on locomotor activity [$q(7) = 0.06$, $p = 0.99$]. These results suggest that excessive grooming in *Dbh*^{-/-} mice is mediated by D1, but not D2 transmission, although we cannot completely rule out a non-specific effect of SCH23390. Vehicle-treated *Dbh*^{-/-} mice exhibited almost no defensive burying (see Fig. 1A), and none of the dopaminergic antagonists altered this outcome (data not shown).

3.5. $\alpha 1AR$ blockade has no effect on behavioral response to predator odor in *Dbh*^{-/-} mice

Our data thus far have indicated that NE transmission was responsible for defensive burying, while DA promoted grooming. However, because we observed a significant reduction in grooming in the *Dbh*^{+/-} mice treated with prazosin, and DA can signal through $\alpha 1AR$ s under some conditions (Cilz et al., 2014; Ozkan et al., 2017; Paladini et al., 2001), we speculated that some of the excessive grooming observed in *Dbh*^{-/-} mice might be driven by DA-mediated $\alpha 1AR$ transmission. To test this hypothesis, we administered prazosin (0.5 mg/kg) to *Dbh*^{-/-} mice in the presence of predator odor, but failed to observe a drug effect on grooming (vehicle: 223.2 ± 49.85 s; prazosin 216.3 ± 45.56 s; $p = 0.8986$). Combined with the outcomes from the DA receptor antagonist experiments, these results suggest that while $\alpha 1AR$ s modestly contribute to predator odor-induced grooming in normal mice, they are not required for the excessive grooming response of *Dbh*^{-/-} mice, which appears to be mediated by excessive activation of D1 receptors.

3.6. *Dbh*^{-/-} mice lack predator odor-induced arousal and aversion

Predator odors typically elicit innate vigilance and aversion in prey that is resistant to habituation. We and others have shown that, when exposed to a novel environment and given access to a cotton nestlet, normal mice rapidly shred the nestlet, build a nest, and fall asleep in it (Lustberg et al., 2020a). To examine the effects of disrupting LC catecholamine homeostasis on these responses to predator odor, we measured nestlet shredding and arousal state in *Dbh*^{+/-} and *Dbh*^{-/-} mice after 90 min of exposure to a novel environment with a nestlet treated with bobcat urine. To control for the influence of a novel odor, a separate cohort of mice experienced the same paradigm except that a non-threatening but unfamiliar odor (lavender oil) was used instead of bobcat urine. In the presence of lavender, most (6/8) *Dbh*^{+/-} mice shredded the nestlet (Fig. 5A) and were asleep (Fig. 5B) in the nest they built by the 90 min time point. As we have shown previously with unscented nestlet (Lustberg et al., 2020a), *Dbh*^{-/-} mice failed to shred the nestlet (0/7) (Fig. 5A) but still fell asleep (6/7) (Fig. 5B). Fisher's exact tests revealed a significant difference between genotypes for shredding ($p = 0.007$) but not sleeping ($p > 0.99$).




Fig. 3. Effects of pharmacological DBH inhibition on behavioral responses to predator odor. *Dbh*^{+/-} mice ($n = 10$) were pretreated with vehicle or the DBH inhibitor nepicastat (100 mg/kg) and exposed to bobcat urine for 10 min, and the amount of time engaged in defensive burying (A) and grooming (B) were assessed. Shown are the mean \pm SEM. * $p < 0.05$, ** $p < 0.01$.




Fig. 4. Effects of dopaminergic drugs on predator odor-induced grooming and locomotor activity in *Dbh*^{-/-} mice. *Dbh*^{-/-} mice were pretreated with vehicle (*n* = 7–13), the non-selective DA receptor antagonist flupentixol (0.25 mg/kg; *n* = 7), the D2 antagonist L-741,626 (10 mg/kg; *n* = 8), or the D1 antagonist SCH-23390 (0.03, 0.01, or 0.006 mg/kg; *n* = 8), or and exposed to bobcat urine 30 min later for 10 min. Shown is the mean \pm SEM time engaged in grooming (A–C) and distance travelled (D). *****p* < 0.0001, ****p* < 0.001, ***p* < 0.01, ns = not significant.

In contrast to lavender odor, none (0/8) of the *Dbh*^{+/-} mice shredded their nestlet (Fig. 5C) or fell asleep (Fig. 5D) in the presence of bobcat urine. The behavior of *Dbh*^{-/-} mice also changed when exposed to predator odor, but in unexpected directions. While *Dbh*^{-/-} mice never engaged in shredding unscented or lavender-scented nestlets within 90 min, a majority (5/9) of them fully (100 %) shredded the bobcat urine-soaked nestlet (Fig. 5C). Those that did build a nest fell asleep in it, and altogether 8/9 of the *Dbh*^{-/-} mice were asleep by the end of the test (Fig. 5D). Fisher’s exact tests revealed significant differences between genotypes for shredding (*p* = 0.0294) and sleeping (*p* = 0.0004). Pilot experiments using different sources of predator odor (fox urine and tufts of domestic cat fur instead of bobcat urine) also revealed paradoxical appetitive responses in knockout mice; *Dbh*^{+/-} animals engaged in typical defensive burying behavior, while most *Dbh*^{-/-} mice instead sat on top of the cat fur and groomed themselves (Supplemental movie 2). These results demonstrate that predator odor elicits powerful arousal and aversion in normal mice, and that loss of DBH activity produces aberrant behavioral responses that include apparent affiliative responses.

3.7. *-fos* immunohistochemistry reveals predator odor-responsive brain regions that differ between *Dbh*^{+/-} and *Dbh*^{-/-} mice

To identify candidate brain regions that are activated by predator odor and facilitate differential responses in *Dbh*^{+/-} and *Dbh*^{-/-} mice, *c-fos* immunohistochemistry was conducted in animals exposed to a novel environment with a bobcat urine-treated nestlet. Analysis focused on the LC and other brain regions that receive innervation from noradrenergic neurons and have been implicated in predator odor responses, including the anterior cingulate cortex (ACC), dorsal BNST (dBNST), periaqueductal gray (PAG), periventricular nucleus of the hypothalamus (PVN), lateral septum (LS), and medial amygdala (MeA) (Fig. 6). Repeated measures 2-way ANOVA showed a main effect of brain region [*F* (3,45) = 76.57, *p* < 0.0001] and a genotype \times brain region interaction [*F* (6,90) = 44.84, *p* < 0.0001]. Sidak’s post hoc tests revealed that *Dbh*^{+/-} mice had significantly more *c-fos*⁺ cells than *Dbh*^{-/-} mice in the ACC (*t* = 8.361, *p* < 0.001), dBNST (*t* = 7.767, *p* < 0.0001), PAG (*t* = 9.897, *p* < 0.0001), and LS (*t* = 7.460, *p* < 0.001), while *Dbh*^{-/-} mice had significantly more *c-fos*⁺ cells in the LC (*t* = 9.554, *p* < 0.0001) and MeA (*t* = 3.679, *p* = 0.0177). No genotype




Fig. 5. Effects of DBH knockout on arousal and nestlet shredding during predator odor exposure. *Dbh* +/- and *Dbh* -/- mice were exposed to lavender oil (A-B; $n = 7,8$) or bobcat urine (C-D; $n = 8,9$) for 90 min, and the fraction of animals that shredded their cotton nesting material (A,C) and fell asleep (B,D) were assessed. * $p < 0.05$, ** $p < 0.01$.

differences were found in the PVN ($t = 0.956$, $p = 0.9542$). Integrated with the effects of genotype and drugs, these results imply that noradrenergic transmission in the ACC, dBNST, PAG, and/or the LS contribute to aversive/defensive effects of predator odor, while D1 signaling in the MeA promotes grooming and appetitive responses.

4. Discussion

4.1. Summary

Predator odors (or components thereof) such as bobcat urine and TMT are ethologically relevant stressors that elicit intense, innate behavioral responses in mice (Ferrero et al., 2011; Janitzky et al., 2015). In this study, we investigated the effects of manipulating catecholamine transmission on predator odor-induced stress in mice by comparing *Dbh* -/- mice, which lack NE and instead produce DA in their “noradrenergic” neurons, to their *Dbh* +/- littermates with normal catecholamine content. We also used pharmacological manipulation to target NE and DA signaling, allowing us to assess the effects of both chronic and acute catecholamine manipulation.

4.2. Adrenergic control of predator odor-induced defensive burying

Defensive burying is considered an active coping strategy in response to stress, in which the mouse attempts to remove or avoid aversive stimuli. In the shock probe defensive burying test, mice will use bedding to bury electrified prods and remove the source of shock pain (Tillage

et al., 2020). The marble burying test has also demonstrated that burying is stress-sensitive, with stressed mice burying more marbles than unstressed mice (Kedia and Chattarji 2014), and we have shown that *Dbh* -/- mice bury fewer marbles than controls (Lustberg et al., 2020a).

We found that both *Dbh* -/- and *Dbh* +/- mice engage in similarly low levels of defensive burying when exposed to a novel cage with a water-soaked cotton nestlet, indicating that this environment is not particularly stressful. *Dbh* +/- mice show increased defensive burying when exposed to predator odor as compared to water, suggesting that this behavior is an innate adaptive defensive response to threat. The combined results from our experiments with *Dbh* -/- mice, acute pharmacological DBH inhibition, and adrenergic receptor antagonists demonstrate that noradrenergic transmission is required for defensive burying during predator odor exposure. We previously showed that NE is critical for digging in the presence of novel odors but did not precisely identify the receptor subtypes responsible (Lustberg et al., 2022).

The effects of β AR and α 1AR activation are partially overlapping but can also be distinct. β AR activation is necessary for many types of anxiety-like and defensive behaviors, including responses to physical threats (e.g. shock) (Bondi et al., 2007), psychological threats (e.g. novel environments) (Lustberg et al., 2022), and anxiogenic drugs (e.g. cocaine) (Schank et al., 2008), while α 1AR signaling promotes both exploratory and aversive behaviors in novel environments (Lustberg et al., 2020a; Stone et al. 1999, 2006). α 1AR blockade decreases defensive behavior in the presence of a traumatic cue in rats (Ketenci et al., 2020), and antagonism of β ARs attenuates cocaine




Fig. 6. Regional c-fos responses to predator odor in *Dbh* +/- and *Dbh* -/- mice. *Dbh* +/- (n = 8) and *Dbh* -/- (n = 9) mice were exposed to bobcat urine for 90 min. Shown are (A) representative images and (B) the mean +/- SEM number of c-fos+ (magenta) cells in the locus coeruleus (LC), anterior cingulate cortex (ACC), dorsal bed nucleus of the stria terminalis (dBNST), periaqueductal gray (PAG), paraventricular nucleus of the hypothalamus (PVN), lateral septum (LS), and medial amygdala (MeA). Images are counterstained with the nuclear marker DAPI (blue) and the noradrenergic terminal marker norepinephrine transporter (NET; green). *p < 0.05, ***p < 0.001, ****p < 0.0001.

withdrawal-induced defensive burying in rats (Harris et al., 2018). Interestingly, the β AR antagonist propranolol blocks the anxiogenic effects of optogenetic LC stimulation in the elevated zero maze, while only the α 1AR antagonist prazosin prevents real-time place aversion (McCall et al., 2015). We have shown that propranolol diminishes nestlet shredding, while prazosin attenuates marble burying in a novel environment (Lustberg et al., 2020a). By contrast, antagonism of either receptor reduces defensive burying in the shock probe test. Thus, the current finding that both β ARs and α 1ARs contribute to defensive burying behavior during predator odor exposure is consistent with previous literature.

The ability of the α 2AR antagonist atipamezole to reduce predator odor-induced defensive burying was more of a surprise. α 2ARs are expressed widely throughout the brain but are enriched in LC neurons, where they function as inhibitory autoreceptors (Aghajanian and VanderMaelen, 1982; Timmermans and van Zwieten 1982). Thus, α 2AR agonists often decrease, while α 2AR antagonists can increase NE transmission and anxiety-like behaviors (Lustberg et al. 2020a, 2020b). In the current study, atipamezole had effects similar to those of propranolol and prazosin, suggesting that it was not primarily acting on α 2AR autoreceptors but rather by blocking α 2AR heteroreceptors on LC target cells. α 2ARs are expressed in stress-sensitive brain regions such as the prefrontal cortex (PFC, which includes the ACC) and the dBNST, where they can have paradoxical excitatory effects on neurons and networks (Harris et al., 2018; Wang et al., 2007). For example, stimulation of postsynaptic α 2ARs in the dBNST enhances neuronal activity and promotes anxiety-like behavior in the elevated plus maze (Harris et al., 2018). We speculate that atipamezole-induced reduction of defensive burying in response to predator odor is mediated by similar mechanisms.

4.3. Dopaminergic control of grooming

Grooming, unlike digging, is not typically considered a stress coping response, as mice will groom in situations of both comfort and distress (Smolinsky et al., 2009). Indeed, cat odor often inhibits grooming in rodents (McGregor et al., 2004). Repetitive, purposeless self-grooming is often observed in animal models of obsessive-compulsive disorder (OCD), Tourette's syndrome (TS), and autism spectrum disorder (ASD) (Lewis 2011; Nordstrom and Burton 2002; Welch et al., 2007). DA signaling has been previously implicated in facilitating grooming behavior. For example, D1, but not D2, receptor signaling contributes to stereotyped self-grooming in rats (Berridge & Aldridge 2000a, 2000b) and novelty-induced grooming in mice (Drago et al., 1999).

As a precursor for NE, DA is produced in noradrenergic neurons. Due to incomplete conversion to NE by DBH, DA can also be released by these cells and mediates some LC functions including specific forms of learning and memory (Chowdhury et al., 2022; Kempadoo et al., 2016; Wagatsuma et al., 2018). *Dbh*^{-/-} mice lack NE and instead exclusively release DA from LC neurons under conditions that normally result in NE transmission. Converging lines of evidence from our study indicate that the excessive grooming displayed by *Dbh*^{-/-} mice is mediated by DA. First, high levels of grooming in the presence of predator odor were observed in control mice treated with the DBH inhibitor nepicastat, but not in mice following AR blockade. Second, grooming behavior in *Dbh*^{-/-} mice was attenuated by DA receptor antagonists.

Our findings that the non-selective DA receptor antagonist flupentixol and the D1-selective antagonist SCH-23390, but not the selective D2 antagonist L-741,626 blocked grooming in *Dbh*^{-/-} mice exposed to predator odor indicates this behavior is mediated by D1 receptors. One caveat to this interpretation is that D1 blockade is well known to suppress many aspects of motor activity, including cocaine-induced hyperactivity (Cabib et al., 1991) and rearing in a novel cage (Dreher and Jackson 1989). Another study using SCH-23390 also showed that it can induce catalepsy (Chinen and Frussa-Filho 1999), although at a much higher dose (0.1 mg/kg) than the one we used. In our experiments, the effects of SCH-23390 on grooming were more potent than those on

locomotion, but there was no dose that significantly reduced grooming without effects on distance travelled; the closest was 0.006 mg/kg, which allowed normal locomotion while almost reaching significance for grooming ($p = 0.06$). Prazosin reduced grooming in *Dbh*^{+/-} but not *Dbh*^{-/-} mice, indicating that NE signaling through α 1ARs promotes this behavior exclusively in normal animals. This is consistent with our previous finding that prazosin reduces amphetamine-induced locomotion in *Dbh*^{+/-} but not *Dbh*^{-/-} mice (Weinshenker et al., 2002), and further implicates D1 receptor mediation in knockouts.

We previously showed that *Dbh*^{-/-} mice groom more than *Dbh*^{+/-} mice in the presence of a novel but non-threatening odorant, and this excessive grooming was reduced by administration of flupentixol (Lustberg et al., 2022). Therefore, the results from our genetic and pharmacological experiments suggest that the neurochemistry and circuitry that drive grooming in the presence of predator odor resemble those controlling grooming responses to novel odors and/or environments.

4.4. *Dbh*^{-/-} mice have impaired arousal and aversion responses to predator odor

Both NE and DA promote arousal. Activation of noradrenergic neurons in the LC and dopaminergic neurons in the ventral tegmental area (VTA) and ventral PAG (vPAG) induce wakefulness, while their suppression reduces arousal (Carter et al., 2010; Cho et al., 2017; Eban-Rothschild et al., 2016; Lu et al., 2006; Poe et al., 2020; Porter-Stransky et al., 2019). Many environmental challenges such as low dose amphetamine, white noise, and human handling that increase wakefulness in normal mice fail to do so in *Dbh*^{-/-} mice (Hunsley & Palmiter 2003, 2004), CRISPR-mediated knockout of DBH in the mouse LC attenuates arousal triggered by rat bedding (Yamaguchi et al., 2018), and AR antagonists can also reduce wakefulness in control mice (Mitchell and Weinshenker 2010; Porter-Stransky et al., 2019). Sustained arousal is an adaptive response to threats such as predator odor. Normal mice exposed to a new cage with an unscented cotton nestlet will initially explore the novel environment, then shred the material and fall asleep in the nest within 60–90 min (Lustberg et al., 2020a). In the current study, we showed that while most control mice shredded a cotton square containing a novel but non-threatening odor (lavender) and fell asleep in the nesting material, none shredded a bobcat urine-soaked nestlet and all remained vigilant for at least 90 min. *Dbh*^{-/-} mice had a paradoxical response to novel and threatening odors. The knockouts fail to shred odorless (Lustberg et al., 2020a) or lavender-scented (this study) material, but remarkably built a nest and fell asleep in it only when it contained bobcat urine. Because defensive burying and grooming responses to predator odor in *Dbh*^{-/-} mice did not differ from those to plain water, it was not initially clear whether *Dbh*^{-/-} mice could even detect bobcat urine. However, the nestlet result refutes that possibility, and odor sensation in general is intact in both *Dbh*^{-/-} mice and people with complete DBH deficiency (Garland et al., 2011; Lustberg et al., 2022; Thomas and Palmiter 1997). Therefore, our findings suggest that suppression of DBH produces aberrant responses of rodents to threats. Crucially, NE transmission is generally anxiogenic and mediates aversive responses to stress, whereas DA signaling is a key component of many appetitive/approach behaviors (Iordanova et al., 2021). Thus, we speculate that the loss of NE suppresses adaptive fearful/defensive behaviors, while the increased DA reverses the valence of predator odor from aversive to appetitive. Further experiments combining genetic and pharmacological manipulations of catecholamine signaling and circuits with real-time and conditioned place preference/aversion approaches will help test this idea.

4.5. Neuroanatomical substrates of NE and DA action

Results from the c-fos immunostaining experiment provide clues about where NE and DA released from noradrenergic neurons may drive

predator odor-induced behaviors (Fig. 7). *Dbh* $-/-$ mice had fewer c-fos $+$ neurons in a field of NET $+$ fibers in the ACC, dBNST, PAG, and LS, suggesting that NE is required in one or more of these brain regions for predator odor-induced defensive behaviors (i.e. burying, arousal). These regions were chosen for examination because they are activated by predator odor, are innervated by noradrenergic neurons, and/or are key nodes in defensive behavior circuits (Janitzky et al., 2015; Jhang et al., 2018; McGregor et al., 2004). Because the contribution of these regions and circuits to predator odor responses is vast and has been reviewed elsewhere (Dielenberg and McGregor 2001; Rosen et al., 2008; Takahashi 2014), we will only highlight a few relevant findings with particular relevance to our data here. For example, microinfusion of either β AR or α 1AR antagonists into the LS reduced defensive burying in the shock probe test, while intra-BNST application of the α 2AR agonist clonidine suppressed TMT-induced NE release and fear responses (Fendt et al., 2005). Furthermore, BNST inactivation not only suppresses aversive responses to cat urine in rats but increases duration of contact with the odor source, suggesting enhanced affiliative behavior similar to those observed in *Dbh* $-/-$ mice (Xu et al., 2012). The PAG is generally implicated in defensive behaviors and fear responses and is of particular interest here because we previously identified a noradrenergic LC-PAG circuit regulating arousal that may be engaged to maintain vigilance under threatening conditions (Porter-Stransky et al., 2019). The lack of NE in *Dbh* $-/-$ mice impairs this circuit and leads to reduced arousal, which could explain why the knockouts fall asleep in the presence of

bobcat urine. Interestingly, inactivation of the ACC increases predator odor-induced freezing, while activation has the opposite effect (Jhang et al., 2018). It is possible that the increased activation of the ACC in our control mice was permissive for active coping (defensive burying) at the expense of passive coping (freezing). We did not observe a high degree of freezing in *Dbh* $-/-$ mice despite the reduced c-fos response, suggesting that other brain regions responsible for grooming dominated in the knockouts.

Two regions, the MeA and LC, had an increased c-fos response in *Dbh* $-/-$ mice. The MeA receives direct and indirect input from the olfactory system, as well as DA neurons in the VTA. Dopaminergic signaling within the MeA mediates social reward and has even been implicated in human bonding (Atzil et al., 2017; Hu et al., 2021; Takahashi 2014). D1 receptor-expressing MeA neurons are activated by cat urine, and facilitating VTA dopaminergic transmission or activating D1 MeA GABAergic outputs to the BNST causes animals to display increased approach to a predator odor threat (McGregor et al., 2004; Miller et al., 2019; Vincenz et al., 2017).

These data are consistent with the idea that bobcat urine elicits paradoxical appetitive rather than aversive responses in *Dbh* $-/-$ mice because of ectopic DA release from LC afferents in the MeA. Because α 2AR inhibitory autoreceptors limit LC activation, it is likely that the increased c-fos response in the LC of *Dbh* $-/-$ mice is due to their lack of NE and α 2AR engagement (Paladini et al., 2007).




Fig. 7. Proposed noradrenergic circuit for defensive behavioral responses to predator odors. The PVN hypothalamus is activated by ancient odorant pathways, in turn exciting the LC by releasing CRH (black arrow). The LC projects to a constellation of brain regions implicated in defensive behavior, where it releases NE in DBH-sufficient animals (blue arrows). Loss of NE results in hypoactivity in PAG, ACC, LS, and dBNST (regions without circles). Interestingly, loss of NE did not affect neuronal activity in the PVN (green circle), suggesting this pathway remains intact. Further, NE deficiency resulted in hyperactivity in the LC and the MeA (magenta circles). Image created with Biorender.

4.6. Translational significance and implications for *T. gondii* infection

It is important to identify the neurobiological underpinnings of species-typical and maladaptive responses to threats because dysregulated behavioral responses to stressful stimuli represent many hallmark features of neuropsychiatric disorders, including posttraumatic stress disorder (PTSD), anxiety, and drug abuse (Dias et al., 2013). Indeed, drugs that impact catecholaminergic transmission are used to treat affective diseases, although the mechanisms of action are not fully understood.

The results of this study may also have implications for *Toxoplasma gondii* infection. As a parasite that reproduces in cats as part of its life cycle, *T. gondii* impairs the innate fear of cat odor in rodents, increasing the chance that the parasite reaches its intermediate (feline) and permanent (larger animals, including humans) host (Kannan et al., 2010; Vyas et al., 2007; Webster et al., 1994). Although the mechanisms by which *T. gondii* changes host behavior are not fully understood, recent studies showed that *T. gondii* infection in rodents profoundly decreases expression of DBH, reducing NE and elevating DA in the brain (Alsaady et al., 2019; Tedford et al., 2023). Our results provide additional evidence that the *T. gondii*-induced reduction of DBH may lead to suppression of defensive behaviors that would normally promote escape from a predator. Indeed, the apparent affiliation of *Dbh* $-/-$ mice to feline scents (i.e. sitting on domestic cat fur, nesting in bobcat urine-soaked cotton) may actually attract them to danger and would make them easy prey in the wild. We predict that at least some of the behavioral effects of *T. gondii* infection would be abolished or occluded in *Dbh* $-/-$ mice. Since NE has anti-inflammatory properties (Chalermphanupap et al., 2013; Feinstein et al., 2016), it is possible that *T. gondii*-induced inflammation would be greater in *Dbh* $-/-$ mice. Consistent with our intriguing finding of increased c-fos in the MeA of *Dbh* $-/-$ mice, reversal of *T. gondii*-induced epigenetic and gene expression signatures in this structure normalized defensive behavior and abolished the gain of attraction to cat odor (Hari Dass and Vyas 2014), further indicating a potential neuroanatomical substrate for the excessive DA in *Dbh* $-/-$ mice.

T. gondii-induced changes in behavioral responses to predator stress is not merely a rodent curiosity; approximately one-third of the human population carries a chronic *T. gondii* infection, and while most infected people are grossly asymptomatic, toxoplasmosis can have devastating effects on the fetal brain and is associated with greater risk for schizophrenia, OCD, ASD, suicide, and homicide in people with latent *T. gondii* infection (Contopoulos-Ioannidis et al., 2022, Cook et al., 2015; Nayeri et al., 2022; Nayeri et al., 2020; Torrey and Yolken 2003; Virus et al., 2021). Thus, further study of NE and DA circuitry underlying *T. gondii*-induced behavioral changes in rodents and of catecholaminergic drugs as therapies for disorders with stress-responsive repetitive behaviors are warranted.

CRediT authorship contribution statement

Joyce Liu: Writing – review & editing, Writing – original draft, Visualization, Validation, Methodology, Investigation, Formal analysis, Conceptualization. **Daniel J. Lustberg:** Writing – review & editing, Writing – original draft, Visualization, Validation, Methodology, Investigation, Formal analysis, Conceptualization. **Abigail Galvez:** Validation, Methodology, Investigation, Formal analysis. **L. Cameron Liles:** Resources. **Katharine E. McCann:** Writing – review & editing, Writing – original draft, Validation, Supervision, Methodology, Investigation, Formal analysis. **David Weinshenker:** Writing – review & editing, Writing – original draft, Visualization, Resources, Project administration, Methodology, Funding acquisition, Formal analysis, Conceptualization.

Declaration of competing interest

DW is co-inventor on a patent concerning the use of selective dopamine β -hydroxylase inhibitors for the treatment of cocaine dependence (US-2010-0105,748-A1; “Methods and Compositions for Treatment of Drug Addiction”). The other authors declare no conflicts of interest.

Acknowledgements

We thank Q. Eastman for helpful editing of the manuscript and Cordelia and Lila for donating fur. This work was supported by the National Institutes of Health (DA049257, AG061175, and AG079199 to DW; NS007480-20 to DL).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.yjnstr.2024.100612>.

References

- Aghajanian, G.K., VanderMaelen, C.P., 1982. Alpha 2-adrenoceptor-mediated hyperpolarization of locus coeruleus neurons: intracellular studies in vivo. *Science* 215, 1394–1396.
- Alsaady, I., Tedford, E., Alsaad, M., Bristow, G., Kohli, S., et al., 2019. Downregulation of the central noradrenergic system by *Toxoplasma gondii* infection. *Infect. Immun.* 87.
- Atzil, S., Touroutoglou, A., Rudy, T., Salcedo, S., Feldman, R., et al., 2017. Dopamine in the medial amygdala network mediates human bonding. *Proc Natl Acad Sci U S A* 114, 2361–2366.
- Beas, B.S., Wright, B.J., Skirzewski, M., Leng, Y., Hyun, J.H., et al., 2018. The locus coeruleus drives disinhibition in the midline thalamus via a dopaminergic mechanism. *Nat. Neurosci.* 21, 963–973.
- Berridge, K.C., Aldridge, J.W., 2000a. Super-stereotypy I: enhancement of a complex movement sequence by systemic dopamine D1 agonists. *Synapse* 37, 194–204.
- Berridge, K.C., Aldridge, J.W., 2000b. Super-stereotypy II: enhancement of a complex movement sequence by intraventricular dopamine D1 agonists. *Synapse* 37, 205–215.
- Bondi, C.O., Barrera, G., Lapiz, M.D., Bedard, T., Mahan, A., Morilak, D.A., 2007. Noradrenergic facilitation of shock-probe defensive burying in lateral septum of rats, and modulation by chronic treatment with desipramine. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 31, 482–495.
- Bourdelat-Parks, B.N., Anderson, G.M., Donaldson, Z.R., Weiss, J.M., Bonsall, R.W., et al., 2005. Effects of dopamine beta-hydroxylase genotype and disulfiram inhibition on catecholamine homeostasis in mice. *Psychopharmacology (Berl)* 183, 72–80.
- Cabib, S., Castellano, C., Cestari, V., Filibeck, U., Puglisi-Allegra, S., 1991. D1 and D2 receptor antagonists differently affect cocaine-induced locomotor hyperactivity in the mouse. *Psychopharmacology (Berl)* 105, 335–339.
- Carter, M.E., Yizhar, O., Chikahisa, S., Nguyen, H., Adamantidis, A., et al., 2010. Tuning arousal with optogenetic modulation of locus coeruleus neurons. *Nat. Neurosci.* 13, 1526–1533.
- Chalermphanupap, T., Kinkead, B., Hu, W.T., Kummer, M.P., Hammerschmidt, T., et al., 2013. Targeting norepinephrine in mild cognitive impairment and Alzheimer's disease. *Alzheimer's Res. Ther.* 5, 21.
- Chinen, C.C., Frussa-Filho, R., 1999. Conditioning to injection procedures and repeated testing increase SCH 23390-induced catalepsy in mice. *Neuropsychopharmacology* 21, 670–678.
- Cho, J.R., Treweek, J.B., Robinson, J.E., Xiao, C., Bremner, L.R., et al., 2017. Dorsal raphe dopamine neurons modulate arousal and promote wakefulness by salient stimuli. *Neuron* 94, 1205–12019 e8.
- Chowdhury, A., Luchetti, A., Fernandes, G., Filho, D.A., Kastellakis, G., et al., 2022. A locus coeruleus-dorsal CA1 dopaminergic circuit modulates memory linking. *Neuron* 110, 3374–33788 e8.
- Cilz, N.I., Kurada, L., Hu, B., Lei, S., 2014. Dopaminergic modulation of GABAergic transmission in the entorhinal cortex: concerted roles of $\alpha 1$ adrenoceptors, inward rectifier K(+), and T-type Ca(2)(+) channels. *Cerebr. Cortex* 24, 3195–3208.
- Contopoulos-Ioannidis, D.G., Gianniki, M., Ai-Nhi Truong, A., Montoya, J.G., 2022. Toxoplasmosis and schizophrenia: a systematic review and meta-analysis of prevalence and associations and future directions. *Psychiatr Res Clin Pract* 4, 48–60.
- Cook, T.B., Brenner, L.A., Cloninger, C.R., Langenberg, P., Igbide, A., et al., 2015. “Latent” infection with *Toxoplasma gondii*: association with trait aggression and impulsivity in healthy adults. *J. Psychiatr.* Res. 60, 87–94.
- Curtis, A.L., Leiser, S.C., Snyder, K., Valentino, R.J., 2012. Predator stress engages corticotropin-releasing factor and opioid systems to alter the operating mode of locus coeruleus norepinephrine neurons. *Neuropharmacology* 62, 1737–1745.
- Day, H.E., Masini, C.V., Campeau, S., 2004. The pattern of brain c-fos mRNA induced by a component of fox odor, 2,5-dihydro-2,4,5-trimethylthiazoline (TMT), in rats,

- suggests both systemic and processive stress characteristics. *Brain Res.* 1025, 139–151.
- De Boer, S.F., Koolhaas, J.M., 2003. Defensive burying in rodents: ethology, neurobiology and psychopharmacology. *Eur. J. Pharmacol.* 463, 145–161.
- De Monte, F.H., Canteras, N.S., Fernandes, D., Assreuy, J., Carobrez, A.P., 2008. New perspectives on beta-adrenergic mediation of innate and learned fear responses to predator odor. *J. Neurosci.* 28, 13296–13302.
- Devoto, P., Flore, G., Saba, P., Frau, R., Gessa, G.L., 2015. Selective inhibition of dopamine-beta-hydroxylase enhances dopamine release from noradrenergic terminals in the medial prefrontal cortex. *Brain Behav* 5, e00393.
- Dias, B.G., Banerjee, S.B., Goodman, J.V., Ressler, K.J., 2013. Towards new approaches to disorders of fear and anxiety. *Curr. Opin. Neurobiol.* 23, 346–352.
- Dielenberg, R.A., McGregor, I.S., 2001. Defensive behavior in rats towards predatory odors: a review. *Neurosci. Biobehav. Rev.* 25, 597–609.
- Dixit, P.V., Sahu, R., Mishra, D.K., 2020. Marble-burying behavior test as a murine model of compulsive-like behavior. *J. Pharmacol. Toxicol. Methods* 102, 106676.
- Drago, F., Contarino, A., Busa, L., 1999. The expression of neuropeptide-induced excessive grooming behavior in dopamine D1 and D2 receptor-deficient mice. *Eur. J. Pharmacol.* 365, 125–131.
- Dreher, J.K., Jackson, D.M., 1989. Role of D1 and D2 dopamine receptors in mediating locomotor activity elicited from the nucleus accumbens of rats. *Brain Res.* 487, 267–277.
- Eban-Rothschild, A., Rothschild, G., Giardino, W.J., Jones, J.R., de Lecea, L., 2016. VTA dopaminergic neurons regulate ethologically relevant sleep-wake behaviors. *Nat. Neurosci.* 19, 1356–1366.
- Feinstein, D.L., Kalinin, S., Braun, D., 2016. Causes, consequences, and cures for neuroinflammation mediated via the locus coeruleus: noradrenergic signaling system. *J. Neurochem.* 139 (Suppl. 2), 154–178.
- Fendt, M., Siegl, S., Steiniger-Brach, B., 2005. Noradrenaline transmission within the ventral bed nucleus of the stria terminalis is critical for fear behavior induced by trimethylthiazoline, a component of fox odor. *J. Neurosci.* 25, 5998–6004.
- Ferrero, D.M., Lemon, J.K., Fluegge, D., Pashkovski, S.L., Korzan, W.J., et al., 2011. Detection and avoidance of a carnivore odor by prey. *Proc Natl Acad Sci U S A* 108, 11235–11240.
- Fucich, E.A., Morilak, D.A., 2018. Shock-probe defensive burying test to measure active versus passive coping style in response to an aversive stimulus in rats. *Bio Protoc* 8.
- Garbe, C.M.K.E., Rawleigh, J.M., 1993. Novel odors evoke risk assessment and suppress appetitive behaviors in mice. *Aggress. Behav.* 19, 447–454.
- Garland, E.M., Raj, S.R., Peltier, A.C., Robertson, D., Biaggioni, I., 2011. A cross-sectional study contrasting olfactory function in autonomic disorders. *Neurology* 76, 456–460.
- Guild, A.L., Dunn, A.J., 1982. Dopamine involvement in ACTH-induced grooming behavior. *Pharmacol. Biochem. Behav.* 17, 31–36.
- Hari Dass, S.A., Vyas, A., 2014. Toxoplasma gondii infection reduces predator aversion in rats through epigenetic modulation in the host medial amygdala. *Mol. Ecol.* 23, 6114–6122.
- Harris, N.A., Isaac, A.T., Gunther, A., Merkel, K., Melchior, J., et al., 2018. Dorsal BNST alpha(2a)-adrenergic receptors produce HCN-dependent excitatory actions that initiate anxiogenic behaviors. *J. Neurosci.* 38, 8922–8942.
- Hayley, S., Borowski, T., Merali, Z., Anisman, H., 2001. Central monoamine activity in genetically distinct strains of mice following a psychogenic stressor: effects of predator exposure. *Brain Res.* 892, 293–300.
- Hu, R.K., Zuo, Y., Ly, T., Wang, J., Meera, P., et al., 2021. An amygdala-to-hypothalamus circuit for social reward. *Nat. Neurosci.* 24, 831–842.
- Hunsley, M.S., Palmiter, R.D., 2003. Norepinephrine-deficient mice exhibit normal sleep-wake states but have shorter sleep latency after mild stress and low doses of amphetamine. *Sleep* 26, 521–526.
- Hunsley, M.S., Palmiter, R.D., 2004. Altered sleep latency and arousal regulation in mice lacking norepinephrine. *Pharmacol. Biochem. Behav.* 78, 765–773.
- Hwa, L.S., Neira, S., Pina, M.M., Pati, D., Calloway, R., Kash, T.L., 2019. Predator odor increases avoidance and glutamatergic synaptic transmission in the prelimbic cortex via corticotropin-releasing factor receptor 1 signaling. *Neuropsychopharmacology* 44, 766–775.
- Iordanova, M.D., Yau, J.O., McDannald, M.A., Corbit, L.H., 2021. Neural substrates of appetitive and aversive prediction error. *Neurosci. Biobehav. Rev.* 123, 337–351.
- Janitzky, K., D'Hanis, W., Krober, A., Schwegler, H., 2015. TMT predator odor activated neural circuit in C57BL/6J mice indicates TMT-stress as a suitable model for uncontrollable intense stress. *Brain Res.* 1599, 1–8.
- Jhang, J., Lee, H., Kang, M.S., Lee, H.S., Park, H., Han, J.H., 2018. Anterior cingulate cortex and its input to the basolateral amygdala control innate fear response. *Nat. Commun.* 9, 2744.
- Kalueff, A.V., Tuohimaa, P., 2005. Mouse grooming microstructure is a reliable anxiety marker bidirectionally sensitive to GABAergic drugs. *Eur. J. Pharmacol.* 508, 147–153.
- Kannan, G., Moldovan, K., Xiao, J.C., Yolken, R.H., Jones-Brando, L., Pletnikov, M.V., 2010. Toxoplasma gondii strain-dependent effects on mouse behaviour. *Folia Parasitol (Praha)* 57, 151–155.
- Kedia, S., Chattarji, S., 2014. Marble burying as a test of the delayed anxiogenic effects of acute immobilisation stress in mice. *J. Neurosci. Methods* 233, 150–154.
- Kemble, E.D., Bolwahn, B.L., 1997. Immediate and long-term effects of novel odors on risk assessment in mice. *Physiol. Behav.* 61, 543–549.
- Kemble, E.D.G.C., Gordon, C., 1995. Effects of novel odors on intermale attack behavior in mice. *Aggress Behav* 21, 293–299.
- Kempadoo, K.A., Mosharov, E.V., Choi, S.J., Sulzer, D., Kandel, E.R., 2016. Dopamine release from the locus coeruleus to the dorsal hippocampus promotes spatial learning and memory. *Proc Natl Acad Sci U S A* 113, 14835–14840.
- Ketenci, S., Acet, N.G., Saridogan, G.E., Aydin, B., Cabadak, H., Goren, M.Z., 2020. The neurochemical effects of prazosin treatment on fear circuitry in a rat traumatic stress model. *Clin Psychopharmacol Neurosci* 18, 219–230.
- Krach, S., Paulus, F.M., Boddien, M., Kircher, T., 2010. The rewarding nature of social interactions. *Front. Behav. Neurosci.* 4, 22.
- Lewis, S., 2011. Autism: grooming mice to model autism. *Nat. Rev. Neurosci.* 12, 248–249.
- Londei, T., Valentini, A.M., Leone, V.G., 1998. Investigative burying by laboratory mice may involve non-functional, compulsive, behaviour. *Behav. Brain Res.* 94, 249–254.
- Lu, J., Zhou, T.C., Saper, C.B., 2006. Identification of wake-active dopaminergic neurons in the ventral periaqueductal gray matter. *J. Neurosci.* 26, 193–202.
- Lustberg, D., Iannitelli, A.F., Tillage, R.P., Pruitt, M., Liles, L.C., Weinschenker, D., 2020a. Central norepinephrine transmission is required for stress-induced repetitive behavior in two rodent models of obsessive-compulsive disorder. *Psychopharmacology (Berl)* 237, 1973–1987.
- Lustberg, D., Tillage, R.P., Bai, Y., Pruitt, M., Liles, L.C., Weinschenker, D., 2020b. Noradrenergic circuits in the forebrain control affective responses to novelty. *Psychopharmacology (Berl)* 237, 3337–3355.
- Lustberg, D.J., Liu, J.Q., Iannitelli, A.F., Vanderhoof, S.O., Liles, L.C., et al., 2022. Norepinephrine and dopamine contribute to distinct repetitive behaviors induced by novel odorant stress in male and female mice. *Horm. Behav.* 144, 105205.
- McCall, J.G., Al-Hasani, R., Siuda, E.R., Hong, D.Y., Norris, A.J., et al., 2015. CRH engagement of the locus coeruleus noradrenergic system mediates stress-induced anxiety. *Neuron* 87, 605–620.
- McGregor, I.S., Hargreaves, G.A., Apfelbach, R., Hunt, G.E., 2004. Neural correlates of cat odor-induced anxiety in rats: region-specific effects of the benzodiazepine midazolam. *J. Neurosci.* 24, 4134–4144.
- McGregor, I.S., Schrama, L., Ambermoon, P., Dielenberg, R.A., 2002. Not all 'predator odours' are equal: cat odour but not 2,4,5 trimethylthiazoline (TMT; fox odour) elicits specific defensive behaviours in rats. *Behav. Brain Res.* 129, 1–16.
- Miller, S.M., Marcotulli, D., Shen, A., Zweifel, L.S., 2019. Divergent medial amygdala projections regulate approach-avoidance conflict behavior. *Nat. Neurosci.* 22, 565–575.
- Mitchell, H.A., Bogenpohl, J.W., Liles, L.C., Epstein, M.P., Bozyczko-Coyne, D., et al., 2008. Behavioral responses of dopamine beta-hydroxylase knockout mice to modafinil suggest a dual noradrenergic-dopaminergic mechanism of action. *Pharmacol. Biochem. Behav.* 91, 217–222.
- Mitchell, H.A., Weinschenker, D., 2010. Good night and good luck: norepinephrine in sleep pharmacology. *Biochem. Pharmacol.* 79, 801–809.
- Nayeri, T., Sarvi, S., Daryani, A., 2022. Toxoplasmosis: targeting neurotransmitter systems in psychiatric disorders. *Metab. Brain Dis.* 37, 123–146.
- Nayeri, T., Sarvi, S., Moosazadeh, M., Hosseini-najad, Z., Sharif, M., et al., 2020. Relationship between toxoplasmosis and autism: a systematic review and meta-analysis. *Microb. Pathog.* 147, 104434.
- Nordstrom, E.J., Burton, F.H., 2002. A transgenic model of comorbid Tourette's syndrome and obsessive-compulsive disorder circuitry. *Mol. Psychiatr.* 7, 617–625, 524.
- Ohkubo, N., Aoto, M., Kon, K., Mitsuda, N., 2019. Lack of zinc finger protein 521 upregulates dopamine beta-hydroxylase expression in the mouse brain, leading to abnormal behavior. *Life Sci.* 231, 116559.
- Ozkan, M., Johnson, N.W., Sehirli, U.S., Woodhall, G.L., Stanford, I.M., 2017. Dopamine acting at D1-like, D2-like and alpha1-adrenergic receptors differentially modulates theta and gamma oscillatory activity in primary motor cortex. *PLoS One* 12, e0181633.
- Paladini, C.A., Beckstead, M.J., Weinschenker, D., 2007. Electrophysiological properties of catecholaminergic neurons in the norepinephrine-deficient mouse. *Neuroscience* 144, 1067–1074.
- Paladini, C.A., Fiorillo, C.D., Morikawa, H., Williams, J.T., 2001. Amphetamine selectively blocks inhibitory glutamate transmission in dopamine neurons. *Nat. Neurosci.* 4, 275–281.
- Petter, E.A., Fallon, I.P., Hughes, R.N., Watson, G.D.R., Meck, W.H., et al., 2023. Elucidating a locus coeruleus-dentate gyrus dopamine pathway for operant reinforcement. *Elife* 12.
- Pina, M.M., Cunningham, C.L., 2014. Effects of dopamine receptor antagonists on the acquisition of ethanol-induced conditioned place preference in mice. *Psychopharmacology (Berl)* 231, 459–468.
- Poe, G.R., Foote, S., Eschenko, O., Johansen, J.P., Bouret, S., et al., 2020. Locus coeruleus: a new look at the blue spot. *Nat. Rev. Neurosci.* 21, 644–659.
- Pond, H.L., Heller, A.T., Gural, B.M., McKissick, O.P., Wilkinson, M.K., Manzini, M.C., 2021. Digging behavior discrimination test to probe burrowing and exploratory digging in male and female mice. *J. Neurosci. Res.* 99, 2046–2058.
- Porter-Stransky, K.A., Centanni, S.W., Karne, S.L., Odil, L.M., Fekir, S., et al., 2019. Noradrenergic transmission at alpha1-adrenergic receptors in the ventral periaqueductal gray modulates arousal. *Biol. Psychiatr.* 85, 237–247.
- Rosen, J.B., Pagani, J.H., Rolla, K.L., Davis, C., 2008. Analysis of behavioral constraints and the neuroanatomy of fear to the predator odor trimethylthiazoline: a model for animal phobias. *Neurosci. Biobehav. Rev.* 32, 1267–1276.
- Sara, S.J., 2009. The locus coeruleus and noradrenergic modulation of cognition. *Nat. Rev. Neurosci.* 10, 211–223.
- Schank, J.R., Liles, L.C., Weinschenker, D., 2008. Norepinephrine signaling through beta-adrenergic receptors is critical for expression of cocaine-induced anxiety. *Biol. Psychiatr.* 63, 1007–1012.
- Schroeder, J.P., Cooper, D.A., Schank, J.R., Lyle, M.A., Gaval-Cruz, M., et al., 2010. Disulfiram attenuates drug-primed reinstatement of cocaine seeking via inhibition of dopamine beta-hydroxylase. *Neuropsychopharmacology* 35, 2440–2449.

- Schwarz, L.A., Luo, L., 2015. Organization of the locus coeruleus-norepinephrine system. *Curr. Biol.* 25, R1051–R1056.
- Sluyter, F., Korte, S.M., Bohus, B., Van Oortmerssen, G.A., 1996. Behavioral stress response of genetically selected aggressive and nonaggressive wild house mice in the shock-probe/defensive burying test. *Pharmacol. Biochem. Behav.* 54, 113–116.
- Smolinsky, A.N.B.C., LaPorte, J.L., Kaluff, A.V., 2009. Analysis of Grooming Behavior and its Utility in Studying Animal Stress, Anxiety, and Depression. Springer, pp. 21–36.
- Stanley, W.C., Li, B., Bonhaus, D.W., Johnson, L.G., Lee, K., et al., 1997. Catecholamine modulatory effects of nopicastat (RS-25560-197), a novel, potent and selective inhibitor of dopamine-beta-hydroxylase. *Br. J. Pharmacol.* 121, 1803–1809.
- Stone, E.A., Yan, L., Ahsan, M.R., Lehmann, M.L., Yeretsian, J., Quartermain, D., 2006. Role of CNS alpha1-adrenoceptor activity in central fos responses to novelty. *Synapse* 59, 299–307.
- Stone, E.A., Zhang, Y., Rosengarten, H., Yeretsian, J., Quartermain, D., 1999. Brain alpha 1-adrenergic neurotransmission is necessary for behavioral activation to environmental change in mice. *Neuroscience* 94, 1245–1252.
- Szot, P., Weinshenker, D., White, S.S., Robbins, C.A., Rust, N.C., et al., 1999. Norepinephrine-deficient mice have increased susceptibility to seizure-inducing stimuli. *J. Neurosci.* 19, 10985–10992.
- Takahashi, L.K., 2014. Olfactory systems and neural circuits that modulate predator odor fear. *Front. Behav. Neurosci.* 8, 72.
- Takeuchi, T., Duzkiewicz, A.J., Sonneborn, A., Spooner, P.A., Yamasaki, M., et al., 2016. Locus coeruleus and dopaminergic consolidation of everyday memory. *Nature* 537, 357–362.
- Tedford, E., Badya, N.B., Laing, C., Asaoka, N., Kaneko, S., et al., 2023. Infection-induced extracellular vesicles evoke neuronal transcriptional and epigenetic changes. *Sci. Rep.* 13, 6913.
- Thomas, S.A., Marck, B.T., Palmiter, R.D., Matsumoto, A.M., 1998. Restoration of norepinephrine and reversal of phenotypes in mice lacking dopamine beta-hydroxylase. *J. Neurochem.* 70, 2468–2476.
- Thomas, S.A., Matsumoto, A.M., Palmiter, R.D., 1995. Noradrenaline is essential for mouse fetal development. *Nature* 374, 643–646.
- Thomas, S.A., Palmiter, R.D., 1997. Impaired maternal behavior in mice lacking norepinephrine and epinephrine. *Cell* 91, 583–592.
- Tillag, R.P., Sciolino, N.R., Plummer, N.W., Lustberg, D., Liles, L.C., et al., 2020. Elimination of galanin synthesis in noradrenergic neurons reduces galanin in select brain areas and promotes active coping behaviors. *Brain Struct. Funct.* 225, 785–803.
- Timmermans, P.B., van Zwieten, P.A., 1982. Alpha 2 adrenoceptors: classification, localization, mechanisms, and targets for drugs. *J. Med. Chem.* 25, 1389–1401.
- Torrey, E.F., Yolken, R.H., 2003. *Toxoplasma gondii* and schizophrenia. *Emerg. Infect. Dis.* 9, 1375–1380.
- Treit, D., Pinel, J.P., Fibiger, H.C., 1981. Conditioned defensive burying: a new paradigm for the study of anxiolytic agents. *Pharmacol. Biochem. Behav.* 15, 619–626.
- Vincenz, D., Wernecke, K.E.A., Fendt, M., Goldschmidt, J., 2017. Habenula and interpeduncular nucleus differentially modulate predator odor-induced innate fear behavior in rats. *Behav. Brain Res.* 332, 164–171.
- Virus, M.A., Ehrhorn, E.G., Lui, L.M., Davis, P.H., 2021. Neurological and neurobehavioral disorders associated with *Toxoplasma gondii* infection in humans. *J. Parasitol Res* 2021, 6634807.
- Vyas, A., Kim, S.K., Giacomini, N., Boothroyd, J.C., Sapolsky, R.M., 2007. Behavioral changes induced by *Toxoplasma* infection of rodents are highly specific to aversion of cat odors. *Proc Natl Acad Sci U S A* 104, 6442–6447.
- Wagatsuma, A., Okuyama, T., Sun, C., Smith, L.M., Abe, K., Tonegawa, S., 2018. Locus coeruleus input to hippocampal CA3 drives single-trial learning of a novel context. *Proc Natl Acad Sci U S A* 115, E310–E316.
- Wang, M., Ramos, B.P., Paspalas, C.D., Shu, Y., Simen, A., et al., 2007. Alpha2A-adrenoceptors strengthen working memory networks by inhibiting cAMP-HCN channel signaling in prefrontal cortex. *Cell* 129, 397–410.
- Webster, J.P., Brunton, C.F., MacDonald, D.W., 1994. Effect of *Toxoplasma gondii* upon neophobic behaviour in wild brown rats, *Rattus norvegicus*. *Parasitology* 109 (Pt1), 37–43.
- Weinshenker, D., Miller, N.S., Blizinsky, K., Laughlin, M.L., Palmiter, R.D., 2002. Mice with chronic norepinephrine deficiency resemble amphetamine-sensitized animals. *Proc Natl Acad Sci U S A* 99, 13873–13877.
- Welch, J.M., Lu, J., Rodriguiz, R.M., Trotta, N.C., Peca, J., et al., 2007. Cortico-striatal synaptic defects and OCD-like behaviours in Sapap3-mutant mice. *Nature* 448, 894–900.
- Xu, H.Y., Liu, Y.J., Xu, M.Y., Zhang, Y.H., Zhang, J.X., Wu, Y.J., 2012. Inactivation of the bed nucleus of the stria terminalis suppresses the innate fear responses of rats induced by the odor of cat urine. *Neuroscience* 221, 21–27.
- Yamaguchi, H., Hopf, F.W., Li, S.B., de Lecea, L., 2018. In vivo cell type-specific CRISPR knockdown of dopamine beta hydroxylase reduces locus coeruleus evoked wakefulness. *Nat. Commun.* 9, 5211.
- Yan, T., Xu, M., Wu, B., Liao, Z., Liu, Z., et al., 2016. The effect of *Schisandra chinensis* extracts on depression by noradrenergic, dopaminergic, GABAergic and glutamatergic systems in the forced swim test in mice. *Food Funct.* 7, 2811–2819.