

## Invited Mini Review

## Regulation of polyubiquitin genes to meet cellular ubiquitin requirement

Seung-Woo Han, Byung-Kwon Jung &amp; Kwon-Yul Ryu\*

Department of Life Science, University of Seoul, Seoul 02504, Korea

Ubiquitin (Ub) is one of the proteins that are highly conserved from yeast to humans. It is an essential core unit of the well-defined post-translational modification, called ubiquitination, which is involved in a variety of biological processes. In metazoans, Ub is encoded by two monoubiquitin genes and two polyubiquitin genes, in which a single Ub is fused to a ribosomal protein or Ub coding units are arranged in tandem repeats. In mice, polyubiquitin genes (*Ubb* and *Ubc*) play a pivotal role to meet the requirement of cellular Ub pools during embryonic development. In addition, expression levels of polyubiquitin genes are increased to adapt to environmental stimuli such as oxidative, heat-shock, and proteotoxic stress. Several researchers have reported about the perturbation of Ub pools through genetic alteration or exogenous Ub delivery using diverse model systems. To study Ub pool changes in a physiologically relevant manner, changing Ub pools via the regulation of endogenous polyubiquitin gene expression has recently been introduced. Furthermore, to understand the regulation of polyubiquitin gene expression more precisely, *cis*-acting elements and *trans*-acting factors, which are regulatory components of polyubiquitin genes, have been analyzed. In this review, we discuss how the role of polyubiquitin genes has been studied during the past decade, especially focusing on their regulation. [BMB Reports 2021; 54(4): 189-195]

## INTRODUCTION

Ubiquitin (Ub) is a highly conserved small eukaryotic protein composed of 76 amino acids, which can be mono- or polyubiquitinated on the substrate by Ub-activating enzyme (E1), Ub-conjugating enzyme (E2), and Ub ligase (E3) (1, 2). Monoubiquitination is known to regulate histone modification, receptor

endocytosis, etc. (3). Polyubiquitination plays different roles depending on the type of Ub chains (4). It is involved in a variety of cellular processes, such as proteasomal degradation, stress response, signal transduction, and membrane protein trafficking (5-8). After exerting its role, Ub conjugated on the substrate can be removed by deubiquitinating enzymes (DUBs) and becomes free Ub (9). Under certain circumstances, Ub itself can also be degraded by the proteasome (10). This Ub recycling is important for maintaining Ub homeostasis inside cells.

In mice, Ub is encoded by four different genes: *Uba52* and *Uba80* (*Rps27a*), which are Ub-ribosomal fusion genes, and *Ubb* and *Ubc*, which are polyubiquitin genes (Fig. 1). Similarly, there are also four different genes, *UBA52*, *UBA80* (*RPS27A*), *UBB*, and *UBC*, in humans (11-13). Ub-ribosomal fusion proteins or Ub polymers translated from these genes are converted into Ub monomers by DUBs, which constitute free Ub pools (14). Polyubiquitin genes are upregulated under stress conditions such as oxidative, heat-shock, and proteotoxic stress, thereby increasing the levels of free Ub pools (Fig. 1) (15, 16). This is important in maintaining cell viability when the cellular stress response pathway is activated (17). If the free Ub level does not increase under stress conditions, for example, because of disruption of the polyubiquitin gene, cell viability can be reduced (18).


It is also important to control the levels of Ub under normal conditions as well as stress conditions. Under normal conditions, the expression of both polyubiquitin genes is tightly regulated, so that it is not upregulated or downregulated significantly, to maintain the basal expression levels. If the levels of Ub decrease under normal conditions, problems may occur in proliferation, differentiation, and even in development (19-22). It has also been reported that chronic Ub overexpression induces synaptic dysfunction in neurons (23, 24). Therefore, the regulation of polyubiquitin gene expression under normal and stress conditions is important, but the relevant molecular mechanism is not known well. Some known facts are that the intron region is important for the regulation of basal expression levels of *UBC*, and the promoter region is important for the upregulation of *UBC* under stress conditions (25-28). Recently, it has been reported that using the inducible CRISPR-activation system, a modified version of the CRISPR/Cas9 system, in which guide RNA was targeted to the intron region of *UBC*,

\*Corresponding author. Tel: +82-2-6490-2674; Fax: +82-2-6490-2664; E-mail: kyryu@uos.ac.kr

<https://doi.org/10.5483/BMBRep.2021.54.4.005>

Received 31 December 2020, Revised 25 January 2021,  
Accepted 3 February 2021

**Keywords:** CRISPR-activation system, Polyubiquitin gene, Stress response, Transcriptional regulation, Ubiquitin



**Fig. 1.** Expression of ubiquitin genes under normal and stress conditions. Ubiquitin (Ub) is encoded by four different genes: *Uba52*, *Uba80*, *Ubb*, and *Ubc*. Two monoubiquitin genes (*Uba52* and *Uba80*) and two polyubiquitin genes (*Ubb* and *Ubc*) are structured as shown. Under normal conditions, four ubiquitin genes are continuously transcribed, translated, and deubiquitinated to produce available free Ub. Under stress conditions, the expression of polyubiquitin genes is upregulated. As a result, cellular free Ub pool levels increase under stress conditions. For simplicity, only mouse ubiquitin genes are shown here.

upregulation of *UBC* could be achieved temporally under normal conditions (29).

Based on these previous studies, we suggest that the regulation of endogenous polyubiquitin gene expression at the transcriptional level is possible under both normal and stress conditions, which is essential to change the levels of Ub pools. In this review, we have mostly focused on the research trend of transcriptional regulation of polyubiquitin genes during the past decade.

### The importance of free Ub pool levels


The cellular Ub pools are composed of free Ub and Ub conjugates, in which Ub is conjugated to the substrate by E1-E3 enzymes. Free Ub pools are maintained from a Ub monomer separated from Ub conjugates by DUBs and also through *de novo* synthesis of Ub by mono- and polyubiquitin genes. Newly synthesized Ub may exist as fusion proteins or polymers, but is converted to free Ub by DUBs right after synthesis (30). Among the important roles of Ub, the most well-known role is to form a K48-linked Ub chain on the substrate, which serves as a signal to be targeted to the 26S proteasome for degradation under both normal and stress conditions (5, 31). Ub also plays an essential role in cell proliferation, differentiation, and stress response. There are many types of Ub chains, for example, K63-linked and K29-linked (10, 32, 33). Although they exert different functions in cells, Ub chains are all formed from free Ub. Therefore, it is important to maintain the homeostasis of available free Ub pools. Either decreased or increased free Ub levels can cause adverse effects on cellular function or survival. Under normal conditions, disruption of Ub homeostasis through knockout of the polyubiquitin gene or ectopic expression of Ub induces dysregulation of proteasomal degradation or UPS (ubiquitin-proteasome system), causing various problems in tissues or cells (18, 23, 24, 34).

In fact, in neural stem cells (NSCs) with free Ub levels reduced because of *Ubb* knockout or knockdown, the degradation of the Notch intracellular domain (NICD) is delayed, leading to the activation of Notch signaling and upregulating the expression of Notch target genes (35, 36). Activation of Notch signaling in the early stages of neuronal development suppresses neurogenesis and promotes premature gliogenesis, causing abnormal differentiation of NSCs, degeneration of defective neurons, and reactive gliosis (35). Reduced free Ub levels also affect the proliferation of NSCs, resulting in a decrease in their numbers (35). In general, Ub deficiency is thought to cause a decrease in cell proliferation by interfering with the degradation of cyclin and CDK inhibitor, which are targets of the proteasome (31).

Cell proliferation decreased in *Ubc* knockout mouse embryonic fibroblasts (MEFs) (37). *Ubc* knockout embryos are lethal as early as 12.5 days post coitum (dpc) because of defects in fetal liver development (37). *Ubc* knockout embryos show smaller livers than do wild-type (WT) controls, suggesting that normal fetal liver development has not occurred, because of the decrease in cell proliferation during embryonic development. In *Ubb* knockout mice, spermatogenesis is arrested in the early pachytene stage of meiosis I, which hampers maturation of germ cells and results in smaller testes than in the WT controls, leading to an infertility phenotype (38). On the other hand, overexpression of Ub can also have an adverse outcome. Investigation of the levels of UPS substrates in the hippocampus of transgenic mice with chronic Ub overexpression has shown that protein levels, but not mRNA levels, of the glutamate ionotropic receptor (GRIA1-4) decreased (24). In addition, analysis of ubiquitinated proteins revealed that ubiquitination of GRIA increased. These results suggest that excessive Ub expression can cause abnormally rapid degradation of proteins through UPS, which leads to dysfunction of neurons (24).

### Maintenance of basal expression levels of polyubiquitin genes

In metazoans, polyubiquitin genes consist of multiple Ub coding units, which are arranged in a head-to-tail tandem array. The polyubiquitin gene *UBC* or *Ubc* has 2 to 3 times more Ub coding units than does the other polyubiquitin gene, *UBB* or *Ubb*. They are composed of the promoter, the non-coding exon (exon 1/2), the intron, and the coding exon (exon 2/2) that encodes Ub (Fig. 2). The putative enhancer can also be found in the intron region, not only in the upstream of the promoter region. For precise understanding of the regulation of polyubiquitin gene expression, analysis of *cis*-acting regulatory elements, located in the intron and the promoter region, and *trans*-acting transcription factors have been carried out. Various transcription factor binding sites, such as NF- $\kappa$ B, Sp1, heat-shock factor 1 (HSF1), and AP-1, have been found in the regulatory region of *UBC*, including the TATA box (13, 39). The mechanism for regulating *UBC* expression may differ depending on the cell type (40). For example, in muscle cells, activation of the mitogen-activated protein kinase (MAPK) signaling pathway by treatment with dexamethasone (glucocorticoid) and Sp1 transcription factor binding are involved in the regulation of *UBC* basal expression (41-43). In addition, the intron region is involved in the regulation of *UBC* basal expression (25, 26). The basal expression levels of *UBC* significantly decreased upon removal of the intron region of *UBC* (25). The transcriptional activity of *UBC* decreased when the Yin Yang 1 (YY1)-binding sequence present in the intron was mutated (26). Therefore, YY1 transcription factor binding to




**Fig. 2.** Structures of human polyubiquitin genes. *UBB* and *UBC* share genetic structures that consist of a promoter, a noncoding exon, an intron, and a Ub-coding exon. *Trans*-acting factors of *UBB* and *UBC* bind *cis*-acting elements located in the putative enhancer as well as in the promoter region. The putative enhancer can be found in the regulatory region including the intron. As an example of *trans*-acting factors, Sp1 and YY1 transcription factors binding to the *UBC* intron region are shown. Promoter and putative enhancer locations are displayed based on the candidate *cis*-acting regulatory elements (cCRE) from the Encyclopedia of DNA Elements (ENCODE) database. TSS, transcription start site.

the intron region plays an important role in the regulation of *UBC* basal expression.

When Ub levels are lowered because of disruption of one polyubiquitin gene, compensatory expression of the other polyubiquitin gene occurs to rescue cellular Ub levels. However, because of different cell or tissue spectra of two polyubiquitin gene expressions, such compensation cannot fully rescue Ub levels (Fig. 3). When Ub levels were increased through Ub overexpression, both polyubiquitin gene expressions were downregulated in mice and humans (24, 44). Intriguingly, when Ub was overexpressed, *UBC* pre mRNA levels were rather increased over those in the control; however, the mRNA levels tended to decrease (44). These results suggest that Ub overexpression contributes to the downregulation of *UBC* expression by affecting the post-transcriptional processing such as splicing.

### Upregulation of polyubiquitin gene expression under stress conditions

Polyubiquitin genes are stress-inducible genes and are upregulated under various stress conditions to increase cellular Ub levels. This upregulation occurs in diverse species, such as yeast, chicken, mouse, and human (15-17). Under oxidative or proteotoxic stress, the proteasome activity is compromised, and many misfolded proteins are generated (45, 46). If these misfolded proteins are not removed by the proteasome in a timely manner, insoluble protein aggregates can be formed that may cause cytotoxicity (47-49). Therefore, when the stress response is induced, it is important to upregulate the



**Fig. 3.** Tissue expression patterns of human polyubiquitin genes. Based on the Genotype-Tissue Expression (GTEx) and Human Protein Atlas database, protein-coding transcripts per million (pTPM) values of *UBB* and *UBC* from 32 tissues are presented. Gene information and raw data are available at <https://www.proteinatlas.org/ENSG00000170315-UBB/tissue> and <https://www.proteinatlas.org/ENSG00000150991-UBC/tissue>.

expression of polyubiquitin genes to meet the required free Ub pool levels, so that the increasing amount of misfolded proteins can be degraded without delay (16, 46). In fact, in cells treated with the proteasome inhibitors or exposed to oxidative stressors, *UBC* expression has been upregulated (28).

When *Ubb*<sup>+/-GFP-puro</sup> and *Ubc*<sup>+/-GFP-puro</sup> heterozygous MEFs, which synthesize GFP by the promoters of endogenous *Ubb* and *Ubc*, are exposed to proteasome inhibitor (MG-132), which causes proteotoxic stress, and arsenite, which causes oxidative stress, the GFP signal increased in a time- and concentration-dependent manner (50). Therefore, the increase in free Ub pool levels under stress conditions is achieved by upregulation of polyubiquitin genes. When *Ubc* knockout MEFs were exposed to oxidative stress, Ub levels were not increased and the viability of cells was significantly reduced (18). Similarly, reduced viability was observed when a polyubiquitin gene *UBI4* was disrupted in yeast (17). These results suggest that the upregulation of polyubiquitin genes under stress conditions and the corresponding increase in Ub levels are essential for stress tolerance.

The stress-inducible polyubiquitin gene *UBC* has heat-shock elements in the promoter region, which play an important role in the upregulation of the polyubiquitin gene under stress conditions (51, 52). In the promoter region of *UBC*, HSF1 binding allows upregulation of *UBC* expression. Under HSF1 deficiency by treatment with *HSF1* shRNA, the upregulation of *UBC* expression was greatly reduced even under stress conditions (27, 28, 39). The expression of HSF1 was increased under oxidative stress and proteasome inhibition, not only under heat-shock stress (53, 54). Thus, HSF1 plays a role to upregulate *UBC* expression in response to various cellular stresses or environmental stimuli.

When MEFs are exposed to oxidative stress, MAPK and nuclear factor erythroid 2-related factor 2 (Nrf2)-Kelch-like ECH-associated protein 1 (Keap1) pathways are activated (55). Stabilized Nrf2 moves into the nucleus as a transcription factor and binds to an antioxidant response element (ARE) to upregulate the expression of antioxidant response genes to resist oxidative stress (56). In *Ubc* promoter, there is a region containing the consensus sequence of ARE, and when this region is deleted, the transcriptional activity decreases even in the presence of stabilized Nrf2 (18). In addition, when MEFs were exposed to arsenite, *Ubc* mRNA levels increased significantly, but much less in Nrf2 knockout MEFs. These results provide evidence that under oxidative stress, Nrf2 may also affect the upregulation of *Ubc*. However, this phenomenon has not been observed in other cells, such as HeLa or NIH3T3 cells (28).

### A novel strategy to induce changes in Ub pools

Free Ub pools are maintained through the regulation of ubiquitin gene expression and by Ub recycling. The expression levels of polyubiquitin genes are tightly regulated under normal conditions without increasing or decreasing

significantly, thereby maintaining the basal levels. However, their expression levels increase dramatically under stress conditions (16, 37, 51, 52). It was not clear whether the expression of endogenous polyubiquitin genes can also be increased under normal conditions. To answer this question, ectopic Ub overexpression has been used to increase Ub levels (37, 57, 58). Conversely, a method used to reduce Ub levels is a knockdown through shRNA or use of cells isolated from knockout mice (36-38, 59, 60). These methods may alter the Ub levels whatever are the expression levels of endogenous genes, although overexpressed Ub has been shown to downregulate the expression of polyubiquitin genes.

Recently, with the development of the CRISPR/Cas9 system, which is widely used as a genome engineering tool in biological research, it has been suggested that Ub levels could potentially be controlled through transcriptional regulation (61, 62). In the CRISPR/Cas9 system, an endonuclease Cas9 binds the target sequence by guide RNA and induces double-strand break (63-65). By introducing double knockout of two polyubiquitin genes using this system, compensatory expression was avoided and more efficient free Ub depletion was achieved (34). In fact, double knockout of *UBB* and *UBC* in HEK293T and HeLa cells showed a dramatic decrease in free Ub levels with reduced proliferation and proteasome activity.

dCas9, a modified protein from which the nuclease activity of Cas9 has been removed, can carry out guide RNA-mediated binding to the target sequence, but does not cause double-strand break. When the transcriptional activator is fused to dCas9, the activator can be recruited only to the promoter of a specific gene, targeted by the guide RNA (65, 66). The transcriptional activator mainly used in this CRISPR-activation system is VP64, a tetramer of the minimal activation domain of VP16. VP16 is a transcription factor of herpes simplex virus type 1 and activates transcription by inducing chromatin remodeling through interaction with other transcription factors (67). The activation system consisting of a simple combination of guide RNA and dCas9-VP64 targeting the *UBC* intron region did not induce *UBC* upregulation (29). These results suggest that *UBC*, with high basal expression levels, may already have a relaxed chromatin structure for transcriptional activation (66). Therefore, to upregulate *UBC*, a modified guide RNA-derived system using an RNA aptamer was introduced (66). *UBC* was successfully upregulated by means of this system, in which the activation domain of HSF1 and p65, the activation domain of NF- $\kappa$ B, were added through RNA aptamer that was part of the modified guide RNA. Thus, modified guide RNA targeting the intron region of *UBC*, dCas9-VP64, and p65-HSF1 were required for the upregulation of *UBC* expression. Because *UBB* was not affected, this modified guide RNA-derived system is quite specific to *UBC* (29). In addition, regulation using this system was inducible using doxycycline and completely reversible (68). These findings suggest that even under normal conditions, it is

possible to induce changes in the free Ub pools by upregulation of endogenous polyubiquitin genes (29).

### Concluding remarks

So far, we have mentioned that the upregulation of polyubiquitin gene expression under stress conditions is important to confer resistance against stress, otherwise cells may exhibit the reduced viability. Not only under stress conditions, regulation of their expression is also important under normal conditions. It is problematic if the state of decreased or increased Ub levels persists because of failure in balancing the regulation of their expression. There is a limitation in that the strategies used in the past studies to change Ub pool levels are irreversible. Under normal conditions, Ub pools do not seem to be regulated beyond the threshold levels. Furthermore, when Ub is overexpressed, polyubiquitin genes are down-regulated, and when one polyubiquitin gene is disrupted, the other polyubiquitin gene is upregulated as a compensatory expression. These changes seem to maintain the free Ub levels inside cells. However, the molecular mechanism that can explain the dependency on Ub levels for the regulation of their expression is not known yet. One hypothesis is that there is a sensor that can bind free Ub directly or indirectly, and this sensor regulates the transcription factor complex assembly in the promoter region of polyubiquitin genes when it is released from free Ub.

Recently, using the CRISPR/Cas9 system, double knockout of *UBB* and *UBC* has been shown to be a way to deplete free Ub pools more efficiently, because compensatory expression can be avoided. In addition, transcriptional regulation using a dCas9-fused effector may be a promising candidate to regulate Ub levels. Especially, the CRISPR-activation system can be advantageous for conferring oxidative stress resistance, and this system regulates endogenous gene expression, which is cell friendly and physiologically relevant. It has been reported that continuous overexpression of Ub can prepare cellular resistance against expected stress conditions (69). However, persistent Ub overexpression may have adverse effects, such as destroying the synaptic function or reducing muscle development (24, 70). Therefore, if polyubiquitin genes can be upregulated temporarily by means of the reversible CRISPR-activation system, this problem could be overcome; so this system is useful under stress conditions. Furthermore, the potential of this system in Ub research is that, if *UBB* can also be upregulated in a similar way, it can be used to upregulate both polyubiquitin genes, *UBB* and *UBC*, for maximal increase of Ub levels, or upregulate one of two polyubiquitin genes in a tissue-specific manner, for efficient increase of Ub levels.

During the past decade, although many studies have been conducted on the regulation of polyubiquitin gene expression, there are still areas that need to be explored. Considering the diverse functions of Ub, it is important to study the mechanism by which polyubiquitin genes are regulated. It seems that the development of various high-throughput sequencing techniques

provided the foundation for studying the mechanism of gene regulation. Understanding the regulatory mechanism of intracellular Ub levels should be beneficial for studying Ub pool dynamics. Furthermore, therapeutic strategies for diseases such as dementia and cancer can be developed using Ub as a target protein.

### ACKNOWLEDGEMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2020R1F1A1070847) to KYR.

### CONFLICTS OF INTEREST

The authors have no conflicting interests.

### REFERENCES

1. Hershko A and Ciechanover A (1998) The ubiquitin system. *Annu Rev Biochem* 67, 425-479
2. Hershko A, Heller H, Elias S and Ciechanover A (1983) Components of ubiquitin-protein ligase system. Resolution, affinity purification, and role in protein breakdown. *J Biol Chem* 258, 8206-8214
3. Hicke L (2001) Protein regulation by monoubiquitin. *Nat Rev Mol Cell Biol* 2, 195-201
4. Schwartz AL and Ciechanover A (1992) Ubiquitin-mediated protein modification and degradation. *Am J Respir Cell Mol Biol* 7, 463-468
5. Hochstrasser M (1996) Ubiquitin-dependent protein degradation. *Annu Rev Genet* 30, 405-439
6. Pickart CM and Fushman D (2004) Polyubiquitin chains: polymeric protein signals. *Curr Opin Chem Biol* 8, 610-616
7. Sun L and Chen ZJ (2004) The novel functions of ubiquitination in signaling. *Curr Opin Cell Biol* 16, 119-126
8. Akutsu M, Dikic I and Bremm A (2016) Ubiquitin chain diversity at a glance. *J Cell Sci* 129, 875-880
9. Komander D, Clague MJ and Urbe S (2009) Breaking the chains: structure and function of the deubiquitinases. *Nat Rev Mol Cell Biol* 10, 550-563
10. Park CW and Ryu KY (2014) Cellular ubiquitin pool dynamics and homeostasis. *BMB Rep* 47, 475-482
11. Baker RT and Board PG (1991) The human ubiquitin-52 amino acid fusion protein gene shares several structural features with mammalian ribosomal protein genes. *Nucleic Acids Res* 19, 1035-1040
12. Redman KL and Rechsteiner M (1989) Identification of the long ubiquitin extension as ribosomal protein S27a. *Nature* 338, 438-440
13. Wiborg O, Pedersen MS, Wind A, Berglund LE, Marcker KA and Vuust J (1985) The human ubiquitin multigene family: some genes contain multiple directly repeated ubiquitin coding sequences. *EMBO J* 4, 755-759
14. Amerik AY and Hochstrasser M (2004) Mechanism and function of deubiquitinating enzymes. *Biochim Biophys*

- Acta 1695, 189-207
15. Bond U and Schlesinger MJ (1986) The chicken ubiquitin gene contains a heat shock promoter and expresses an unstable mRNA in heat-shocked cells. *Mol Cell Biol* 6, 4602-4610
  16. Fornace AJ Jr, Alamo I Jr, Hollander MC and Lamoreaux E (1989) Ubiquitin mRNA is a major stress-induced transcript in mammalian cells. *Nucleic Acids Res* 17, 1215-1230
  17. Finley D, Ozkaynak E and Varshavsky A (1987) The yeast polyubiquitin gene is essential for resistance to high temperatures, starvation, and other stresses. *Cell* 48, 1035-1046
  18. Kim MN, Choi J, Ryu HW and Ryu KY (2015) Disruption of polyubiquitin gene Ubc leads to attenuated resistance against arsenite-induced toxicity in mouse embryonic fibroblasts. *Biochim Biophys Acta* 1853, 996-1009
  19. Buckley SM, Aranda-Orgilles B, Strikoudis A et al (2012) Regulation of pluripotency and cellular reprogramming by the ubiquitin-proteasome system. *Cell Stem Cell* 11, 783-798
  20. Guo W, Shang F, Liu Q, Urim L, Zhang M and Taylor A (2006) Ubiquitin-proteasome pathway function is required for lens cell proliferation and differentiation. *Invest Ophthalmol Vis Sci* 47, 2569-2575
  21. Xu P, Duong DM, Seyfried NT et al (2009) Quantitative proteomics reveals the function of unconventional ubiquitin chains in proteasomal degradation. *Cell* 137, 133-145
  22. Chen PC, Bhattacharyya BJ, Hanna J et al (2011) Ubiquitin homeostasis is critical for synaptic development and function. *J Neurosci* 31, 17505-17513
  23. Hallengren J, Chen PC and Wilson SM (2013) Neuronal ubiquitin homeostasis. *Cell Biochem Biophys* 67, 67-73
  24. Vaden JH, Tian T, Golf S, McLean JW, Wilson JA and Wilson SM (2019) Chronic over-expression of ubiquitin impairs learning, reduces synaptic plasticity, and enhances GRIA receptor turnover in mice. *J Neurochem* 148, 386-399
  25. Bianchi M, Crinelli R, Giacomini E, Carloni E and Magnani M (2009) A potent enhancer element in the 5'-UTR intron is crucial for transcriptional regulation of the human ubiquitin C gene. *Gene* 448, 88-101
  26. Bianchi M, Crinelli R, Giacomini E, Carloni E, Radici L and Magnani M (2013) Yin Yang 1 intronic binding sequences and splicing elicit intron-mediated enhancement of ubiquitin C gene expression. *PLoS One* 8, e65932
  27. Crinelli R, Bianchi M, Radici L, Carloni E, Giacomini E and Magnani M (2015) Molecular dissection of the human ubiquitin C promoter reveals heat shock element architectures with activating and repressive functions. *PLoS One* 10, e0136882
  28. Bianchi M, Crinelli R, Arbore V and Magnani M (2018) Induction of ubiquitin C (UBC) gene transcription is mediated by HSF1: role of proteotoxic and oxidative stress. *FEBS Open Bio* 8, 1471-1485
  29. Han SW, Jung BK, Park SH and Ryu KY (2019) Reversible regulation of polyubiquitin gene UBC via modified inducible CRISPR/Cas9 system. *Int J Mol Sci* 20, 3168
  30. Ryu HW and Ryu KY (2015) Ubiquitin homeostasis: from neural stem cell differentiation to neuronal development. *Neural Regen Res* 10, 1209-1210
  31. Hershko A (2005) The ubiquitin system for protein degradation and some of its roles in the control of the cell division cycle. *Cell Death Differ* 12, 1191-1197
  32. Komander D and Rape M (2012) The ubiquitin code. *Annu Rev Biochem* 81, 203-229
  33. Komander D (2009) The emerging complexity of protein ubiquitination. *Biochem Soc Trans* 37, 937-953
  34. Park CW, Bae JS and Ryu KY (2020) Simultaneous disruption of both polyubiquitin genes affects proteasome function and decreases cellular proliferation. *Cell Biochem Biophys* 78, 321-329
  35. Ryu HW, Park CW and Ryu KY (2014) Disruption of polyubiquitin gene Ubb causes dysregulation of neural stem cell differentiation with premature gliogenesis. *Sci Rep* 4, 7026
  36. Jung BK, Park CW and Ryu KY (2018) Temporal down-regulation of the polyubiquitin gene Ubb affects neuronal differentiation, but not maturation, in cells cultured in vitro. *Sci Rep* 8, 2629
  37. Ryu KY, Maehr R, Gilchrist CA et al (2007) The mouse polyubiquitin gene UbC is essential for fetal liver development, cell-cycle progression and stress tolerance. *EMBO J* 26, 2693-2706
  38. Ryu KY, Sinnar SA, Reinholdt LG et al (2008) The mouse polyubiquitin gene Ubb is essential for meiotic progression. *Mol Cell Biol* 28, 1136-1146
  39. Nenoï M, Mita K, Ichimura S et al (1996) Heterogeneous structure of the polyubiquitin gene UbC of HeLa S3 cells. *Gene* 175, 179-185
  40. Marinovic AC, Zheng B, Mitch WE and Price SR (2007) Tissue-specific regulation of ubiquitin (UbC) transcription by glucocorticoids: in vivo and in vitro analyses. *Am J Physiol Renal Physiol* 292, F660-666
  41. Bell LM, Leong ML, Kim B et al (2000) Hyperosmotic stress stimulates promoter activity and regulates cellular utilization of the serum- and glucocorticoid-inducible protein kinase (Sgk) by a p38 MAPK-dependent pathway. *J Biol Chem* 275, 25262-25272
  42. Chupreta S, Du M, Todisco A and Merchant JL (2000) EGF stimulates gastrin promoter through activation of Sp1 kinase activity. *Am J Physiol Cell Physiol* 278, C697-708
  43. Marinovic AC, Zheng B, Mitch WE and Price SR (2002) Ubiquitin (UbC) expression in muscle cells is increased by glucocorticoids through a mechanism involving Sp1 and MEK1. *J Biol Chem* 277, 16673-16681
  44. Bianchi M, Crinelli R, Giacomini E et al (2019) A negative feedback mechanism links UBC gene expression to ubiquitin levels by affecting RNA splicing rather than transcription. *Sci Rep* 9, 18556
  45. Caballero M, Liton PB, Epstein DL and Gonzalez P (2003) Proteasome inhibition by chronic oxidative stress in human trabecular meshwork cells. *Biochem Biophys Res Commun* 308, 346-352
  46. Shang F and Taylor A (2011) Ubiquitin-proteasome pathway and cellular responses to oxidative stress. *Free Radic Biol Med* 51, 5-16
  47. Taylor A and Davies KJ (1987) Protein oxidation and loss of protease activity may lead to cataract formation in the aged lens. *Free Radic Biol Med* 3, 371-377

48. Julien JP (2001) Amyotrophic lateral sclerosis. unfolding the toxicity of the misfolded. *Cell* 104, 581-591
49. Shang F, Nowell TR Jr and Taylor A (2001) Removal of oxidatively damaged proteins from lens cells by the ubiquitin-proteasome pathway. *Exp Eye Res* 73, 229-238
50. Ryu HW and Ryu KY (2011) Quantification of oxidative stress in live mouse embryonic fibroblasts by monitoring the responses of polyubiquitin genes. *Biochem Biophys Res Commun* 404, 470-475
51. Vihervaara A, Sergelius C, Vasara J et al (2013) Transcriptional response to stress in the dynamic chromatin environment of cycling and mitotic cells. *Proc Natl Acad Sci U S A* 110, E3388-3397
52. Bianchi M, Giacomini E, Crinelli R, Radici L, Carloni E and Magnani M (2015) Dynamic transcription of ubiquitin genes under basal and stressful conditions and new insights into the multiple UBC transcript variants. *Gene* 573, 100-109
53. Pirkkala L, Alastalo TP, Zuo X, Benjamin IJ and Sistonen L (2000) Disruption of heat shock factor 1 reveals an essential role in the ubiquitin proteolytic pathway. *Mol Cell Biol* 20, 2670-2675
54. Morimoto RI (1998) Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes Dev* 12, 3788-3796
55. Taguchi K, Motohashi H and Yamamoto M (2011) Molecular mechanisms of the Keap1-Nrf2 pathway in stress response and cancer evolution. *Genes Cells* 16, 123-140
56. Kaspar JW, Niture SK and Jaiswal AK (2009) Nrf2:Keap1 signaling in oxidative stress. *Free Radic Biol Med* 47, 1304-1309
57. Peloponese JM Jr, Iha H, Yedavalli VR et al (2004) Ubiquitination of human T-cell leukemia virus type 1 tax modulates its activity. *J Virol* 78, 11686-11695
58. Mosesson Y, Shtiegman K, Katz M et al (2003) Endocytosis of receptor tyrosine kinases is driven by mono-ubiquitylation, not polyubiquitylation. *J Biol Chem* 278, 21323-21326
59. Oh C, Park S, Lee EK and Yoo YJ (2013) Downregulation of ubiquitin level via knockdown of polyubiquitin gene Ubb as potential cancer therapeutic intervention. *Sci Rep* 3, 2623
60. Scarpa ES, Tasini F, Crinelli R, Ceccarini C, Magnani M and Bianchi M (2020) The ubiquitin gene expression pattern and sensitivity to UBB and UBC knockdown differentiate primary 23132/87 and metastatic MKN45 gastric cancer cells. *Int J Mol Sci* 21, 5435
61. Perez-Pinera P, Kocak DD, Vockley CM et al (2013) RNA-guided gene activation by CRISPR-Cas9-based transcription factors. *Nat Methods* 10, 973-976
62. Qi LS, Larson MH, Gilbert LA et al (2013) Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell* 152, 1173-1183
63. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA and Zhang F (2013) Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* 8, 2281-2308
64. Wang T, Wei JJ, Sabatini DM and Lander ES (2014) Genetic screens in human cells using the CRISPR-Cas9 system. *Science* 343, 80-84
65. Shalem O, Sanjana NE and Zhang F (2015) High-throughput functional genomics using CRISPR-Cas9. *Nat Rev Genet* 16, 299-311
66. Konermann S, Brigham MD, Trevino AE et al (2015) Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* 517, 583-588
67. Hirai H, Tani T and Kikyo N (2010) Structure and functions of powerful transactivators: VP16, MyoD and FoxA. *Int J Dev Biol* 54, 1589-1596
68. Gossen M, Freundlieb S, Bender G, Muller G, Hillen W and Bujard H (1995) Transcriptional activation by tetracyclines in mammalian cells. *Science* 268, 1766-1769
69. Chen Y and Piper PW (1995) Consequences of the overexpression of ubiquitin in yeast: elevated tolerances of osmotic stress, ethanol and canavanine, yet reduced tolerances of cadmium, arsenite and paromomycin. *Biochim Biophys Acta* 1268, 59-64
70. Vaden JH, Watson JA, Howard AD, Chen PC, Wilson JA and Wilson SM (2015) Distinct effects of ubiquitin overexpression on NMJ structure and motor performance in mice expressing catalytically inactive USP14. *Front Mol Neurosci* 8, 11