

Research article

Open Access

## Cyclin A1 promoter hypermethylation in human papillomavirus-associated cervical cancer

Nakarin Kitkumthorn<sup>1,2</sup>, Pattamawadee Yanatatsanajit<sup>1</sup>, Sorapop Kiatpongsan<sup>3</sup>, Chureerat Phokaew<sup>1</sup>, Surang Triratanachat<sup>3</sup>, Prasert Trivijitsilp<sup>3</sup>, Wichai Termrungruanglert<sup>3</sup>, Damrong Tresukosol<sup>3</sup>, Somchai Niruthisard<sup>3</sup> and Apiwat Mutirangura\*<sup>1,4</sup>

Address: <sup>1</sup>Molecular Biology and Genetics of Cancer Development Research Unit, Faculty of Medicine, Chulalongkorn University, Rama IV., Bangkok 10330, Thailand, <sup>2</sup>Department of Microbiology, Faculty of Medicine, Chulalongkorn University, Rama IV., Bangkok 10330, Thailand, <sup>3</sup>Department of Obstetrics and Gynecology, Faculty of Medicine, Chulalongkorn University, Rama IV., Bangkok 10330, Thailand and <sup>4</sup>Department of Anatomy, Faculty of Medicine, Chulalongkorn University, Rama IV., Bangkok 10330, Thailand

Email: Nakarin Kitkumthorn - Art\_khuan@hotmail.com; Pattamawadee Yanatatsanajit - Nuchjoe@hotmail.com; Sorapop Kiatpongsan - Ksorapop@yahoo.com; Chureerat Phokaew - Hoshleys@hotmail.com; Surang Triratanachat - Surangtrirattanachat@hotmail.com; Prasert Trivijitsilp - Tprasert@chula.ac.th; Wichai Termrungruanglert - Wichaiterm@hotmail.com; Damrong Tresukosol - Dtresukosol@hotmail.com; Somchai Niruthisard - Ndsomchai@hotmail.com; Apiwat Mutirangura\* - mapiwat@chula.ac.th

\* Corresponding author

Published: 08 March 2006

Received: 23 November 2005

BMC Cancer 2006, 6:55 doi:10.1186/1471-2407-6-55

Accepted: 08 March 2006

This article is available from: <http://www.biomedcentral.com/1471-2407/6/55>

© 2006 Kitkumthorn et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** The aim of this study was to evaluate epigenetic status of *cyclin A1* in human papillomavirus-associated cervical cancer. Y. Tokumaru et al., *Cancer Res* **64**, 5982-7 (Sep 1, 2004) demonstrated in head and neck squamous-cell cancer an inverse correlation between *cyclin A1* promoter hypermethylation and *TP53* mutation. Human papillomavirus-associated cervical cancer, however, is deprived of *TP53* function by a different mechanism. Therefore, it was of interest to investigate the epigenetic alterations during multistep cervical cancer development.

**Methods:** In this study, we performed duplex methylation-specific PCR and reverse transcriptase PCR on several cervical cancer cell lines and microdissected cervical cancers. Furthermore, the incidence of *cyclin A1* methylation was studied in 43 samples of white blood cells, 25 normal cervixes, and 24, 5 and 30 human papillomavirus-associated premalignant, microinvasive and invasive cervical lesions, respectively.

**Results:** We demonstrated *cyclin A1* methylation to be commonly found in cervical cancer, both in vitro and in vivo, with its physiological role being to decrease gene expression. More important, this study demonstrated that not only is *cyclin A1* promoter hypermethylation strikingly common in cervical cancer, but is also specific to the invasive phenotype in comparison with other histopathological stages during multistep carcinogenesis. None of the normal cells and low-grade squamous intraepithelial lesions exhibited methylation. In contrast, 36.6%, 60% and 93.3% of high-grade squamous intraepithelial lesions, microinvasive and invasive cancers, respectively, showed methylation.

**Conclusion:** This methylation study indicated that *cyclin A1* is a potential tumor marker for early diagnosis of invasive cervical cancer.

## Background

Cervical cancer (CC) is an important health problem and is a leading cause of cancer mortality worldwide in women. [1] When exposed to and infected by one of the high-risk human papillomaviruses (HPV), vulnerable cervical epithelium may enter a complex multistep process and develop an invasive carcinoma. [2-4] The spectrum of histologic alterations during the intricate processes of multistep carcinogenesis can be classified as premalignant lesions, including low-grade and high-grade squamous intraepithelial lesions (SILs), and malignant invasive cervical cancers. [5] Despite its strong association with CC, HPV infection alone is not sufficient for the cervical epithelium to fully develop an invasive cervical cancer. Persistent HPV infection contributes to the development of SILs, with viral oncoproteins facilitating the dysregulation of cellular proliferation and the apoptotic process. However, additional accumulation of mutations, as well as epigenetic alterations in the crucial oncogenes and tumor suppressor genes, is required before these premalignant lesions fully transform into invasive cancers. [6]

The aim of this study was to evaluate DNA methylation status of *cyclin A1* (*CCNA1*) in HPV-associated CC. *CCNA1*, a second A-type cyclin, has been shown to be essential for entry into metaphase of male meiosis I [7,8] Consistent with this function, *CCNA1* is highly expressed in testis and hematopoietic progenitor cells, but is present at low levels in most other tissues. [9] No phenotype other than male infertility has been reported in mice lacking *CCNA1*. [10] Surprisingly, several lines of evidence suggest that *CCNA1* may be a potential epithelial tumor suppressor gene. First, the expression of *CCNA1* has been demonstrated to be downregulated in several cancers, such as nasopharyngeal carcinoma and head and neck squamous-cell cancer (HNSCC). [11-13] Second, *CCNA1* plays an important role in DNA double-strand break repair following radiation damage by activation of the non-homologous end-joining process that confers DNA stability. [14] Finally, the promoter, similar to several key tumor suppressor genes, is frequently hypermethylated in colon cancer and HNSCC. [13,15]

Expression of *CCNA1* has been shown to be correlated with the activation of *TP53*. In a HNSCC model, there is an inverse relationship between *CCNA1* promoter methylation and *TP53* mutation status in HNSCC tissues. [13] Similar to HNSCC, the majority of CC is of squamous cell origin and its molecular carcinogenesis strongly correlates with impaired *TP53* function. [16-18] However, unlike HNSCC, the functional loss of *TP53* in CC is not ascribed to gene mutation, but is processed by viral and host protein-protein interaction. CC is strongly associated with infection by high-risk HPV types and its oncoprotein E6 has the ability to associate with and neutralize the func-

tion of *TP53*. [17,18] E6 binds to *TP53* and catalyzes multi-ubiquitination and degradation of *TP53*. Consequently, the majority of CC cells have a wild-type *TP53*, but the protein levels are decreased. Therefore, in comparison with HNSCC, it was of interest to determine if *CCNA1* is methylated in HPV-associated squamous cell CC.

## Methods

### Cell lines and tissue samples

SiHa and two HeLa CC cell lines from different sources were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum. All three cells were purchased from ATCC. SiHa, HeLa (S), and HeLa (K) were grown and maintained in laboratories of Dr. Ponglikitmongkol M, Mahidol University, Dr. Gutkind JS, NIH, USA and Dr. Ruxrungthum K, Chulalongkorn University, respectively.

With approval of ethical committee, faculty of medicine, Chulalongkorn university, normal cervical tissues, cancer tissues and blood samples were obtained and prepared as previously described. [19,20] Cervical tissues were obtained by punch biopsy of lesions under direct visualization or under colposcopic examination. Specimens were divided in two. The first sample was submitted to routine histological examination, and the second was reserved for DNA isolation. Blood samples were obtained by venipuncture from CC patients and healthy blood donors. All HPV-positive premalignant lesions were exfoliated cells, selected from routine cytological screening. In brief, cervical cells were collected with a cervical sampler (Digene Corporation, Gaithersburg, MD, USA) using the cervical cytobrush technique, and were divided into three parts. The first was reserved for routine cytological diagnosis. The second was tested for the presence of high-risk HPV (types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, and 68) DNA by Hybrid Capture 2 (Digene Corporation, Gaithersburg, MD, USA). [21] In cases of positive high-risk HPV and complete histological tissue evaluation, the third part was subjected to *CCNA1* methylation analysis. DNA extraction was performed using Tris/SDS and proteinase K at 50°C overnight, followed by phenol/chloroform extraction and ethanol precipitation.

Cervical biopsy specimens and Papanicolaou smears were examined and reviewed by at least two gynecologic pathologists to ensure good quality control of the final pathology results. All CCs contained 20–95% malignant cells. The histological diagnoses distinguished among normal epithelium, low-grade SILs, high-grade SILs, microinvasive and invasive cancer. In case of invasive cancer, only those samples classified as squamous-cell lesions were used for further analysis.

**Table 1: Oligonucleotide sequences and conditions for PCR analyses**

Primer	Sequence	Amplicon size (bp)	Annealing temperature (°C)
CCNA1metF	TTTCGAGGATTTTCGCGTCGT	46	53
CCNA1metR	CTCCTAAAAACCCCTAACTCGA		
CCNA1unmetF	TTAGTGTGGGTAGGGTGT	67	53
CCNA1unmetR	CCCTAACTCAAAAAACAACACA		
CCNA1cloningF	TGGGTAGGGCGCTCGTAGTT	196	55
CCNA1cloningR	GCCCCGACCTAAAAAAA		
CCNA1cDNAF	ATTCATTAAGTGAAATTGTGC	170	47
CCNA1cDNAR	CTTCCATTTCAGAACTTATTG		
GAPDH	GTGGGCAAGGTATCCCTG	460	52
GAPDHR	GATTCAGTGTGGTGGGGGAC		

Additional six OTC-embedded frozen CCs and five normal cervixes, obtained from hysterectomy specimens, were microdissected as previously described.<sup>22</sup> Histologically normal epithelium, connective tissue and malignant cells were subjected to *CCNA1* methylation and expression studies.

#### HPV detection and typing

HPV L1, E6 gene amplification and dot blot hybridization were performed as previously described [19,22,23]. Briefly, each L1 amplification reaction contained the L1 degenerate primers MY11 and MY09. The E6 reactions contained WD72, WD66, WD154, WD67 and WD76. Both reactions were used to amplify genomic DNA during 40 PCR cycles. To analyze the amplicons for the presence of high-risk HPV, we applied dot blot hybridization using the HPV type-specific oligo probes, WD170, WD132, RR1, RR2, WD103, WD165, WD, consensus L1, MY12/13, WD126, WD128, MY16, WD133/134, MY14 and WD174. The membranes were subjected to analysis by a phosphorimager. Results for L1 and E6 dot blots were scored independently. Duplicate filters were prepared for all specimens.

#### Sodium bisulfite modification and duplex methylation-specific PCR (MSP)

The DNA samples were subjected to bisulfite treatment. [24,25] Briefly, 2 µg of genomic DNA was denatured with NaOH (final concentration 0.2 M). Subsequently, 10 mM hydroquinone and 3 M sodium bisulfite were added and incubated at 50°C for 16 h. The modified DNA was then purified using Wizard DNA purification resin (Promega, Madison, WI, USA) followed by ethanol precipitation. Duplex MSPs were performed to identify the *CCNA1* methylation status of all samples. The duplex PCR mixtures contained 10× PCR buffer (Qiagen, Chuo-ku, Tokyo), deoxynucleotide triphosphates (0.2 mM), primers CCNA1metF, CCNA1metR, CCNA1unmetF and CCNA1unmetR (final concentration 0.4 µM each per reaction) (Table 1), 1 U of HotStarTaq (Qiagen, Chuo-ku,

Tokyo) and bisulfited DNA (80 ng). The amplification reaction was carried out for 30 cycles in a 2400 Perkin Elmer thermal cycler. Then 10-µl aliquots of the PCR products were stained with cyber green, run on an 8% non-denaturing polyacrylamide gel. The band intensity was visualized and measured by using a phosphorimager.

#### RNA preparation and analysis

Expression of *CCNA1* in the CC cell lines was examined by RT-PCR. Total RNA was extracted using the TRIZOL reagent (Invitrogen, Singapore) according to the manufacturer's specifications and 5 µg of each sample was subjected to cDNA synthesis using MMLV reverse transcriptase (Fermentas, Hanover, MD, USA). PCR mixtures contained 10× PCR buffer, 0.2 mM dNTPs, 0.4 µM each of primers CCNA1cDNAF and CNA1cDNAR, 1 U of HotStartaq and 80 ng cDNA. *GAPDH* served as the internal control (Table 1). Aliquots of 10 µl of the PCR products were subjected to electrophoresis on a 2% agarose gel stained with ethidium bromide on preparation, and were visualized by a UV trans-illuminator.


#### Bisulfite genome sequence analysis

Some *CCNA1* methylation-positive CCs were selected for sequence analysis. The bisulfited DNAs were amplified using CCNA1cloningF and CCNA1cloningR (Table 1). The amplified fragments were cloned using the PGemT easy vector and sequenced.

#### Results

The aim of this study was to determine if the *CCNA1* promoter is methylated in CC and to elucidate how the epigenetic alteration occurs during multistep CC development. The experiments conducted comprised of: first, establishment of *CCNA1* MSP; second, identification of the methylation status and correlation with expression in CC cell lines, normal cervix and CC; and finally, investigation of the frequency of methylation in normal tissues, high-risk HPV-associated low SILs, high SILs, microinvasive and invasive squamous cell CC.





**Figure 2**  
 Intra- and inter-assay variation of the duplex MSP. (A) Duplex MSP of a mixture of *CCNA1* complete and non-methylated CC cell lines, SiHa and HeLa(S), respectively. M, DNA size marker; Neg, negative; 0, 25, 50, 75, 100 Met (%) represent the proportion of SiHa DNA in the mixture, varied from 0 to 100%, respectively. The upper and lower bands are non-methylated and methylated bands, respectively, indicated by labeled arrows. (B) Graphical comparison between measured *CCNA1* methylation, percentage intensity of methylation amplicon (x-axis), and actual methylation, the proportion of SiHa DNA (y-axis). The bar height indicates the mean and error bars, T, represent standard deviation (SD) across experiments.

***CCNA1* methylation incidence during multistep cervical carcinogenesis**


Cervical intraepithelial neoplasia provides a crucial model to study the multistep process of carcinogenesis. Therefore, we evaluated the frequency of *CCNA1* methylation in several cervical epithelial tissues with a distinctive degree of malignant transformation, normal cervix, CIN, microinvasive and CC, respectively. We selected 43, 25 and 30 cases of white blood cells (WBC), normal cervical biopsies and invasive CCs, respectively (Table 2). Among these samples, 13 WBC samples and 6 normal cervical samples, located at least 3 cm from the tumor margin and showing the absence of HPV DNA, originated from CC patients. For all cases, when a methylated amplicon was visible and the methylation percentage measured exceeded 5%, the test was deemed positive. All selected CCs were squamous and positive for HPV. Of the cases, 24 harbored HPV type 16, 4 had HPV type 18 and 2 cases displayed unclassifiable HPV types. Interestingly, a high frequency of methylation was exclusively present in CCs, i.e., 28 cases or 93.3% (Fig. 4A,B and Table 2). To reveal multistep carcinogenesis, we included 24 cases of SILs and 5

microinvasive cancers from exfoliated cervical cells. All cases were positive for oncogenic HPV, analyzed by Hybrid Capture 2. Whereas 60% and 36.6% of the microinvasive cancers and high SILs, respectively, demonstrated *CCNA1* methylation, none of the HPV-associated low SILs exhibited these epigenetic changes (Fig. 4B and Table 2).

**Discussion**

This study demonstrated that: (i) *CCNA1* promoter hypermethylation in HPV-associated squamous cell CC is unusually common; (ii) it is specific to CC; and (iii) the methylation is more common in invasive phenotypes compared to other histopathological stages during multistep carcinogenesis. This finding identifies both the interesting biology of CC and a potential clinical application of *CCNA1* methylation as an additional molecular marker for the early diagnosis of invasive CC.

Annual cytology screening has dramatically increased the effectiveness of early CC detection. Nonetheless, additional tests will help to improve the sensitivity and specificity of a single Papanicolaou smear for histological



**Figure 3**  
*CCNA1* methylation and expression in microdissected cervical tissues. (A) Duplex MSP and *CCNA1* PCR; E and CNT are epithelium and connective tissue cells from normal cervix; N and T are adjacent histological normal and cancer cervical epithelium from CC, respectively. Arrows indicate non-methylated, methylated, *CCNA1* cDNA and *GAPDH* cDNA, respectively. (B) Bisulfite sequencing at the *CCNA1* promoter, with circles denoting the methylation status of each selected clone. Black and white circles are methylated CG dinucleotides, and non-methylated CpG dinucleotides and TG dinucleotides, respectively.

analysis. Recently, testing for oncogenic HPVs has been introduced to aid in the triage of women with atypical squamous cells of undetermined significance (ASCUS). [28] However, because the majority of patients with HPV-associated lesions do not progress to invasive cancer, several studies have attempted to add a panel of tumor suppressor gene methylations to improve the effectiveness of molecular cytological diagnosis. [29,30] Since the frequency of *CCNA1* methylation is high and specific to invasive CC, this gene should be a good candidate to increase the coverage rate for early cancer detection.

In HNSCC, *CCNA1* promoter hypermethylation is inversely related to *TP53* mutation. [13] Nonetheless, the frequency of *CCNA1* promoter hypermethylation in CC is high, whereas the function of *TP53* in CC is usually impaired as a consequence of protein degradation induced by binding of the viral E6 protein. [18] This observation may be due to either differences in tissue types or pathophysiological outcomes of *TP53* between mutations and diminution of the protein function subsequent to E6 binding. We prefer the latter hypothesis, since *TP53* and *CCNA1* have been shown to augment each


**Table 2: CCNA1 methylation and clinico-pathological correlation**

Histological characteristics	Total number of cases	CCNA1 promoter hypermethylation	
		Absent	Present
WBC	43	43	0
Normal cervix	25	25	0
Low-grade SIL	13	13	0
High-grade SIL	11	7	4
Microinvasive cancer	5	2	3
Squamous cell CC	30	2	28
FIGO stage I-IIA	6	0	6
FIGO stage IIB-IV	24	2	22
Grade 1, keratinized type	9	0	9
Grade 2, non-keratinized type	21	2	19

FIGO, International Federation of Gynecology and Obstetrics.

other's expression. [13,14] Consequently, the CCNA1 protein could help to increase physiologic TP53 to counter the function of E6, except for cases of TP53 mutation. In other words, alterations of both CCNA1 and TP53 in HNSCC will be redundant. In contrast, in CC, a decrease in CCNA1 protein should prevent the increment of TP53 that would have compensated for the protein destruction by E6.

Multistep process analysis revealed that CCNA1 methylation is remarkably specific for cervical carcinogenesis. The biological function of CCNA1 is to activate DNA breakage repair by mechanisms depending on CDK2 activity and Ku proteins. [14] It is interesting to hypothesize why the genomic instability, triggered by impairment of the CCNA1 function, is crucial as an early event in CC development. Perhaps the rate of spontaneous mutations in cervical epithelial cells is too low to accumulate sufficient malignancy-transformation-dependent oncogene and



**Figure 4**

Schematic representation of methylation-specific PCR in CC. (A) PCR analysis of CC: M, DNA size marker; Neg, water; N and T, matched normal cervixes and tumors, respectively. (B) Bar graph demonstrating the frequency of DNA methylation. Numbers on the y-axis are the percentage of positive methylation cases. Sample types are on the x-axis. WBC, normal cervix, Low-grade SIL, High-grade SIL, microinvasive cancer and squamous cell CCs number are 43, 25, 13, 11, 5, and 30, respectively. The methylation frequencies of each tissue type are represented by the height of each rectangular bar.

tumor suppressor gene mutations if the cells possess fully functional CCNA1. Therefore, the frequency of invasive CC devoid of CCNA1 methylation is limited.

## Conclusion

This study demonstrates the strong association between CCNA1 promoter hypermethylation and invasive HPV-associated CC indicates that this gene could serve as an effective molecular marker. Moreover, our finding, in comparison with previous reports, [13,14] also suggests that there is a possible molecular link between oncogenic HPVs, TP53 and CCNA1 promoter hypermethylation.

## Abbreviations

CC: cervical cancer, CCNA1: cyclin A1, SILs: squamous intraepithelial lesions, HPV: *Human papillomavirus*, WBC: white blood cell

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

NK: Perform all experiments, data analysis and write the article. PY: set up duplex MSP experiment, CP: collecting and HPV analysis of CIN, SK, ST, PT, WT, DT and SN: collect clinical samples and data, AM: Hypothesize, design and analyze the experiments and write the article

## Acknowledgements

We would like to thank the entire staff of the Department of Obstetrics and Gynecology for their assistance in collecting and providing the tissue samples, Drs. Virote Sriuranpong and Gutkind JS, Kiat Ruxrunghum, and Mathurose Ponglikitmongkol for the CC cell lines, HeLa(S), HeLa(K) and SiHa, respectively, peripheral nerve research unit for assisting in microdissection and Dr. Virote Sriuranpong, Ms Petra Hirsch, Mrs. Nita Suvarnses-takorn and Asia Science Editing for language editing the manuscript. This work has been supported by the National Center for Biotechnology and Genetic Engineering (Thailand), The Royal Golden Jubilee Ph.D. program, the Thailand Research Funds and the Molecular Biology and Genetics of Cancer Development research unit, Chulalongkorn University.

## References

- Pisani P, Parkin DM, Bray F, Ferlay J: **Estimates of the worldwide mortality from 25 cancers in 1990.** *Int J Cancer* 1999, **83(1)**:18-29.
- Durst M, Gritz D, Schneider A, zur Hausen H: **Human papillomavirus type 16 (HPV 16) gene expression and DNA replication in cervical neoplasia: analysis by in situ hybridization.** *Virology* 1992, **189(1)**:132-140.
- Bosch FX, Manos MM, Munoz N, Sherman M, Jansen AM, Peto J, Schiffman MH, Moreno V, Kurman R, Shah KV: **Prevalence of human papillomavirus in cervical cancer: a worldwide perspective. International biological study on cervical cancer (IBSCC) Study Group.** *J Natl Cancer Inst* 1995, **87(11)**:796-802.
- zur Hausen H: **Papillomaviruses and cancer: from basic studies to clinical application.** *Nat Rev Cancer* 2002, **2(5)**:342-350.
- The 1988 Bethesda System for reporting cervical/vaginal cytological diagnoses. National Cancer Institute Workshop.** *Jama* 1989, **262(7)**:931-934.
- Ferencyz A, Franco E: **Persistent human papillomavirus infection and cervical neoplasia.** *Lancet Oncol* 2002, **3(1)**:11-16.
- Sweeney C, Murphy M, Kubelka M, Ravnik SE, Hawkins CF, Wolgemuth DJ, Carrington M: **A distinct cyclin A is expressed in germ cells in the mouse.** *Development* 1996, **122(1)**:53-64.
- Liu D, Matzuk MM, Sung WK, Guo Q, Wang P, Wolgemuth DJ: **Cyclin A1 is required for meiosis in the male mouse.** *Nat Genet* 1998, **20(4)**:377-380.
- Yang R, Morosetti R, Koeffler HP: **Characterization of a second human cyclin A that is highly expressed in testis and in several leukemic cell lines.** *Cancer Res* 1997, **57(5)**:913-920.
- van der Meer T, Chan WY, Palazon LS, Nieduszynski C, Murphy M, Sobczak-Thepot J, Carrington M, Colledge WH: **Cyclin A1 protein shows haplo-insufficiency for normal fertility in male mice.** *Reproduction* 2004, **127(4)**:503-511.
- Maxwell SA, Davis GE: **Differential gene expression in p53-mediated apoptosis-resistant vs. apoptosis-sensitive tumor cell lines.** *Proc Natl Acad Sci U S A* 2000, **97(24)**:13009-13014.
- Sriuranpong V, Mutirangura A, Gillespie JW, Patel V, Amornphimoltham P, Molinolo AA, Kerekhanjanarong V, Supanakorn S, Supiyaphun P, Rangdaeng S, Voravud N, Gutkind JS: **Global gene expression profile of nasopharyngeal carcinoma by laser capture microdissection and complementary DNA microarrays.** *Clin Cancer Res* 2004, **10(15)**:4944-4958.
- Tokumar Y, Yamashita K, Osada M, Nomoto S, Sun DI, Xiao Y, Hoque MO, Westra WH, Califano JA, Sidransky D: **Inverse correlation between cyclin A1 hypermethylation and p53 mutation in head and neck cancer identified by reversal of epigenetic silencing.** *Cancer Res* 2004, **64(17)**:5982-5987.
- Muller-Tidow C, Ji P, Diederichs S, Potratz J, Baumer N, Kohler G, Cauvet T, Choudary C, van der Meer T, Chan WY, Nieduszynski C, Colledge WH, Carrington M, Koeffler HP, Restle A, Wiesmuller L, Sobczak-Thepot J, Berdel WE, Serve H: **The cyclin A1-CDK2 complex regulates DNA double-strand break repair.** *Mol Cell Biol* 2004, **24(20)**:8917-8928.
- Xu XL, Yu J, Zhang HY, Sun MH, Gu J, Du X, Shi DR, Wang P, Yang ZH, Zhu JD: **Methylation profile of the promoter CpG islands of 31 genes that may contribute to colorectal carcinogenesis.** *World J Gastroenterol* 2004, **10(23)**:3441-3454.
- Somers KD, Merrick MA, Lopez ME, Incognito LS, Schechter GL, Casey G: **Frequent p53 mutations in head and neck cancer.** *Cancer Res* 1992, **52(21)**:5997-6000.
- Tommasino M, Accardi R, Caldeira S, Dong W, Malanchi I, Smet A, Zehbe I: **The role of TP53 in Cervical carcinogenesis.** *Hum Mutat* 2003, **21(3)**:307-312.
- Thomas M, Pim D, Banks L: **The role of the E6-p53 interaction in the molecular pathogenesis of HPV.** *Oncogene* 1999, **18(53)**:7690-7700.
- Mutirangura A, Sriuranpong V, Termrungruanglert W, Tresukosol D, Lertsaguansinchai P, Voravud N, Niruthisard S: **Telomerase activity and human papillomavirus in malignant, premalignant and benign cervical lesions.** *Br J Cancer* 1998, **78(7)**:933-939.
- Pornthanakasem W, Shotelersuk K, Termrungruanglert W, Voravud N, Niruthisard S, Mutirangura A: **Human papillomavirus DNA in plasma of patients with cervical cancer.** *BMC Cancer* 2001, **1(1)**:2.
- Hubbard RA: **Human papillomavirus testing methods.** *Arch Pathol Lab Med* 2003, **127(8)**:940-945.
- Resnick RM, Cornelissen MT, Wright DK, Eichinger GH, Fox HS, ter Schegget J, Manos MM: **Detection and typing of human papillomavirus in archival cervical cancer specimens by DNA amplification with consensus primers.** *J Natl Cancer Inst* 1990, **82(18)**:1477-1484.
- Bauer HM, Ting Y, Greer CE, Chambers JC, Tashiro CJ, Chimera J, Reingold A, Manos MM: **Genital human papillomavirus infection in female university students as determined by a PCR-based method.** *Jama* 1991, **265(4)**:472-477.
- Herman JG, Graff JR, Myohanen S, Nelkin BD, Baylin SB: **Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands.** *Proc Natl Acad Sci U S A* 1996, **93(18)**:9821-9826.
- Chalitchagorn K, Shuangshoti S, Hourpai N, Kongruttanachok N, Tangkijvanich P, Thong-ngam D, Voravud N, Sriuranpong V, Mutirangura A: **Distinctive pattern of LINE-1 methylation level in normal tissues and the association with carcinogenesis.** *Oncogene* 2004, **23(54)**:8841-8846.
- Muller-Tidow C, Bornemann C, Diederichs S, Westermann A, Klumpen S, Zuo P, Wang W, Berdel WE, Serve H: **Analyses of the**



- genomic methylation status of the human cyclin A1 promoter by a novel real-time PCR-based methodology. *FEBS Lett* 2001, **490(1-2)**:75-78.
27. Muller C, Readhead C, Diederichs S, Idos G, Yang R, Tidow N, Serve H, Berdel WE, Koefler HP: **Methylation of the cyclin A1 promoter correlates with gene silencing in somatic cell lines, while tissue-specific expression of cyclin A1 is methylation independent.** *Mol Cell Biol* 2000, **20(9)**:3316-3329.
  28. Levi AW, Kelly DP, Rosenthal DL, Ronnett BM: **Atypical squamous cells of undetermined significance in liquid-based cytologic specimens: results of reflex human papillomavirus testing and histologic follow-up in routine practice with comparison of interpretive and probabilistic reporting methods.** *Cancer* 2003, **99(4)**:191-197.
  29. Widschwendter A, Gatringer C, Ivarsson L, Fiegl H, Schneitter A, Ramoni A, Muller HM, Wiedemair A, Jerabek S, Muller-Holzner E, Goebel G, Marth C, Widschwendter M: **Analysis of aberrant DNA methylation and human papillomavirus DNA in cervicovaginal specimens to detect invasive cervical cancer and its precursors.** *Clin Cancer Res* 2004, **10(10)**:3396-3400.
  30. Feng Q, Balasubramanian A, Hawes SE, Toure P, Sow PS, Dem A, Dembele B, Critchlow CW, Xi L, Lu H, McIntosh MW, Young AM, Kiviat NB: **Detection of hypermethylated genes in women with and without cervical neoplasia.** *J Natl Cancer Inst* 2005, **97(4)**:273-282.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/6/55/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

