

Screening for body dysmorphic disorder among patients pursuing cosmetic surgeries in Saudi Arabia

Hatan Mortada, MBBS, Hadeel Seraj, MBBS, Amal Bokhari, MD, FRCS.

ABSTRACT

الأهداف: اضطراب التشوه الجسمي هو اضطراب وسواسي (نفسي) منتشر بين المرضى الذين يبحثون عن عمليات تجميلية متكررة. ومع ذلك، فإن فحص المرض ليس ممارسة منتشرة. أجريت هذه الدراسة لتحديد مدى انتشار مرض اضطراب التشوه الجسمي بين مراجعي عيادات جراحات التجميل وجراحة تجميل العيون.

المنهجية: هذه الدراسة المقطعية تم إجراؤها عن طريق مقابلات شخصية مع مراجعي عيادات الجراحة التجميلية وجراحة تجميل العيون في مستشفى جامعة الملك عبد العزيز بجدة بين مارس 2019 ومارس 2020. تم التحقق من صحة استبيان اضطراب تشوه الجسم (BDDQ). تم منح الموافقة الأخلاقية من قبل لجنة أخلاقيات البحث. تم إجراء جميع الدراسات التحليلية باستخدام برنامج SPSS الإصدار 24.

النتائج: حجم العينة المشاركة = 344، بمتوسط عمر 39.66 ± 13.76 سنة. ومن بينهم 296 (86%) من النساء و 298 (86.6%) من السعوديين. كان انتشار الفحص الإيجابي لمرض اضطراب التشوه الجسمي 19.2%. أكثر العمليات المطلوبة من قبل المرضى كانت شد البطن (21.2%) وإزالة الآفات الجلدية (21.2%). وجد أن التدخين مرتبط بشكل كبير بـ BDD مع 21.2% من المدخنين الذين يعانون منه ($P > 0.010$).

الخلاصة: لم يتم التعرف على انتشار مرض اضطراب التشوه الجسمي بين المرضى الذين يتابعون الجراحة التجميلية وجراحة تجميل العيون. خمس المرضى الذين يطلبون إجراءات تجميلية هم حالات محتملة من مرض اضطراب التشوه الجسمي وتتطلب تقييمًا وعلاجًا نفسيًا وليس التدخل الجراحي. نوصي بتنفيذ بروتوكولات الفحص لتحديد الحالات المحتملة قبل التدخل الجراحي.

Objectives: To determine the prevalence of positive screening of body dysmorphic disorder (BDD) among patients seeking cosmetic surgeries in plastic surgery and oculoplastic surgery clinics.

Methods: The survey of this cross-sectional study was self-administrated and distributed among adults pursuing cosmetic surgeries in plastic surgery and oculoplastic surgery clinics at King Abdulaziz University Hospital, Jeddah, Saudi Arabia, between March 2019 and March 2020. The BDD questionnaire was validated, and a highly sensitive and specific tool was used to identify patients with BDD.

Ethical approval was granted by the Research Ethics Committee. All analytic studies were performed using IBM SPSS, version 24.

Results: A total of 344 patients participated in this study with a mean age of 39.66 ± 13.76 years. Of these, 296 (86%) were women and 298 (86.6%) were Saudi. The prevalence of positive screening for BDD was 19.2%. The most commonly requested procedures were abdominoplasty (21.2%) and skin lesion removal (21.2%). Smoking was found to be significantly associated with BDD with 21.2% of smokers having it ($p < 0.010$).

Conclusion: Body dysmorphic disorder was unrecognized among patients pursuing cosmetic surgeries. One-fifth of patients requesting cosmetic procedures are potential cases of BDD requiring psychiatric evaluation and treatment. We recommend implementing screening protocols to identify cases before surgical plans.

Keywords: body dysmorphic disorder, plastic surgery, oculoplastic surgery, prevalence, screening

*Saudi Med J 2020; Vol. 41 (10): 1111-1120
doi: 10.15537/smj.2020.10.25380*

From the Department of Plastic Surgery & Burn Unit (Mortada), King Saud Medical City, Riyadh; and from the Department of Ophthalmology (Seraj, Bokhari), Faculty of Medicine, King Abdulaziz University, Jeddah, Kingdom of Saudi Arabia.

Received 13th May 2020 Accepted 29th August 2020.

*Address correspondence and reprint request to: Dr. Hatan Mortada, Department of Plastic Surgery & Burn Unit, King Saud Medical City, Riyadh, Kingdom of Saudi Arabia. E-mail: Hatanmortada@gmail.com
ORCID ID: <https://orcid.org/0000-0003-1283-4136>*

Body dysmorphic disorder (BDD), also recognized as dysmorphophobia, is a DSM-V somatoform condition characterized by concerns about an imagined defect in appearance causing clinically major distress or functional damage that may extend to suicide.¹⁻⁴ Patients tend to exhibit repetitive skin picking, mirror checking, and grooming behavior. While different body parts can be involved, BDD mostly involves parts related to appearance, such as the face.² Moreover, most individuals have minor dissatisfactions related to personal image, which is not considered to be serious unless this dissatisfaction consumes most of the individual's thoughts and actions that might progress to severe disorders such as self-harm, anxiety, depression, and undergoing unneeded plastic surgery procedures.⁵ Furthermore, almost 76% of BDD patients visit a plastic surgeon to undergo cosmetic surgery, and more than half (60%) of patients with BDD undergo unneeded cosmetic surgery due to apparent imperfections.^{6,7} Most plastic surgeons (84%) reported that they had previously operated on a query BDD patient, and a vast majority (82%) of patients with BDD had poor results.^{8,9} Multiple previous studies have shown that patients with BDD usually have negative outcomes as well as dissatisfactions after the procedure.^{7,10} Consequently, it is vital to correctly identify BDD patients in any cosmetic surgery-related practices because the appropriate management for these individuals involves psychiatric evaluation rather than plastic surgery.^{11,12} The overall prevalence of patients with BDD among the general population is about 2.4%.¹³ However, among cosmetic-related practices, the prevalence of BDD is substantially higher, with a prevalence of 13% among patients pursuing plastic surgery clinics, 14% among patients in cosmetic dermatology, 6.9% in an oculoplastic surgery setting, and 10% in the maxillofacial clinic.¹⁴⁻¹⁷ The aforementioned prevalence of BDD in patients differ dramatically across practices, likely due to different patient populations and unpredictable screening approaches.¹⁰⁻¹³ Moreover, due to the uncommon and unusual screening of patients with BDD, there is usually an underdetection of patients with BDD.^{6,8,18} Body dysmorphic disorder patients who undergo cosmetic surgery usually consume the surgeon's effort and time with multiple requests for multiple procedures and consultations.

Moreover, patients with BDD are also more prone to consider lawsuits against their surgeons or even become physically violent.¹⁸⁻²⁰ These major concerns emphasize the importance of accurately diagnosing patients with BDD attending cosmetic management. Cross-sectional studies are required to offer data on the prevalence of BDD in plastic surgery and oculoplastic surgery clinics. A better understanding of this condition in patients seeking facial and oculoplastic surgeries can decrease the economic burden on health care systems and help redirect patients with BDD to seek the care they need, particularly if it is psychiatric doctor rather than plastic surgeon. As psychological intervention by mental health professionals is an important treatment approach, it mainly focuses on altering negative thoughts and behavior through specific supported strategies. Different screening tools were identified in the literature for screening of BDD, including the Dysmorphic Concern Questionnaire (DCQ). This questionnaire contains 7 items that examine physical appearance concerns. It has been validated with good internal consistency (Cronbach's alpha: 0.88).²¹ However, the sensitivity (72%) and specificity (90.7%) with a cut-off value of ≥ 14 (35) were much lower than the screening tool used in this study (Body Dysmorphic Disorder Questionnaire [BDDQ]), which has a sensitivity of 100% and specificity of 89-93%.²² Another available screening tool for BDD is the Body Dysmorphic Symptom Scale (BDSS); this screening tool is a self-reported 10-item questionnaire, focusing on patient physical appearance concerns. However, there is no specific accepted cut-off point for positive BDD screening in this questionnaire. Furthermore, it lacks validity.²³ For these reasons, this was not the screening tool used in this study. Another available screening tool is the BDDQ-Dermatology Version (BDDQ-DV). However, as the name implies, has most commonly been used in dermatology clinics. Little is known about such a disorder in patients from Jeddah. This study aim at identifying the prevalence of positive screening for BDD among patients pursuing cosmetic surgeries in plastic surgery and oculoplastic surgery clinics, and the association between BDD and the desired cosmetic surgical procedure at King Abdulaziz University Hospital (KAUH), Jeddah, Saudi Arabia.

Methods. After obtaining ethical approval from the ethics committee, this cross-sectional study was distributed randomly between March 2019 and March 2020 among adult patients pursuing cosmetic surgery in plastic surgery and oculoplastic surgery practice at KAUH. All patients aged 18 years or older who

Disclosure. Authors have no conflict of interests, and the work was not supported or funded by any drug company.

arrived for a new consultation to pursue cosmetic surgery in plastic surgery and oculoplastic surgery clinics were included in the screening survey during the pre-consultation process. Participants with medical indications for surgery for example visually significant ptosis and recurrent infected chalazions were excluded from the study. The main reason for undergoing surgery in our participants was for cosmetic purposes.

Individuals who agreed to fill out the questionnaire were offered a self-administered questionnaire. Randomly selected medical students volunteered as data collectors, informed patients about the study, and asked for the consent. Patients completed the questionnaire in the waiting room. After completing the survey, patients proceeded to their respective clinic appointments. The total sample size was calculated as 376 while keeping the margin of error as 5%, confidence level as 95%, population size as 20,000, and response distribution as 50% by using raosoft sample size calculator.²⁴ Participants completed a validated, anonymous, self-administered questionnaire.

Study variables. The questionnaire of this study had 3 main categories including: i) Sociodemographic (age, nationality, marital status, educational level, monthly household income, smoking status, comorbidities); ii) Previous oculoplastic/plastic surgery clinic visit; iii) most desired cosmetic surgical procedure; and iv) a validated and reliable questionnaire (such as BDDQ).

The BDDQ. The questionnaire used in this study was the BDDQ due to its high sensitivity (100%) and specificity (89-93%) for BDD screening in cosmetics clinics; also, it also showed good validity.²⁵ The BDDQ was the primary survey used to identify patients who exhibited symptoms consistent with BDD. The BDDQ is a validated, brief, self-administered questionnaire that has been extracted from DSM-V diagnostic criteria for BDD developed by Phillips.²⁶ It uses close-ended variables and asks the participants whether their general concerns about appearance are a source of preoccupation. Consequently, it also assesses the grade to which these concerns cause distress or interfere with the person's social or work-related functions. The survey was originally developed among psychiatric patients, but since then it has been validated in a number of different patients, including facial plastic and oculoplastic surgery patient population. Among the plastic surgery and oculoplastic surgery patient population, the survey was established as having a high sensitivity (100%) and high specificity (90.3%).²⁷ The English version of the BDDQ was translated by an expert bilingual and tested for validity, which showed reasonable internal consistency (Cronbach's $\alpha = 0.659$). Body dysmorphic disorder was

diagnosed according to the DSM-V criteria. The scores were calculated in **Table 1**.

The severity scale of BDD was used from 0 to 4 (0 = level 0, 1 = level 1, 2 = level 2, 3 = level 3, 4 = level 4). A score of 4 was regarded as BDD. A score of 0 to 3 was considered negative for BDD. The reliability of the BDD score measurement was calculated using Cronbach's alpha. An alpha value of 0.758 was observed, which indicated an acceptable reliability (**Table 2**).

Furthermore, it is worth noting that BDDQ is not a diagnostic tool. It is a screening tool and a patient who screen positive is a possible case of BDD. A mental health specialist has to confirm the case.

Ethical considerations. This study was carried out after receiving ethical approval from the research ethics committee of King Abdulaziz University, Jeddah, Saudi Arabia. The contribution of patients in this study was entirely on a voluntary basis. Prior to participation in the research, written informed consent was obtained from all patients.

Statistical analysis. Data were checked for completeness and correctness. Categorical data were presented as frequencies and percentages, and

Table 1 - Scoring of body dysmorphic disorder questionnaire (BDDQ).

Question no.	BDDQ question	Answer	BDDQ score
1	Are you very concerned about the appearance of some part(s) of your body that you consider especially unattractive?	Yes	1
2	Do these concerns preoccupy you? That is, do you think about them a lot and wish you could think about them less?	Yes	2
3	Do other people remark your defect?		
4	Has your defect(s) caused you a lot of distress, torment or pain?		
5	Has your defect(s) significantly interfered with your social life?	Yes, to at least one question	3
6	Has your defect(s) significantly interfered with your school work, your job or your ability to function in your role?		
7	How much time do you spend thinking about your defect(s) per day on average?	≥1 hour	4

Table 2 - Reliability of the body dysmorphic disorder questionnaire (BDDQ).

Questions	Item-total statistics				
	Scale mean if item deleted	Scale variance if item deleted	Corrected item-total correlation	Squared multiple correlation	Cronbach's alpha if item deleted
If yes, do these concerns preoccupy you? That is, do you think about them a lot and wish you could think about them less?	2.24	3.005	0.258	0.130	0.773
Do other people remark your defect?	2.63	2.336	0.525	0.338	0.716
Has your defect(s) caused you a lot of distress, torment or pain?	2.50	2.353	0.542	0.336	0.711
as your defect(s) significantly interfered with your social life?	2.86	2.368	0.602	0.410	0.696
Has your defect(s) significantly interfered with your school work, your job or your ability to function in your role?	2.83	2.330	0.607	0.396	0.693
How much time do you spend thinking about your defect(s) per day on average? (time ≥1 hour)	2.59	2.431	0.457	0.229	0.736

Cronbach's alpha: 0.758, Cronbach's alpha based on standardized items: 0.751, number of items: 7

continuous data were presented as mean ± standard deviation. An alpha value of 0.758 was observed, which indicated acceptable reliability. Prevalence rate for all cases and different subgroups were calculated. The age of the respondents were normally distributed according to Shapiro Wilk test. Age was compared between BDD cases and non BDD cases; clinic visit (plastic and reconstructive surgery versus oculoplastic surgery) by independent samples t-test. The rest of the variables were categorical and were compared between the above-mentioned subgroups by Chi-square test. A binary logistic regression was carried out to identify the factors responsible for the development of BDD. The analysis was carried out in 95% confidence interval using the Statistical Package for Social Sciences, version 24.0 (IBM, Armonk, N.Y., USA).

Results. A total of 344 cases were included in this cross-sectional study, and 66 of them had BDD (prevalence rate of 19.2%). The prevalence of BDD among oculoplasty clinic visitors (13.3%) was lower than plastic and reconstructive surgery clinic visitors (21.3%) (Table 3). The majority, 278 cases, did not have BDD, but they had level 1 to 3 BDD as presented in Figure 1. The prevalence of chronic disease was 33.72% among all cases and the most common chronic disease was diabetes mellitus, present in 13.37% cases (Figure 2). All included cases underwent plastic and reconstructive surgery or oculoplastic surgery. The most popular

Table 3 - Prevalence of body dysmorphic disorder among all cases.

Groups/subgroups	Frequency	Prevalence rate
All cases	66	19.2
Oculoplasty	12	13.3
Plastic and reconstructive surgery	54	21.3




Figure 1 - Body dysmorphic disorder questionnaire levels. BDD: body dysmorphic disorder

surgeries underwent or will be undergone by the BDD cases were abdominoplasty (21.2%), and skin lesion removal (21.2%). The most popular surgery among the non-BDD cases were skin lesion removal (16.5%), and blepharoplasty (15.1%) (Figure 3).

The mean age of all cases was 39.66 ± 13.76 years. Age distribution between BDD cases and non-BDD cases was not statistically significantly different, $p=0.260$. A considerably higher percentage of females (93.9%) had BDD compared to males (6.1%), (OR=0.343, $p=0.040$). Smoking was found to be significantly associated with BDD, 21.2% of smokers had it, $p=0.010$. The prevalence of BDD among those who had previous surgery was staggering, 60.6%, $p=0.002$. However, nationality, marital status, education level, monthly household income, and city of residence was not significantly associated with BDD (all p -values of >0.050). Among all cases, 254 (73.8%) were visiting plastic and reconstructive surgery clinic, and the remaining 90 (26.2%) cases were visiting oculoplastic

surgery clinic. These 2 groups were compared in terms of all socio-demographic characteristics. The important observations were as follows: both clinics were predominantly visited by female cases but slightly higher percentage of females visited oculoplastic surgery clinic (93.3%) compared to plastic and reconstructive surgery clinic, (OR=2.774, $p=0.020$). Almost half of the plastic and reconstructive surgery visitors had previous oculoplastic/plastic surgery clinic visit, 126 (49.6%) whereas only 24 (26.7%) of the oculoplastic surgery clinic visitors had previous oculoplastic/plastic surgery clinic visit, (OR=2.707, $p<0.001$). It was also observed that, higher percentage (60.6%) BDD cases previously visited oculoplastic/plastic surgery clinic compared to non-BDD cases (39.6%) ($p=0.002$). The other studied socio-demographic variables were not significantly different between 2 clinic visitors (all p -values of >0.050) (Table 4).

The percentage of cases underwent specific breast-related procedures and face-related procedures were as follows: i) mastopexy 7.6%, breast augmentation 4.7%, breast reconstruction 12.8%; ii) blepharoplasty 13.4%, ptosis repair 3.5%, rhinoplasty 3.5%, chalazion removal 5.2%, and dacryocystorhinoplasty 4.1%. None of these specific surgeries were associated with positive BDD ($p>0.050$) (Table 5). A binary logistic regression was performed to find out the factors collectively predictive of developing BDD. Variables included in the regression analysis were clinical type, age, nationality, gender, marital status, education level, income, smoking, and




Figure 2 - Prevalence of chronic disease among all cases. CHD: coronary heart disease, HTN: hypertension




Figure 2 - Distribution of body dysmorphic disorder (BDD) and no BDD cases by procedure about to undergo/ have undergone /consult the doctor about.

Table 4 - Association between BDD (yes/ no) and sociodemographic variables; clinic type and sociodemographic variables.

Characteristics	All cases	With BDD	Without BDD	P-value	Plastic & reconstructive surgery	Oculoplastic surgery	P-value
Age (mean ± SD)	39.66±13.76	37.94±12.83	40.06±13.97	0.260	38.92±13.30	41.73±14.88	0.096
Nationality							
Saudi	29 (86.6)	60 (90.9)	238 (85.6)	0.317	220 (86.6)	78 (86.7)	0.990
Non-Saudi	46 (13.4)	6 (9.1)	40 (14.4)		34 (13.4)	12 (13.3)	
Gender							
Male	48 (14.0)	4 (6.1)	44 (15.8)	0.040	42 (16.5)	6 (6.7)	0.020
Female	296 (86.0)	62 (93.9)	234 (84.2)		212 (83.5)	84 (93.3)	
Marital status							
Single	102 (29.7)	14 (21.2)	88 (31.7)	0.132	74 (29.1)	28 (31.1)	0.847
Married	188 (54.7)	36 (54.5)	152 (54.7)		142 (55.9)	46 (51.1)	
Divorced	32 (9.3)	10 (15.2)	22 (7.9)		22 (8.7)	10 (11.1)	
Widowed	22 (6.4)	6 (9.1)	16 (5.8)		16 (6.3)	6 (6.7)	
Education level							
Below high school	68 (19.8)	14 (21.2)	54 (19.4)	0.775	40 (15.7)	28 (31.1)	<0.001
High school	78 (22.7)	16 (24.2)	62 (22.3)		54 (21.3)	24 (26.7)	
Bachelor's degree	196 (57.0)	36 (54.5)	160 (57.6)		160 (63.0)	36 (40.0)	
Master/doctorate degree	2 (0.6)	0 (0.0)	2 (0.7)		0 (0.0)	2 (2.2)	
Monthly household income							
<5000 SR	130 (37.8)	28 (42.4)	102 (36.7)	0.522	94 (37.0)	36 (40.0)	0.906
5000-10000 SR	98 (28.5)	14 (21.2)	84 (30.2)		72 (28.3)	26 (28.9)	
10000-20000 SR	78 (22.7)	16 (24.2)	62 (22.3)		60 (23.6)	18 (20.0)	
>20000 SR	38 (11.0)	8 (12.1)	30 (10.8)		28 (11.0)	10 (11.1)	
City							
Jeddah	282 (82.0)	56 (84.8)	226 (81.3)	0.151	212 (83.5)	70 (77.8)	0.015
Makkah	44 (12.8)	10 (15.2)	34 (12.2)		28 (11.0)	16 (17.8)	
Medina	6 (1.7)	0 (0.0)	6 (2.2)		6 (2.4)	0 (0.0)	
Taif	4 (1.2)	0 (0.0)	4 (1.4)		4 (1.6)	0 (0.0)	
Yanbu	2 (0.6)	0 (0.0)	2 (0.7)		0 (0.0)	2 (2.2)	
Dammam	6 (1.7)	0 (0.0)	6 (2.2)		4 (1.6)	2 (2.2)	
Smoking status							
Yes	38 (11.0)	14 (21.2)	24 (8.6)	0.010	36 (14.2)	2 (2.2)	0.003
No	290 (84.3)	48 (72.7)	242 (87.1)		204 (80.3)	86 (95.6)	
Ex-smoker	16 (4.7)	4 (6.1)	12 (4.3)		14 (5.5)	2 (2.2)	
Previous oculoplastic/plastic surgery clinic visit							
Yes	150 (43.6)	40 (60.6)	110 (39.6)	0.002	126 (49.6)	24 (26.7)	<0.001
No	194 (56.4)	26 (39.4)	168 (60.4)		128 (50.4)	66 (73.3)	

Data are presented as numbers and percentages (%). BDD: body dysmorphic disorder, SR: Saudi Riyals

history of previous surgery. The analysis showed that the age (OR=0.943, $p<0.001$) and marital status - single (OR=0.076, $p=0.002$) were the significant predictors of the model (Table 6).

Discussion. Plastic surgeries are often sought by individuals with BDD. This is consistent with the intense surge in the last decade of cosmetic surgical procedures in the United States, Germany, England, and

Table 5 - Relationship between specific breast related procedure, facial procedure and positive body dysmorphic disorder (BDD).

Procedures	Cases underwent %	Have BDD %	Odds ratio	P-value
<i>Breast related</i>				
Mastopexy	7.6	1.16	0.751	0.609
Breast augmentation	4.7	0.58	0.589	0.487
Breast reconstruction	12.8	1.74	0.632	0.317
<i>Face related</i>				
Blepharoplasty	13.4	1.16	0.363	0.052
Ptosis repair	3.5	0.58	0.838	0.822
Rhinoplasty	3.5	0.58	0.838	0.822
Chalazion removal	5.2	1.74	2.217	0.117
Dacryocystorhinoplasty	4.1	0.00	-	0.063

Table 6 - Binary logistic regression analysis to identify the factors responsible for developing body dysmorphic disorder (BDD).

Variables	S.E.	P-value	Odds ratio	95% CI	
				Lower	Upper
Clinic	0.396	0.316	0.672	0.309	1.461
Age	0.017	0.000	0.943	0.913	0.974
Nationality	0.515	0.426	1.507	0.549	4.134
Gender	0.597	0.016	0.237	0.073	0.763
Marital status		0.007			
Single	0.817	0.002	0.076	0.015	0.375
Married	0.603	0.060	0.321	0.098	1.048
Divorced	0.720	0.265	0.448	0.109	1.837
Education level		0.695			
Below high school	0.453	0.394	1.472	0.605	3.577
High school	0.412	0.753	1.139	0.507	2.555
Income		0.178			
<5000 Saudi Riyals	0.566	0.927	0.949	0.313	2.876
5000 -10000 Saudi Riyals	0.561	0.093	0.389	0.130	1.169
>10000-20000 Saudi Riyals	0.535	0.421	0.650	0.228	1.855
Smoking		0.003			
Non-smoker	0.733	0.443	1.755	0.417	7.381
Ex-smoker	0.646	0.158	0.402	0.113	1.424
Previous surgery	0.320	0.081	0.572	0.305	1.072

Brazil.²⁸⁻³¹ However, studies show that most of these patients have a poor level of satisfaction after surgery and very little benefit.^{27,32} Body dysmorphic disorder is considered to be a relative contraindication for undergoing plastic surgery; however, routine screening in practice is not common. This could be attributed to clinicians' underestimation of BDD prevalence and a lack of knowledge of available validated screening tools.²⁷ In our study, the prevalence of patients with positive screening for BDD was 19.2%. This is in contrast to another study conducted in the United States, where the prevalence was 9.7%.³³ This discrepancy might be explained by the fact that their study population included a smaller number of female patients compared to our study. Furthermore, in our study, we found that 93% of female subjects screened positive for BDD. This is consistent with another study by Ishigooka et al,¹⁹ who found a clear relationship between body image and identity in females compared to males. Moreover, it was found by Salehahmadi et al³³ that education levels were the most important influential factor in undergoing plastic surgery followed by age and gender. This agrees with our results, since more than half (54.4%) of the participants who screened positive for BDD in our study population had a bachelor's degree. Of the positively screened participants in our study, 72.7% were non-smokers; this is in contrast to another study conducted in the United States,³⁴ where they found a relationship between perceptions of physical appearance and smoking. They suggested that female smokers are more likely to evaluate both their physical health and appearance negatively. This discrepancy could be due to the higher number of female smokers in the United States (12%),³⁵ compared to Saudi Arabia (9%).³⁶ Interestingly, we found a statistically significant difference between the prevalence of BDD and recurrent visits to oculoplastic/plastic surgery clinics as 60.6% of participants with a positive BDD screening test reported having at least one previous visit to the clinic. This is consistent with the findings of Joseph et al,²⁷ who concluded that most BDD patients were unhappy with the outcome of their surgeries,³¹ which results in recurrent visits and appointments to clinics, seeking further opinions and possible unnecessary cosmetic surgeries. In our study, chronic chalazion removal had the highest association with BDD in the oculoplastic clinic (9.1%), as most of the patients screened positive for BDD reported being inpatient for the conservative methods used for treating such conditions. The second highest association with BDD in the oculoplastic clinic was blepharoplasty (6.1%), which is carried out to alter either the natural anatomy or age-related changes.

On the other hand, we found that in plastic surgery clinics, abdominoplasty and skin lesion removal had the highest associations with BDD (both, 21.2%) followed by liposuction (12.1%). These patients reported being inpatient for seeking the desired body shapes with conservative methods such as diet and exercise. They reported being self-isolated and ashamed with suicidal thoughts. In recent years, with an increasing prevalence of obesity (35.6%),³⁷ and sedentary lifestyles in Saudi Arabia, an increase in the number of BDD patients seeking such surgeries after massive weight loss is expected.

In Saudi Arabia, there are several studies that were conducted on the prevalence of BDD. The first study by AlShahwan,³⁸ concluded that BDD was common among female patients seeking dermatology clinics at King Khalid University hospital. Another study by Kattan et al,³⁹ discovered that cosmetic treatment providers were actually knowledgeable about BDD. In addition, a study that was carried out in Al Ahsa reported the prevalence of BDD among individuals undergoing facial plastic surgery was 14.19%.⁴⁰

Study limitations. Although our sample size was reasonable, and the study objectives were accomplished, this study has a number of limitations. First, this research was performed among patients in a single center; therefore, there is a limitation in the generalizability of the findings. Second, our research was cross-sectional, which covers a limited period of time. Consequently, the study population may have not represented the entire population in Saudi Arabia. We recommend future studies that include different institutes from different regions in Saudi Arabia. In addition, we recommend establishing a routine BDD screening tool in oculoplastic/plastic surgery clinics prior to undergoing any procedures. For health professionals who cannot screen with a tool such as the BDDQ, they can ask appearance specific questions. For example, "does your appearance worry you?" "Do you spend much time worrying about your appearance?"

If a procedure is to be performed, patients should be informed of the possibility of little benefit and poor satisfaction postoperatively. It is recommended for physicians not to make a decision to operate based on patient preference alone, and to prioritize clinical judgment. We recommend future randomized control trial studies to identify high risk groups for BDD. Lastly, we also recommend referring patients with positive BDD screening to psychiatrists for proper evaluation and management.

In conclusion, BDD is most likely an undetected mental health issue among our patients pursuing

cosmetic surgeries in plastic surgery and oculoplastic surgery clinics and a potential source of post-operative patient dissatisfaction. Cosmetic surgeons should consider screening their patients as a standard protocol to guarantee the best outcomes.

Acknowledgment. *The authors gratefully Editage (www.editage.com) for the English language editing.*

References

- Ribeiro RVE. Prevalence of body dysmorphic disorder in plastic surgery and dermatology patients: a systematic review with meta-analysis. *Aesthetic Plast Surg* 2017; 41: 964-970.
- Phillips KA, McElroy SL, Keck PE, Pope HG, Hudson JL. Body dysmorphic disorder: 30 cases of imagined ugliness. *Am J Psychiatr* 1993; 150: 302-308.
- Phillips KA. Quality of life for patients with body dysmorphic disorder. *J Nerv Ment Dis* 2000; 185: 170-175.
- Veale D, Boockock A, Gournay K, Dryden W, Shah F, Willson R, et al. Body dysmorphic disorder. a survey of fifty cases. *Brit J Psychiatr* 1996; 169: 196-201.
- Rosenfield E. Overview of DSM-5 changes. Massachusetts General Hospital OCD and Related Disorders Program 2015. [Updated 2013. Accessed 2020 April 15]. Available from URL: <https://mghocd.org/dsm-5/>
- Phillips KA, Grant J, Siniscalchi J, Albertini RS. Surgical and nonpsychiatric medical treatment of patients with body dysmorphic disorder. *Psychosomatics* 2001; 42: 504-510.
- Cunningham SJ, Bryant CJ, Manisali M, Hunt NP, Feinmann C. Dysmorphophobia: recent developments of interest to the maxillofacial surgeon. *Br J Oral Maxillofac Surg* 1996; 34: 368-374.
- Sarwer DB. Awareness and identification of body dysmorphic disorder by aesthetic surgeons: results of a survey of American society for aesthetic plastic surgery members. *Aesthet Surg J* 2002; 22: 531-535.
- Sarwer DB, Spitzer JC, Sobanko JF, Beer KR. Identification and management of mental health issues by dermatologic surgeons: a survey of American Society for Dermatologic Surgery members. *Dermatol Surg* 2015; 41: 352-357.
- Wright MR. The male aesthetic patient. *Arch Otolaryngol Head Neck Surg* 1987; 113: 724-727.
- de Brito MJ, Neto SM, de Oliveira MF, Cordas TA, Duarte LS, Rosella MF, et al. Yale-brown obsessive compulsive scale modified for body dysmorphic disorder (BDD-YBOCS): Brazilian Portuguese translation, cultural adaptation and validation. *Braz J Psychiatry* 2015; 37: 310-316.
- Morselli PG, Micai A, Boriani F. Eumorphic plastic surgery: expectation versus satisfaction in body dysmorphic disorder. *Aesthetic Plast Surg* 2016; 40: 592-601.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (DSM-5[®]). 5th ed. Washington (DC): American Psychiatric Publishing; 2013. p. 79
- Bouman TK, Mulken S, van der Lei B. Cosmetic professionals' awareness of body dysmorphic disorder. *Plast Reconstr Surg* 2017; 139: 336-342.
- Woolley AJ, Perry JD. Body dysmorphic disorder: Prevalence and outcomes in an oculofacial plastic surgery practice. *Am J Ophthalmol* 2015; 159: 1058.e1-1064.e1.
- Phillips KA, Dufresne RG. Body dysmorphic disorder. A guide for dermatologists and cosmetic surgeons. *Am J Clin Dermatol* 2000; 1: 235-243.
- Dey JK, Ishii M, Phillis M, Byrne PJ, Boahene KD, Ishii LE. Body dysmorphic disorder in a facial plastic and reconstructive surgery clinic: measuring prevalence, assessing comorbidities, and validating a feasible screening instrument. *JAMA Facial Plast Surg* 2015; 17: 137-143.
- Crerand CE, Sarwer DB, Magee L, Gibbons LN, Lowe MR, Barlett SP, et al. Rate of body dysmorphic disorder among patients seeking facial plastic surgery. *Psychiatr Ann* 2004; 34: 958-965.
- Ishigooka J, Iwao M, Suzuki M, Fukuyama Y, Murasaki M, Miura S. Demographic features of patients seeking cosmetic surgery. *Psychiat Clin Neuros* 1998; 52: 283-287.
- Vindigni V, Pavan C, Semenzin M, Gambaro F, Marini M, Bassetto F, et al. The importance of recognizing body dysmorphic disorder in cosmetic surgery patients: do our patients need a preoperative psychiatric evaluation? *Eur J Plast Surg* 2002; 25: 305-308.
- Oosthuizen P, Lambert T, Castle, DJ. Dysmorphic concern: prevalence and associations with clinical variables. *Aust N Z J Psychiatry* 1998; 32: 129-132.
- International OCD Foundation. Assessment Tools for BDD - BDD. [Updated 2020. Accessed 2020 April 17]. Available from URL: <https://bdd.iocdf.org/professionals/assessment-tools-for-bdd/>
- Kaymak Y, Taner E, Şimşek I. Body dysmorphic disorder in university students with skin diseases compared with healthy controls. *Actan Derm Venereol* 2009; 89: 281-284.
- Raosoftware. Sample Size Calculator. [Updated 2008. Accessed 2020 June 17]. Available from URL: <http://www.raosoftware.com/samplesize.html>
- Brohede S, Wingren G, Wijma B, Wijma K. Validation of the body dysmorphic disorder questionnaire in a community sample of Swedish women. *Psychiatry Res* 2013; 210: 647-652.
- Phillips KA. *The Broken Mirror: Understanding and Treating Body Dysmorphic Disorder*. New York (NY): Oxford University Press; 2005. p. 162
- Joseph AW, Ishii L, Joseph SS, Smith JI, Su P, Bater K, et al. Prevalence of body dysmorphic disorder and surgeon diagnostic accuracy in facial plastic and oculoplastic surgery clinics. *JAMA Facial Plast Surg* 2017; 19: 269-274.
- Conrado LA, Hounie AG, Diniz JB, Fossaluza V, Torres AR, Miguel EC, et al. Body dysmorphic disorder among dermatologic patients: prevalence and clinical features. *J Am Acad Dermatol* 2010; 63: 235-243.
- Harth W, Hermes B. Psychosomatic disturbances and cosmetic surgery. *J Dtsch Dermatol Ges* 2007; 5: 736-743.
- Thompson CM, Durrani AJ. An increasing need for early detection of body dysmorphic disorder by all specialties. *J R Soc Med* 2007; 100: 61-62.
- Finger C. Brazilian beauty: Brazil's cosmetic surgery industry is thriving, but why is beauty so important? *Lancet* 2003; 362: 1560.
- Science Daily. Does cosmetic surgery help body dysmorphic disorder? Science Daily 2010. [Updated 2020. Accessed 2020 April 12]. from URL: <https://www.sciencedaily.com/releases/2010/08/100811101424.htm>
- Salehahmadi Z, Rafie SR. Factors affecting patients undergoing cosmetic surgery in Bushehr, Southern Iran. *World J Plast Surg* 2012; 1: 99-106.

34. Strickney S, Black DR. Physical self-perception, body dysmorphic disorder, and smoking behavior. *Am J Health Behav* 2008; 32: 295-304.
35. Centers for Disease Control and Prevention. Current Cigarette Smoking Among Adults in the United States. [Updated 2019 November 18. Accessed 2020 April 10]. Available from URL: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/adult_data/cig_smoking/index.htm
36. Bassiony MM. Smoking in Saudi Arabia. *Saudi Med J* 2009; 30: 876-881.
37. Al-Nozha MM, Al-Mazrou YY, Al-Maatouq MA, Arafah MR, Khalil MZ, Khan NB, et al. Obesity in Saudi Arabia. *Saudi Med J* 2005; 26: 824-829.
38. AlShahwan MA. Prevalence and characteristics of body dysmorphic disorder in Arab dermatology patients. *Saudi Med J* 2020; 41: 73-78.
39. Kattan AE, Alnujaim NH, Barasain O, Bouman TK, AlHammad R, Van der Lei B. Awareness and experiences of cosmetic treatment providers with body dysmorphic disorder in Saudi Arabia. *PeerJ* 2020; 8: e8959.
40. Al Shuhayb ZS. Prevalence of body dysmorphic disorder among Saudis seeking facial plastic surgery. *Saudi Surg J* 2019; 7: 83-86.