

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

4-Carboxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazideHriday Bera,<sup>a</sup> Anton V. Dolzhenko,<sup>a\*</sup> Geok Kheng Tan,<sup>b</sup> Lip Lin Koh<sup>b</sup> and Wai Keung Chui<sup>a</sup><sup>a</sup>Department of Pharmacy, Faculty of Science, National University of Singapore, 18 Science Drive 4, Singapore 117543, Singapore, and <sup>b</sup>Department of Chemistry, Faculty of Science, National University of Singapore, 3 Science Drive 3, Singapore 117543, Singapore

Correspondence e-mail: phada@nus.edu.sg


Received 20 April 2010; accepted 27 April 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.081;  $wR$  factor = 0.218; data-to-parameter ratio = 16.5.

The molecular structure of the title compound,  $\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$ , (systematic name: ethyl *N*-[2-[4-(dimethylamino)benzoyl]hydrazinethiocarbonyl]carbamate) is stabilized by intramolecular  $\text{N}-\text{H}\cdots\text{O}=\text{C}$  hydrogen bonding arranged in an  $S(6)$  graph-set motif. In the crystal, inversion dimers connected *via* intermolecular  $\text{N}-\text{H}\cdots\text{S}=\text{C}$  hydrogen bonds [ $R_2^2(8)$  graph-set motif] form sheets parallel to the  $(\bar{1}21)$  plane. Dimers are also formed by the molecules *via* weak intermolecular  $\text{N}-\text{H}\cdots\text{S}=\text{C}$  hydrogen bonds [ $R_2^2(10)$  graph-set motif] connecting the sheets.

## Related literature

For examples of bioactive 1,4-diacyl substituted thiosemicarbazides and their metal complexes, see: Angelusiu *et al.* (2009); Cunha *et al.* (2007); Qandil *et al.* (2006). For 4-aryloxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazides as high affinity anion receptors, see: Liu & Jiang (2008). For the structures of related carbethoxythioureas, see: Dolzhenko *et al.* (2010*a,b*). For the structures of related 1,4-diacyl thiosemicarbazides, see: Ali *et al.* (2004); Xue *et al.* (2006); Yamin & Yusof (2003); Yusof *et al.* (2003). For the graph-set analysis of hydrogen bonding, see: Bernstein *et al.* (1995).



## Experimental

## Crystal data

$\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$   
 $M_r = 310.37$   
 Triclinic,  $P\bar{1}$   
 $a = 7.876$  (4) Å

$b = 8.184$  (4) Å  
 $c = 12.086$  (6) Å  
 $\alpha = 82.290$  (12)°  
 $\beta = 74.769$  (11)°

$\gamma = 84.469$  (11)°  
 $V = 743.3$  (7) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation

$\mu = 0.23$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.24 \times 0.10 \times 0.08$  mm

## Data collection

Bruker SMART APEX CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 2001)  
 $T_{\min} = 0.946$ ,  $T_{\max} = 0.982$

5201 measured reflections  
 3379 independent reflections  
 2839 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.081$   
 $wR(F^2) = 0.218$   
 $S = 1.18$   
 3379 reflections  
 205 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.84$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.45$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{S1}^{\text{i}}$	0.82 (6)	2.53 (6)	3.342 (3)	173 (5)
$\text{N3}-\text{H3N}\cdots\text{S1}^{\text{ii}}$	0.80 (4)	2.64 (5)	3.385 (4)	156 (4)
$\text{N2}-\text{H2N}\cdots\text{O2}$	0.86 (5)	2.02 (5)	2.653 (4)	130 (4)

Symmetry codes: (i)  $-x + 1, -y + 2, -z + 2$ ; (ii)  $-x, -y + 2, -z + 2$ .

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the National Medical Research Council, Singapore (NMRC/NIG/0019/2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DS2029).

## References

- Ali, H., Yusof, M. S., Khamis, N. A., Mardi, A. S. & Yamin, B. M. (2004). *Acta Cryst.* **E60**, o1656–o1658.
- Angelusiu, M. V., Almajan, G. L., Rosu, T., Negoiu, M., Almajan, E.-R. & Roy, J. (2009). *Eur. J. Med. Chem.* **44**, 3323–3329.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2001). SMART and SAINT. Bruker AXS GmbH, Karlsruhe, Germany.
- Cunha, S., Macedo, F. C. Jr, Costa, G. A. N., Rodrigues, M. T. Jr, Verde, R. B. V., de Souza Neta, L. C., Vencato, I., Lariucci, C. & Sa, F. P. (2007). *Monatsh. Chem.* **138**, 511–516.
- Dolzhenko, A. V., Tan, G. K., Koh, L. L., Dolzhenko, A. V. & Chui, W. K. (2010*a*). *Acta Cryst.* **E66**, o425.
- Dolzhenko, A. V., Tan, G. K., Koh, L. L., Dolzhenko, A. V. & Chui, W. K. (2010*b*). *Acta Cryst.* **E66**, o549–o550.
- Liu, W.-X. & Jiang, Y.-B. (2008). *J. Org. Chem.* **73**, 1124–1127.
- Qandil, A. M., Tumah, H. N. & Hassan, M. A. (2006). *Acta Pharm. Sci.* **48**, 95–107.
- Sheldrick, G. M. (2001). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Xue, S.-J., Bian, W.-D., Wang, Q.-D. & Chai, A. (2006). *Acta Cryst.* **E62**, o4842–o4843.
- Yamin, B. M. & Yusof, M. S. M. (2003). *Acta Cryst.* **E59**, o124–o126.
- Yusof, M. S. M., Yamin, B. M. & Shamsuddin, M. (2003). *Acta Cryst.* **E59**, o810–o811.

**supplementary materials**

*Acta Cryst.* (2010). E66, o1241 [ doi:10.1107/S1600536810015576 ]

## 4-Carboethoxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazide

H. Bera, A. V. Dolzhenko, G. K. Tan, L. L. Koh and W. K. Chui

### Comment

1,4-Diacyl substituted thiosemicarbazides and their metal complexes have been demonstrated to possess a potent antimicrobial activity (Angelusiu *et al.*, 2009; Cunha *et al.*, 2007; Qandil *et al.*, 2006). In continuation of our structural investigations of the carbethoxythioureas derivatives (Dolzhenko *et al.*, 2010*a,b*), we report herein molecular and crystal structure of 4-carboethoxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazide (Figure 1 and 2). The compound is a structural analogue of 4-aryloyl-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazides reported recently as high affinity anion receptors (Liu & Jiang, 2008).

4-Carboethoxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazide was prepared by nucleophilic addition of 4-(*N,N*-dimethylamino)benzhydrazide to ethoxycarbonyl isothiocyanate in DMF at room temperature (Figure 3).

The molecule of 4-carboethoxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazide adopts similar to the previously reported (Ali *et al.*, 2004; Xue *et al.*, 2006; Yamin & Yusof, 2003; Yusof *et al.*, 2003) for the related 1,4-diacyl substituted thiosemicarbazides configuration with the thiocarbonyl group pointed to the side opposite of the carbonyl groups. In the thiourea fragment, (*E*)- and (*Z*)-configurations observed across the C4—N1 and C4—N2 bonds, respectively. This configuration is stabilized by the strong intramolecular hydrogen bonding between N(2)—H and O2=C3 arranged in the *S*(6) graph-set motif (Bernstein *et al.*, 1995).

The thiourea C4—N2 bond is significantly shorter (1.315 (5) Å) than other C—N bonds of the molecule. The planarity of the molecule is affected by some twisting at the hydrazine N2—N3 fragment [—C4—N2—N3—C6— torsion angle is 166.5 (33)°].

In the crystal, the molecules form sheets parallel to the (T21) plane (Figure 2). In the sheets, atom N1 of one molecule is involved in a intermolecular N(1)—H...S=C interaction with the thiocarbonyl atom S1 of adjacent molecule making pair with the  $R_2^2(8)$  graph-set motif. Dimmers are also formed by molecules *via* weak intermolecular N(3)—H...S=C hydrogen bonds arranged in  $R_2^2(10)$  graph-set motifs connecting the sheets between each other.

### Experimental

To a fine suspension of 4-(*N,N*-dimethylamino)benzhydrazide in (0.54 g, 3.0 mmol) in anhydrous DMF (4 ml), ethoxycarbonyl isothiocyanate (0.37 ml, 3.3 mmol) was added. After stirring the mixture for 5 h at ambient temperature, cold water (50 ml) was added. The precipitated product was filtered, washed with cold water and recrystallized from toluene. Yield 0.83 g (89%), m.p. 201 °C (PhMe).

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  1.26 (t, 3H, CH<sub>3</sub>, *J* 7.2 Hz), 2.99 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 4.21 (q, 2H, CH<sub>2</sub>, *J* 7.2 Hz), 6.74 (d, 2H, Ar, *J* 8.7 Hz), 7.78 (d, 2H, Ar, *J* 8.7 Hz), 10.56 (s, 1H, NH), 11.34 (s, 1H, NH), 11.41 (s, 1H, NH).

## supplementary materials

$^{13}\text{C}$  NMR (75 MHz, DMSO- $d_6$ ):  $\delta$  15.2, 40.1 (2 C), 62.7, 111.3 (2 C), 118.6, 129.6 (2 C), 153.1, 153.8, 165.0, 179.8.

### Refinement

All the H atoms attached to the carbon atoms were constrained in a riding motion approximation [0.95 Å for  $\text{C}_{\text{aryl}}\text{—H}$ , 0.99 Å for methylenic protons and 0.98 Å for methyl groups;  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}_{\text{aryl}})$ ,  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}_{\text{methylenic}})$  and  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ ] while the N-bound H atoms were located in a difference map and refined freely.

### Figures




Fig. 1. The molecular structure of 4-carbethoxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazide, showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.



Fig. 2. Crystal packing in the cell (view along axis  $c$ )



Fig. 3. Synthesis of 4-carbethoxy-1-[4-(*N,N*-dimethylamino)benzoyl]thiosemicarbazide

### Ethyl *N*-{2-[4-(dimethylamino)benzoyl]hydrazinethiocarbonyl}carbamate

#### Crystal data

$\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$

$M_r = 310.37$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 7.876$  (4) Å

$b = 8.184$  (4) Å

$c = 12.086$  (6) Å

$\alpha = 82.290$  (12)°

$\beta = 74.769$  (11)°

$\gamma = 84.469$  (11)°

$V = 743.3$  (7) Å<sup>3</sup>

$Z = 2$

$F(000) = 328$

$D_x = 1.387$  Mg m<sup>-3</sup>

Melting point: 474 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 644 reflections

$\theta = 2.5\text{--}27.4^\circ$

$\mu = 0.23$  mm<sup>-1</sup>

$T = 100$  K

Rod, colourless

$0.24 \times 0.10 \times 0.08$  mm

#### Data collection

Bruker SMART APEX CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\varphi$  and  $\omega$  scans

3379 independent reflections

2839 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 1.8^\circ$

Absorption correction: multi-scan  
(SADABS; Sheldrick, 2001)  $h = -9 \rightarrow 10$   
 $T_{\min} = 0.946$ ,  $T_{\max} = 0.982$   $k = -10 \rightarrow 10$   
 5201 measured reflections  $l = -12 \rightarrow 15$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.081$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.218$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.18$	$w = 1/[\sigma^2(F_o^2) + (0.0987P)^2 + 1.3625P]$
3379 reflections	where $P = (F_o^2 + 2F_c^2)/3$
205 parameters	$(\Delta/\sigma)_{\max} < 0.001$
0 restraints	$\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.45 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.25106 (11)	0.89905 (11)	1.00357 (7)	0.0123 (3)
O1	0.6976 (3)	1.2062 (3)	0.7276 (2)	0.0147 (6)
O2	0.4814 (3)	1.1658 (3)	0.6437 (2)	0.0152 (6)
O3	0.1155 (3)	0.9455 (3)	0.6162 (2)	0.0146 (6)
N1	0.4741 (4)	1.0685 (4)	0.8327 (3)	0.0106 (6)
H1N	0.534 (7)	1.072 (7)	0.878 (5)	0.037 (15)*
N2	0.2296 (4)	0.9931 (4)	0.7882 (3)	0.0123 (6)
H2N	0.262 (6)	1.043 (6)	0.719 (4)	0.017 (11)*
N3	0.0781 (4)	0.9079 (4)	0.8084 (3)	0.0138 (7)
H3N	0.005 (6)	0.926 (5)	0.866 (4)	0.011 (10)*
N4	-0.6011 (4)	0.5754 (4)	0.7533 (3)	0.0218 (8)
C1	0.9562 (5)	1.3511 (5)	0.6447 (3)	0.0151 (8)
H1A	0.9156	1.4354	0.6979	0.023*

## supplementary materials

---

H1B	1.0341	1.3997	0.5730	0.023*
H1C	1.0207	1.2596	0.6803	0.023*
C2	0.7995 (5)	1.2877 (5)	0.6182 (3)	0.0138 (7)
H2A	0.7274	1.3800	0.5879	0.017*
H2B	0.8385	1.2083	0.5601	0.017*
C3	0.5467 (5)	1.1491 (4)	0.7259 (3)	0.0109 (7)
C4	0.3184 (4)	0.9900 (4)	0.8667 (3)	0.0119 (7)
C6	0.0248 (4)	0.8950 (4)	0.7109 (3)	0.0109 (7)
C7	-0.1384 (4)	0.8092 (4)	0.7270 (3)	0.0108 (7)
C8	-0.2115 (5)	0.8174 (4)	0.6328 (3)	0.0122 (7)
H8	-0.1558	0.8778	0.5619	0.015*
C9	-0.3619 (5)	0.7403 (5)	0.6401 (3)	0.0137 (7)
H9	-0.4079	0.7473	0.5743	0.016*
C10	-0.4491 (5)	0.6506 (4)	0.7447 (3)	0.0127 (7)
C11	-0.3746 (5)	0.6417 (5)	0.8393 (3)	0.0145 (7)
H11	-0.4295	0.5811	0.9104	0.017*
C12	-0.2221 (5)	0.7201 (5)	0.8304 (3)	0.0124 (7)
H12	-0.1742	0.7129	0.8955	0.015*
C13	-0.6896 (5)	0.4845 (5)	0.8622 (4)	0.0228 (9)
H13A	-0.6084	0.3962	0.8853	0.034*
H13B	-0.7933	0.4363	0.8527	0.034*
H13C	-0.7266	0.5600	0.9219	0.034*
C14	-0.6706 (5)	0.5756 (5)	0.6535 (3)	0.0172 (8)
H14A	-0.6649	0.6861	0.6103	0.026*
H14B	-0.7935	0.5456	0.6789	0.026*
H14C	-0.6004	0.4953	0.6037	0.026*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0110 (4)	0.0190 (5)	0.0080 (4)	-0.0039 (3)	-0.0034 (3)	-0.0013 (3)
O1	0.0134 (12)	0.0172 (13)	0.0144 (13)	-0.0080 (10)	-0.0040 (10)	0.0009 (10)
O2	0.0169 (13)	0.0166 (13)	0.0141 (13)	-0.0069 (10)	-0.0066 (10)	0.0008 (10)
O3	0.0172 (13)	0.0185 (13)	0.0092 (12)	-0.0073 (10)	-0.0032 (10)	-0.0018 (10)
N1	0.0119 (14)	0.0123 (14)	0.0092 (14)	-0.0051 (11)	-0.0045 (12)	-0.0004 (11)
N2	0.0122 (14)	0.0179 (15)	0.0072 (14)	-0.0068 (12)	-0.0028 (11)	0.0024 (12)
N3	0.0067 (13)	0.0220 (17)	0.0128 (16)	-0.0060 (12)	-0.0001 (12)	-0.0032 (13)
N4	0.0201 (17)	0.0303 (19)	0.0176 (17)	-0.0149 (15)	-0.0078 (14)	0.0034 (14)
C1	0.0134 (16)	0.0133 (17)	0.0183 (19)	-0.0044 (13)	-0.0034 (14)	0.0004 (14)
C2	0.0167 (17)	0.0143 (17)	0.0107 (17)	-0.0068 (14)	-0.0019 (14)	-0.0010 (13)
C3	0.0129 (16)	0.0051 (15)	0.0149 (17)	-0.0016 (12)	-0.0018 (13)	-0.0045 (12)
C4	0.0110 (16)	0.0113 (16)	0.0133 (17)	0.0005 (13)	-0.0031 (13)	-0.0024 (13)
C6	0.0109 (15)	0.0101 (16)	0.0122 (17)	-0.0015 (13)	-0.0016 (13)	-0.0050 (13)
C7	0.0091 (15)	0.0122 (16)	0.0104 (16)	-0.0028 (13)	-0.0004 (13)	-0.0012 (13)
C8	0.0130 (16)	0.0089 (16)	0.0142 (17)	0.0012 (13)	-0.0012 (13)	-0.0048 (13)
C9	0.0164 (17)	0.0180 (18)	0.0108 (17)	-0.0025 (14)	-0.0078 (14)	-0.0064 (14)
C10	0.0129 (16)	0.0119 (16)	0.0144 (18)	-0.0030 (13)	-0.0040 (14)	-0.0025 (13)
C11	0.0159 (17)	0.0163 (18)	0.0106 (17)	-0.0069 (14)	-0.0024 (14)	0.0028 (14)

C12	0.0111 (16)	0.0180 (18)	0.0096 (16)	-0.0035 (13)	-0.0049 (13)	-0.0004 (13)
C13	0.0180 (19)	0.025 (2)	0.026 (2)	-0.0121 (16)	-0.0032 (16)	-0.0026 (17)
C14	0.0189 (18)	0.0167 (18)	0.021 (2)	-0.0042 (14)	-0.0125 (15)	-0.0029 (15)

*Geometric parameters (Å, °)*

S1—C4	1.690 (4)	C2—H2A	0.9900
O1—C3	1.325 (4)	C2—H2B	0.9900
O1—C2	1.465 (4)	C6—C7	1.479 (5)
O2—C3	1.219 (4)	C7—C12	1.394 (5)
O3—C6	1.221 (4)	C7—C8	1.396 (5)
N1—C3	1.374 (5)	C8—C9	1.372 (5)
N1—C4	1.379 (4)	C8—H8	0.9500
N1—H1N	0.82 (6)	C9—C10	1.414 (5)
N2—C4	1.315 (5)	C9—H9	0.9500
N2—N3	1.390 (4)	C10—C11	1.407 (5)
N2—H2N	0.86 (5)	C11—C12	1.389 (5)
N3—C6	1.371 (5)	C11—H11	0.9500
N3—H3N	0.80 (4)	C12—H12	0.9500
N4—C10	1.372 (5)	C13—H13A	0.9800
N4—C14	1.450 (5)	C13—H13B	0.9800
N4—C13	1.458 (5)	C13—H13C	0.9800
C1—C2	1.507 (5)	C14—H14A	0.9800
C1—H1A	0.9800	C14—H14B	0.9800
C1—H1B	0.9800	C14—H14C	0.9800
C1—H1C	0.9800		
C3—O1—C2	116.3 (3)	O3—C6—C7	123.2 (3)
C3—N1—C4	127.1 (3)	N3—C6—C7	116.7 (3)
C3—N1—H1N	112 (4)	C12—C7—C8	118.4 (3)
C4—N1—H1N	121 (4)	C12—C7—C6	123.7 (3)
C4—N2—N3	122.1 (3)	C8—C7—C6	117.9 (3)
C4—N2—H2N	124 (3)	C9—C8—C7	121.6 (3)
N3—N2—H2N	114 (3)	C9—C8—H8	119.2
C6—N3—N2	114.1 (3)	C7—C8—H8	119.2
C6—N3—H3N	119 (3)	C8—C9—C10	120.6 (3)
N2—N3—H3N	114 (3)	C8—C9—H9	119.7
C10—N4—C14	121.0 (3)	C10—C9—H9	119.7
C10—N4—C13	120.2 (3)	N4—C10—C11	121.3 (3)
C14—N4—C13	118.7 (3)	N4—C10—C9	121.0 (3)
C2—C1—H1A	109.5	C11—C10—C9	117.7 (3)
C2—C1—H1B	109.5	C12—C11—C10	121.0 (3)
H1A—C1—H1B	109.5	C12—C11—H11	119.5
C2—C1—H1C	109.5	C10—C11—H11	119.5
H1A—C1—H1C	109.5	C11—C12—C7	120.7 (3)
H1B—C1—H1C	109.5	C11—C12—H12	119.7
O1—C2—C1	106.0 (3)	C7—C12—H12	119.7
O1—C2—H2A	110.5	N4—C13—H13A	109.5
C1—C2—H2A	110.5	N4—C13—H13B	109.5
O1—C2—H2B	110.5	H13A—C13—H13B	109.5

## supplementary materials

C1—C2—H2B	110.5	N4—C13—H13C	109.5
H2A—C2—H2B	108.7	H13A—C13—H13C	109.5
O2—C3—O1	126.1 (3)	H13B—C13—H13C	109.5
O2—C3—N1	125.2 (3)	N4—C14—H14A	109.5
O1—C3—N1	108.7 (3)	N4—C14—H14B	109.5
N2—C4—N1	116.4 (3)	H14A—C14—H14B	109.5
N2—C4—S1	123.9 (3)	N4—C14—H14C	109.5
N1—C4—S1	119.7 (3)	H14A—C14—H14C	109.5
O3—C6—N3	120.0 (3)	H14B—C14—H14C	109.5
C4—N2—N3—C6	-166.5 (3)	N3—C6—C7—C8	169.8 (3)
C3—O1—C2—C1	176.1 (3)	C12—C7—C8—C9	0.0 (5)
C2—O1—C3—O2	-4.1 (5)	C6—C7—C8—C9	179.5 (3)
C2—O1—C3—N1	176.4 (3)	C7—C8—C9—C10	0.6 (5)
C4—N1—C3—O2	1.2 (6)	C14—N4—C10—C11	-175.8 (3)
C4—N1—C3—O1	-179.3 (3)	C13—N4—C10—C11	0.3 (6)
N3—N2—C4—N1	174.8 (3)	C14—N4—C10—C9	4.3 (6)
N3—N2—C4—S1	-6.0 (5)	C13—N4—C10—C9	-179.5 (4)
C3—N1—C4—N2	-0.2 (5)	C8—C9—C10—N4	178.9 (3)
C3—N1—C4—S1	-179.4 (3)	C8—C9—C10—C11	-0.9 (5)
N2—N3—C6—O3	5.2 (5)	N4—C10—C11—C12	-179.1 (4)
N2—N3—C6—C7	-178.5 (3)	C9—C10—C11—C12	0.8 (5)
O3—C6—C7—C12	165.6 (4)	C10—C11—C12—C7	-0.3 (6)
N3—C6—C7—C12	-10.6 (5)	C8—C7—C12—C11	-0.1 (5)
O3—C6—C7—C8	-13.9 (5)	C6—C7—C12—C11	-179.6 (3)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N $\cdots$ S1 <sup>i</sup>	0.82 (6)	2.53 (6)	3.342 (3)	173 (5)
N3—H3N $\cdots$ S1 <sup>ii</sup>	0.80 (4)	2.64 (5)	3.385 (4)	156 (4)
N2—H2N $\cdots$ O2	0.86 (5)	2.02 (5)	2.653 (4)	130 (4)

Symmetry codes: (i)  $-x+1, -y+2, -z+2$ ; (ii)  $-x, -y+2, -z+2$ .



Fig. 1




Fig. 2




Fig. 3

