

POSTER PRESENTATION

Open Access

Impact of biologics on growth in children with juvenile idiopathic arthritis

F Uettwiller^{1*}, G Pinto², M Polak², P Quartier³

From 18th Pediatric Rheumatology European Society (PReS) Congress
Bruges, Belgium. 14-18 September 2011

Background

Juvenile idiopathic arthritis (JIA) is sometimes associated with growth impairment due to inflammation and/or steroid treatment. Biologics allow a better control of inflammation and may therefore have a positive effect on growth.

Aim

To assess growth of JIA children on biologics.

Methods

JIA patients followed in one reference center, who were prepubescent at onset of biologic treatment, were included. We collected data about their growth and their disease (course, treatments, activity). Heights were expressed in standard deviations (SD) corrected with target height, at disease onset, at onset of biologics and at last follow-up.

Results

100 patients were included. 29% were systemic. Mean age and height were 4.26 years (0.7; 13.7) and 0.31 SD (-2.47; 5.46) respectively at disease onset, 7.1 years (1.6; 15.7) and -0.24 SD (-3.63; 2.90) at onset of biologics ($p<0.05$), 11 years (2.3; 19.5) and -0.15 SD (-4.95; 3.52) at last follow-up. 93 had received anti-TNF α , 35 anti-IL1, 12 anti-IL-6 and 9 CTLA4-Ig. 45% had an active disease. More than one biological agent was statistically linked with poor growth. At last follow-up, 10 children had a growth delay and 6 were treated with growth hormone.

Conclusion

Growth delay was observed before biologics were started. On biologic treatment, growth velocity normalized, however there was no correction of growth delay.

The discrepancy with previous studies, which showed a growth restoration on anti-TNF, might be linked to a higher proportion of systemic and severe polyarticular JIA patients. In a subset of patients, growth hormone treatment has to be discussed even at the era of biologics.

Author details

¹Unité de Spécialités Pédiatriques, CHU Clocheville, Tours, France.

²Endocrinologie Pédiatrique, Necker Enfants Malades, Paris, France. ³Unité d'Immunologie, d'Hématologie et de Rhumatologie Pédiatriques, Necker Enfants Malades, Paris, France.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P139

Cite this article as: Uettwiller et al.: Impact of biologics on growth in children with juvenile idiopathic arthritis. *Pediatric Rheumatology* 2011 **9** (Suppl 1):P139.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Unité de Spécialités Pédiatriques, CHU Clocheville, Tours, France
Full list of author information is available at the end of the article