

Chemopreventive Action of Anthocyanin-rich Black Soybean Fraction in $APC^{Min/+}$ Intestinal Polyposis Model

ORIGINAL
ARTICLEMi-Young Park^{1,*}, Jung-Mi Kim^{2,*}, Jong-Sang Kim³, Myoung-Gun Choung⁴, Mi-Kyung Sung²

¹Department of Food and Nutrition Education, Graduate School of Education, Soonchunhyang University, Asan, ²Department of Food and Nutrition, College of Human Ecology, Sookmyung Women's University, Seoul, ³Department of Animal Science and Biotechnology, The Graduate School of Kyungpook National University, Daegu, ⁴National Yeongnam Agricultural Experiment Station, Rural Development Administration, Miryang, Korea

Background: Anthocyanins have been shown to inhibit cancer cell growth by suppressing oxidative stress and inflammatory responses. The purpose of this study was to investigate the effects of an anthocyanin-rich extract (AE) from black soybean coat on intestinal carcinogenesis.

Methods: $APC^{Min/+}$ mice were fed a diet of 0.2% or 0.5% AE for 7 weeks. We analyzed the number of intestinal tumors, oxidative stress and inflammatory markers associated with β -catenin and cytosolic phospholipase A₂ (cPLA₂) signals. The number of intestinal tumors, and cellular expression of β -catenin were determined.

Results: The number of intestinal tumors was significantly lower in mice fed a 0.5% AE diet compared to those of the other groups. Cytosolic β -catenin expression was significantly decreased in the AE supplemented groups compared to that of the control animals. In addition, mucosa expression of cyclooxygenase-2 and cPLA₂ were also significantly decreased in the 0.5% AE group, by 32% and 62%, respectively, compared to the control group.

Conclusions: These results suggest that dietary AE reduced the development of intestinal tumors, possibly through the ability to suppress oxidative stresses, decreasing inflammatory responses mediated by β -catenin associated signals.

(J Cancer Prev 2015;20:193-201)

Key Words: Anthocyanins, $APC^{Min/+}$, Black soybean, Inflammation, Intestinal tumors

INTRODUCTION

Environmental factors, including a western style diet rich in animal fat and protein, increase the risk of colon, breast and prostate cancers. Colorectal cancer was the fourth highest cause of cancer deaths worldwide in 2012.¹ Although plant food-based diets are known to reduce the risk of developing colorectal cancer,^{2,3} the cause-effect evidence is not strong enough to show a negative association. This may be due to complex interactions between individual plant food components to exert biological responses. In addition, most of the efficacy evaluation studies used an in vitro cell culture system, which does not accurately

reflect complex biological systems.

Soybeans are one of the most frequently studied legumes for their biological activities. Black soybeans have been widely used as materials in oriental medicine. A distinctive difference between yellow and black soybeans is their anthocyanin contents, which are responsible for the dark-blue colors exhibited by the plants. Anthocyanins are especially abundant in the epidermis palisade layer of the black soybean coat,⁴ synthesized via the phenylpropanoid pathway.⁵ The most abundant anthocyanins in plant foods are cyanidin-3-glucoside, delphinidin-3-glucoside and melvidin-3-glucoside, all of which are known to possess strong antioxidant and anticarcinogenic activity.⁴ It is reported that

Received July 16, 2015. Revised August 11, 2015. Accepted August 11, 2015

Correspondence to: Mi-Kyung Sung

Department of Food and Nutrition, College of Human Ecology, Sookmyung Women's University, 100 Cheongpa-ro 47-gil, Yongsan-gu, Seoul 04310, Korea
Tel: +82-2-710-9395, Fax: +82-2-710-9453, E-mail: mksung@sm.ac.kr, ORCID: Mi-Kyung Sung, <http://orcid.org/0000-0002-3575-5628>

*These authors contributed equally to this work as co-first authors.

Copyright © 2015 Korean Society of Cancer Prevention

©This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

administration of bilberry anthocyanins decreased tumor proliferation in colorectal cancer patients.⁶ Previous studies also has been reported that anti-carcinogenesis effects of anthocyanins from several plant foods such as sweet potato, grapes, tart cherry and berries in colon cancer animal models.⁷⁻¹¹ However, few mechanistic explanations on the inhibitory effects of anthocyanins on tumor growth in vivo have been provided.

We previously demonstrated that anthocyanidins and anthocyanin-rich extract (AE) from black soybean coat exerted anti-inflammatory and anti-proliferative effects in colon cancer cells.¹² The biological properties of anthocyanins are due to their peculiar chemical structure, as they are excellent scavengers with reactivity towards reactive oxygen species (ROS).¹³ One of the best described cellular effects of ROS is the oxidative modification of fatty acids within membrane phospholipids, altering membrane fluidity, protein structure and cell signaling.¹⁴ ROS also activate nuclear transcription factors such as AP-1 and NF- κ B, which in turn induce the expression of inflammatory target proteins, including cytosolic phospholipase A₂ (cPLA₂).¹⁵ cPLA₂ hydrolyzes the *sn*-2 ester bond in phospholipids, releasing free fatty acids and lysophospholipids.¹⁶ Increased cPLA₂ activity gives rise to a number of biologically active mediators, such as arachidonic acid (AA)-derived metabolites, including prostaglandins and lysophosphatidylcholine. For this reason, cPLA₂ has an important role in the modulation of membrane stability, as well as in the formation of inflammatory prostaglandins.

Apc^{Min/+} mice exhibit a germ-line nonsense mutation at codon 850 of the adenomatous polyposis coli (*APC*) gene, causing the spontaneous development of multiple polyps in the small and large intestines at the age of 10 to 12 weeks.¹⁷ Mutations in the *APC* tumor suppressor gene occur very early in cancer, and are found in a majority of sporadic colorectal tumors, as well as in familial adenomatous polyposis (FAP), a hereditary form of intestinal cancer.¹⁸ The main tumor suppressing activity of the APC protein is due to its ability to down regulate cellular β -catenin.¹⁹ Cytosolic β -catenin can be translocated into the nucleus, where it interacts with the transcriptional factor Tcf/Lef to stimulate transcription of inflammatory response genes, including cyclooxygenase-2 (COX-2). Enhanced levels of β -catenin and several inflammatory mediators are favored factors for the development of colorectal cancer.²⁰ Previous studies have also indicated that ROS-induced activation of nuclear transcription factors increase the expression of COX-2. COX-2 converts AA to prostaglandin E₂ (PGE₂), mediating inflammatory responses in colonic carcinogenesis.^{21,22} AA is liberated from membrane phospholipids by the action of cPLA₂. Since the expression of

cPLA₂ is shown to be up-regulated by ROS,²³ decreasing the oxidative stress by intake of anthocyanins may reduce the synthesis of PGE₂, suppressing colon carcinogenesis.

In the present study, we investigated the inhibitory activity and the potential underlying mechanism of action of an AE from black soybean coat against intestinal tumorigenesis in *Apc*^{Min/+} mice.

MATERIALS AND METHODS

1. Animals

The mice were bred in Sookmyung Women's University, from inbred mice originally purchased from the Jackson Laboratory (Bar Harbor, ME, USA). The *Apc*^{Min/+} pedigree was maintained by mating *Apc*^{+/+} females with *Apc*^{Min/+} males, and procedures to secure inbreeding were followed. The *Apc*^{Min/+} mice were identified by allele-specific PCR on DNA isolated from the tail. All animals were housed in wire cages (3 mice/cage) with free access to drinking water and basal diet CE-2 (CLEA Japan Inc., Tokyo, Japan), under controlled conditions of humidity (50% \pm 10%), lighting (12 hours light/dark cycle), and temperature (23°C \pm 2°C), which were maintained until termination of the study.

2. Preparation of anthocyanin-rich extract from black soybean seed coat

AE was prepared as previously described.⁶ Briefly, the seed coats of black soybean were peeled manually after heating at 105°C for 2 hours. The separated seed coats were extracted three times with 1,000 mL of 1% HCl-40% CH₃OH at 4°C for 24 hours. The combined extracts were filtered and concentrated at 30°C in vacuo. The main anthocyanin contents in the seed coats of the black soybeans were determined by measurement of the HPLC peak area, monitored at 530 nm. Cyanidin-3-glucoside, delphinidin-3-glucoside, and petunidin-3-glucoside content were determined to be 0.38, 0.16, and 0.05 mg/g black soybean seed coat, respectively.

3. Study design

A total of 30 *Apc*^{Min/+} male mice (5 weeks old) were used in this study. The *Apc*^{Min/+} mice were randomly assigned to three groups (ten mice per group) following a 1 week acclimatization period. The control group of *Apc*^{Min/+} mice were fed the AIN-93G control diet, and 2 groups of *Apc*^{Min/+} mice were fed AE diets at different concentrations (0.2% and 0.5%, respectively). Diets were prepared by replacing sucrose in the AIN-93G diet with powdered AE. All mice were fed these diets for 7 weeks. Food intake were measured every 2nd day after diet is replaced with fresh diet. Body weight

is measured every other day. At the end of the experimental period, the mice were sacrificed, and the intestines and colons were harvested. After scoring of tumors, the small intestinal and colonic mucosa was scraped with blades and kept at -80°C to use for mRNA and protein analysis. Blood was also obtained for biochemical analysis. The experiment was approved by the Animal Care and Use Committee of the Department of Food and Nutrition of Sookmyung Women's University (Seoul, Korea).

4. Tumor scoring

All organs were removed at sacrifice, and the small and large intestines were cut open along the longitudinal axis. The small intestine and colon were then spread flat on a microscope slide, and the number of tumors was counted under an inverse light microscope at a magnification of 1×10 , while blind to the dietary treatment.

5. Measurement of cytosolic phospholipase A₂ activity

cPLA₂ activity was determined in the cytosolic fraction using a cPLA₂ assay kit (Cayman Chemical, Ann Arbor, MI, USA). Arachidonoyl thio-PC (1-*O*-hexadecyl-2-deoxy-2-thio-*R*-(arachidonoyl)-*sn*-glycerol-3-phosphoryl-choline) was used as a synthetic substrate to detect PLA₂ activity. Hydrolysis of the arachidonoyl thioester bond at the *sn*-2 position by PLA₂ releases free thiol, which is detected by Ellman's reagent. Activity was reported as nanomol per minute per milligram of cytosolic protein.

6. RNA extraction and semi-quantitative reverse transcription-PCR

Total RNA was isolated from mouse tissues using Trizol (Invitrogen, Carlsbad, CA, USA), following the manufacturer's recommendations. Single-stranded cDNAs were generated from total RNA using the superscript first-strand synthesis system (Invitrogen). An aliquot of 100 ng of RNA was used for reverse transcription, and subsequent amplification was carried out using 1 μL of cDNA in a PCR at a final volume of 20 μL with an MJ Research PTC-0150 MiniCycler (MJ Research Inc., Waltham, MA, USA), using HotStarTaq (Intron Biotechnology Inc, Seongnam, Korea). PCR oligonucleotide sequences for intestinal mucosa used were as follows: COX-2 (forward 5'-TGTATCCCCACAGTCAAAGACAC-3', reverse 5'-GTGCTCCCGAAGCCAGATGG-3'), cPLA₂ (forward 5'-GTGTCTGGGCAGTGCCTTT-3', reverse 5'-GTTGAAAATGGCGATTCTGGG-3'), β -actin (forward 5'-TGTGAGGTGGGAATGGGTCAG-3', reverse 5'-TTTGATGTCACGCACGATTTCC-3'). The conditions of PCR for the target genes were as follows: COX-2 gene: 30 cycles at 94°C for 1 minute, 62°C for 1 minute, and 72°C for 2 minutes;

cPLA₂ gene: 30 cycles at 94°C for 1 minute, 60°C for 1 minute, and 72°C for 2 minutes. The termination cycle included a prolonged extension at 72°C for 5 minutes. PCR products were confirmed by 1.6% agarose gel electrophoresis, and visualized by UV transillumination (Bio-Rad Laboratories Inc., Hercules, CA, USA). All signals were normalized to the mRNA levels of the housekeeping gene β -actin, and expressed as a ratio.

7. Biochemistry of intestinal mucosa

Because the subcellular localization of β -catenin is considered to be an important determinant in cancer, β -catenin was analyzed separately from cytosolic and membranous fractions of adenoma and normal-appearing mucosa. Small intestinal and colonic mucosa homogenates were centrifuged for 10 minutes at 200 *g* to remove intact cells. Cleared supernatants were centrifuged further for 10 minutes at 1,000 *g*. Supernatants were used for cytosolic and membranous fraction isolations, and pellets for nuclear fraction isolations. Ice-cold homogenization buffer was added to the supernatants to make a final volume of 5 mL. These mixtures were centrifuged for 60 minutes at 100,000 *g*. Newly formed supernatants were collected and used as the cytosolic fractions of the mucosal epithelial cells. The pellets were resuspended in 5 mL of ice-cold homogenization buffer containing 2 mL/L Triton X-100, incubated for 30 minutes and centrifuged for 60 minutes at 100,000 *g*. The resulting supernatants were used as the membranous fractions of the mucosal epithelial cells. The pellets formed in the first step of the procedure containing the nuclear fractions were resuspended in 1.8 mL of ice-cold homogenization buffer containing 2 mL/L Triton X-100, incubated for 20 minutes and centrifuged for 10 minutes at 15,000 *g*. The resulting supernatants were used as the nuclear fractions of the mucosal epithelial cells. All steps were performed at 4°C . All fractions were concentrated using Amicon Ultra-4 Centrifugal Filter Devices (Millipore, Bedford, MA, USA), and the protein concentrations of the resulting samples were measured using a Bio-Rad protein assay reagent (Bio-Rad Laboratories Inc.).

8. Western blot analysis

For the immunoblotting analysis of β -catenin, 30 μg of the sample proteins were electrophoresed through 7.5% SDS PAGE and transferred to polyvinylidenedifluoride membranes (Amersham, Arlington Heights, IL, USA). The transferred membrane was blocked using 2% skim milk to inhibit non-specific proteins, and primary antibodies were added. β -catenin (BD Transduction, San Diego, CA, USA), β -actin, and β -tubulin (Cell Signaling, Beverly, MA, USA) were used as the primary antibodies. Protein detection

was carried out using secondary anti-mouse HPR antibodies (Zymed Laboratories, South San Francisco, CA, USA). Each protein band was then confirmed and quantified using an enhanced chemiluminescence system (Amersham). Blots were scanned and analyzed using a multiple image analyzer and the Quantity One program (Bio-Rad Laboratories Inc.).

9. Measurement of serum prostaglandin E₂ level

The PGE₂ levels in the serum were measured using commercially available ELISA kits (R&D, Ann Arbor, MI, USA). This assay is based on the competitive binding technique, in which the PGE₂ present in a sample competes with a fixed amount of horseradish peroxidase labeled PGE₂ for the sites on a mouse monoclonal antibody. The absorbance was determined at 450 nm, with the wavelength correction set at 540 nm.

10. Statistical analysis

For the results of all experimental analyses, means and standard errors in each group were calculated using an SAS program (SAS Institute, Cary, NC, USA). To determine the statistical significance of differences between the control and AE-treated groups, one-way analysis of variance was performed, followed by the Student's *t*-test. *P*-values less than 0.05 were considered to be statistically significant.

RESULTS

1. Effect of dietary anthocyanin-rich extract on food intake, body weight, intestinal length, liver weight and tumor formation

As indicated in Table 1, there was no significant differences in the body weight gain between the groups. Further, there were no considerable changes in liver weight and intestinal length (Table 1). The food intake was measured every 2nd day after replacing the feed with fresh diet. The average food consumption for 2 days were as follows: control group, 7.84 ± 0.47 g; 0.2% AE group, 8.00

Table 1. Effects of dietary AE on body weight, intestinal length and liver weight in *APC^{Min/+}* mice

Group	Body weight (g)	Intestinal length (cm)	Liver weight (g)
Control	30.64 ± 1.04 ^a	39.31 ± 1.11 ^a	9.34 ± 0.31 ^a
0.2% AE	33.11 ± 1.13	38.73 ± 1.18	9.02 ± 0.19
0.5% AE	26.30 ± 1.72	38.74 ± 1.14	9.09 ± 0.16

Values are presented as mean ± SE. The experimental mice were fed 0.2% or 0.5% anthocyanin-rich extract (AE) supplemented diets for 7 weeks. Statistical significance of the differences was evaluated by *t*-test. ^aNS, not significant.

± 0.51 g; 0.5% AE group, 8.10 ± 0.55 g and there is no significant difference. The effect of dietary AE from black soybean coats on tumor formation was assessed at week 7 after supplying the different experimental diets. Results showed that the average number of tumors were 41.3 ± 8.14, 34.7 ± 14.51, and 28.3 ± 11.50 per mouse fed either control diet, 0.2% AE diet or 0.5% AE diet, respectively (Fig. 1). Although we did not differentially counted the tumor numbers from small and large intestine separately, more than 90% of the tumors were located in the small intestine. We could find only 2 to 3 polyps from large intestine. The reduction observed with the 0.2% anthocyanin extract diet was not statistically significant; however, mice fed the 0.5% AE diet had a significant decrease in the number of intestinal tumors (*P* < 0.05; Fig. 1).

2. Effect of dietary anthocyanin-rich extract on intestinal β-catenin expression

Cytosolic and nucleic β-catenin protein expression levels were determined, revealing that the cytosolic β-catenin expression was significantly decreased in animals fed either 0.2% AE (*P* < 0.01) or 0.5% AE (*P* < 0.001; Fig. 2A), compared to the control. However, nucleic β-catenin protein level was not significantly different between the groups (Fig. 2B).

3. Effect of dietary anthocyanin-rich extract on intestinal cytosolic phospholipase A₂ activity and mRNA expression

The effects of an AE supplemented diet on cPLA₂ activity and cPLA₂ mRNA expression in the intestinal mucosa were determined.




Figure 1. Effect of dietary anthocyanin-rich extract (AE) on tumor formation in *APC^{Min/+}* mice. The mice were fed 0.2% or 0.5% AE supplemented diets for 7 weeks. After 7 weeks, the number of tumors was scored. Each bar represents the mean ± SE; n = 10 mice for each group. Statistical significance of the differences was evaluated by *t*-test (**P* < 0.05). NS, not significant.




Figure 2. Effect of dietary anthocyanin-rich extract (AE) on (A) cytosolic β -catenin protein and (B) nuclear β -catenin protein level in $Apc^{Min/+}$ mice. The mice were fed 0.2% or 0.5% AE supplemented diets for 7 weeks. After 7 weeks, (A) cytosolic β -catenin protein and (B) nuclear β -catenin protein level in the intestines were measured. The white bars indicate non-supplemented diets, while the black bars represents diets supplemented with AE. Each bar represents the mean \pm SE; $n = 10$ mice for each group. The experiments are repeated 3 times. Statistical significance of differences was evaluated by t -test (** $P < 0.01$, *** $P < 0.001$). NS, not significant.

Results showed that cPLA₂ activity was decreased in both the 0.2% and 0.5% AE supplemented group; however, the results were not statistically significant (Fig. 3A). In contrast to activity, the cPLA₂ mRNA expression was significantly reduced in the 0.5% AE-fed group by 62% compared to that of the control group ($P < 0.05$, Fig. 3B).

4. Effect of dietary anthocyanin-rich extract on intestinal COX-2 mRNA expression

COX-2 transforms membrane AA to cause a pro-inflammatory response. Figure 4A shows that the diet of 0.5% AE significantly decreased the expression of intestinal COX-2 mRNA by 32% ($P < 0.05$) compared with the control diet.

5. Effect of dietary anthocyanin-rich extract on serum prostaglandin E₂ and nitric oxide

COX-2 generates PGE₂ and production of PGE₂ is related to various pathogenic processes, and to the stability of the cell membrane. The results of this study showed that serum PGE₂ levels were not statistically significant in AE supplemented groups (Fig. 4B).

DISCUSSION

We previously reported that anthocyanins or anthocyanin-rich black soybeans inhibited colon cancer cell growth and the formation of precancerous aberrant crypt foci (ACF) by suppressing

inflammatory response.¹² In this study, we further investigated the AE of black soybean coat on colon tumor formation in $APC^{Min/+}$ mice.

The results of this study suggest that the dietary administration of AE in this study significantly decreased the cytosolic β -catenin level, with no significant difference in nuclear β -catenin expression. Although β -catenin nuclear translocation has been used as a marker for colon cancer in many studies, the relationship between cytosolic and nuclear β -catenin accumulation in intestinal tumor development has not been clearly defined. In a β -catenin expression pattern study, most colorectal tumors demonstrated increased β -catenin in the cytosolic fraction, while nuclear β -catenin expression was found in only 64% of primary colorectal tumors and 21% of corresponding liver metastases,²⁴ suggesting intestinal tumor formation and growth are correlated more with the cytosolic levels of β -catenin than with the nuclear levels. It was also previously suggested that nuclear accumulation of β -catenin is the very first event of tumor development, followed by a substantial increase of β -catenin in the entire cytosol.²⁵ Another study showed that the nuclear β -catenin level was significantly correlated with the number of adenomas at the early stage of tumor development, while the cytosolic, but not nuclear or membranous β -catenin, was strong correlated with adenoma growth at the later developmental stages in $APC^{Min/+}$ mice.¹⁹ In this regard, it should be noted that cytoplasmic expression of β -catenin may be the most frequent abnormality in human ACF, adenomas, and carcinomas development.²⁶ There




Figure 3. Effect of dietary anthocyanin-rich extract (AE) on (A) cytosolic phospholipase A2 (cPLA₂) activity and (B) cPLA₂ mRNA expression in *Apc^{Min/+}* mice. The mice were fed 0.2% or 0.5% AE supplemented diets for 7 weeks. After 7 weeks, (A) cPLA₂ activity and (B) cPLA₂ mRNA expression in the intestinal mucosa were measured. Each bar represents the mean \pm SE; $n = 10$ mice for each group. The experiments are repeated 3 times. Statistical significance of the differences was evaluated by *t*-test (* $P < 0.05$). NS, not significant.

are few studies that show that the direct effects of anthocyanins on the activation of β -catenin. Two previous studies showed that anthocyanins from black raspberry and fruits of *Vitis coignetiae* Pulliat effectively suppressed β -catenin expression in colon cancer cells and uterine cervical cancer cells, respectively.^{27,28} Here, we found that the AE diet specifically down regulated the cytosolic β -catenin level, which may play a critical role in the intestinal tumor development.

cPLA₂ is an enzyme which releases AA from cell membrane phospholipids, which is an upstream event in the eicosanoid synthesis pathway involving COX-2.²⁹ COX-2 overexpression is also observed in human colorectal adenocarcinomas and adenomas.³⁰ In




Figure 4. Effect of dietary anthocyanin-rich extract (AE) on (A) intestinal COX-2 and (B) serum levels of prostaglandin E₂ (PGE₂) in *Apc^{Min/+}* mice. The mice were fed 0.2% or 0.5% AE supplemented diets for 7 weeks. After 7 weeks, (A) intestinal COX-2 and (B) serum PGE₂ level were measured. The white bars represent non-supplemented diets, and black bars indicate diets supplemented with AE. Each bar represents the mean \pm SE; $n = 10$ mice for each group. The experiments are repeated 3 times. Statistical significance of the differences was evaluated by *t*-test (* $P < 0.05$). NS, not significant.

this study, although the 0.5% AE diet significantly decreased cPLA₂ mRNA expression, the cPLA₂ activity was not changed.

Meanwhile COX2 mRNA expression in the intestinal mucosa was effectively decreased by 0.5% AE diet compared to the control diet (Fig. 4A). It is reported that AEs inhibited COX-2 mRNA expression in colon cancer model.¹⁰ These results indicate that the inhibitory effects of an AE diet on colon tumor formation may be partially attributed by the inhibition of COX-2. Anthocyanins were also shown to regulate various cellular signaling pathways involving the PLA₂, PI3K/Akt and NF- κ B pathways in several pathological conditions.³¹ Cai et al.⁸ showed that expression of

Akt was reduced in *Apc*^{Min/+} mice received red grape pomace extract. These data suggest that the protective effects of anthocyanins on cancer are not due to the antioxidant activity alone, but also to the direct blockage of signaling pathways. In this study, a tendency towards lower serum concentrations of PGE₂, a metabolite of COX-2, was observed by AE supplementation. We assume that higher amounts of and/or longer periods of AE supplementation may cause a significant reduction in the serum PGE₂ concentration.

The highest content of anthocyanin in AE was cyanidin-3-glycoside (C3G), followed by delphinidin-3-glucoside and petunidin-3-glucoside. In a pharmacokinetic study, C57BL/6J mice received an oral administration of 500 mg C3G/kg body weight and gastrointestinal (GI) mucosa contained 10⁻⁴ to 10⁻³ M C3G which is similar to effective concentrations exerting anti-proliferative effects against cancer cells in vitro.³² The authors showed that these levels were higher than those seen in any other of the tissues. Moreover, the elimination rate constants and half-lives in mucosa were 0.39 h⁻¹ and 1.8 h, respectively. These data indicated that C3G stay in the tissue long enough to exert biological effects. In our previous study, cyanidin and delphinidin exerted significant growth inhibitory effects at ≥1 μM concentrations in human colon cancer cells.¹² These data indicated that the anti-

proliferative effects of these phenolics would be in part responsible for the anticancer activity of AE diet.

Supplementation at levels of 0.2% to 0.5% crude anthocyanin extract corresponds to 2 to 5 g crude anthocyanins per day assuming we consume about 1 kg of food material a day. These levels of supplementation will reach around 0.1 mM to 1 mM in GI mucosa of a 70 kg person since it has been suggested that GI mucosa contained 0.1 mM to 1 mM of cyanidin-3-glucoside when 500 mg/kg body weight of cyanidin-3-glucoside was administered.^{32,33} However, these concentrations might not be accomplishable by consuming black soybeans by diet especially because 100% of anthocyanins are present in soybean coat which is often removed before consumption. Therefore, the use of black soybean coat anthocyanin may be more appropriate as a dietary supplement rather than the use of whole beans. Since we did not analyze all of the constituents present in black soybean coat extract, other bioactive constituents may be present. Black soybean coat also contains several phenolic compounds, isoflavones, and tannins. Since it was reported that above constituents also possess anti-carcinogenic activities against colon cancer,³⁴⁻⁴⁵ anthocyanins might not be the only group of compounds suppressing tumor formation. However, we used the extraction method specifically extract crude anthocyanins from black




Figure 5. The hypothetical scheme of the mechanisms by which an anthocyanin-rich extract (AE) diet inhibits tumor formation in *Apc*^{Min/+} mice. The adenomatous polyposis coli (APC) mutation in *Apc*^{Min/+} mice leads to accumulation of cytosolic and nuclear β -Cat which leads to Wnt-specific target gene translation. Reactive oxygen species (ROS) activate AA release from membrane, leading to increase of inflammatory mediators such as COX-2 and prostaglandin E₂ (PGE₂) which accelerate tumorigenesis. AE diet effectively inhibits cytosolic β -Cat accumulation and suppresses ROS-induced COX-2 and PGE₂ expressions. β -Cat, β -catenin; AA, arachidonic acid; AXIN, Axis inhibition protein; COX-2, cyclooxygenase-2; cPLA₂, cytosolic phospholipase A₂; ERK, extracellular signal-regulated kinase; GSK3 β , glycogen synthase kinase 3 β ; MAPK, mitogen-activated protein kinase; PL, phospholipid.

soybean coat⁴⁶ and we assumed major constituents in the extract are anthocyanins. A study reported by Xu and Chang⁴⁷ indicated that seed coat of black soybean contains less than 10% of phenolic acids and flavonoids including isoflavones compared to dehulled beans. Also, 100% of anthocyanins are present in seed coat. Presuming these phytochemicals possess similar potential to exert anti-carcinogenic activities, anthocyanins might be major anti-carcinogenic compounds in seed coat especially in this crude anthocyanin extract. To elucidate the contribution of each soybean coat components regarding the anticancer activities, a further study will be needed.

In conclusion, dietary AE possesses chemopreventive action in intestinal carcinogenesis by reducing the cytosolic levels of β -catenin, and by suppressing inflammatory responses (Fig. 5). These results may have implications for dietary intervention in human FAP and colorectal cancer. Further studies are required to provide explanations for the precise mechanisms of action.

ACKNOWLEDGMENTS

This work was supported by the Korea Science & Engineering Foundation (R01-2005-000-10602-0 (2006) and by the Basic Science Research Program thorough the National Research Foundation of Korea (NRF), funded by the Ministry of Education, Science and Technology (2010-0001886).

CONFLICTS OF INTEREST

No potential conflicts of interest were disclosed.

REFERENCES

- World Health Organization (WHO), 2013. Cancer Incidence and Mortality Worldwide: IARC. <http://www.who.int/mediacentre/factsheets/fs297/en/>. Accessed June 13, 2015.
- Lizarraga D, Vinardell MP, Noé V, van Delft JH, Alcarraz-Vizán G, van Breda SG, et al. A lyophilized red grape pomace containing proanthocyanidin-rich dietary fiber induces genetic and metabolic alterations in colon mucosa of female C57BL/6J mice. *J Nutr* 2011;141:1597-604.
- Tantamango YM, Knutsen SF, Beeson WL, Fraser G, Sabate J. Foods and food groups associated with the incidence of colorectal polyps: the Adventist Health Study. *Nutr Cancer* 2011;63:565-72.
- Kim HJ, Tsou I, Park JM, Chung JI, Shin SC, Chang KC. Anthocyanins from soybean seed coat inhibit the expression of TNF- α -induced genes associated with ischemia/reperfusion in endothelial cell by NF- κ B-dependent pathway and reduce rat myocardial damages incurred by ischemia and reperfusion in vivo. *FEBS Lett* 2006;580:1391-7.
- Ververidis F, Trantas E, Douglas C, Vollmer G, Kretzschmar G, Panopoulos N. Biotechnology of flavonoids and other phenylpropanoid-derived natural products. Part I: chemical diversity, impacts on plant biology and human health. *Biotechnol J* 2007;2:1214-34.
- Thomasset S, Berry DP, Cai H, West K, Marczylo TH, Marsden D, et al. Pilot study of oral anthocyanins for colorectal cancer chemoprevention. *Cancer Prev Res (Phila)* 2009;2:625-33.
- Lim S, Xu J, Kim J, Chen TY, Su X, Standard J, et al. Role of anthocyanin-enriched purple-fleshed sweet potato p40 in colorectal cancer prevention. *Mol Nutr Food Res* 2013;57:1908-17.
- Cai H, Marczylo TH, Teller N, Brown K, Steward WP, Marko D, et al. Anthocyanin-rich red grape extract impedes adenoma development in the Apc(Min) mouse: pharmacodynamic changes and anthocyanin levels in the murine biophase. *Eur J Cancer* 2010;46:811-7.
- Bobe G, Wang B, Seeram NP, Nair MG, Bourquin LD. Dietary anthocyanin-rich tart cherry extract inhibits intestinal tumorigenesis in APC(Min) mice fed suboptimal levels of sulindac. *J Agric Food Chem* 2006;54:9322-8.
- Lala G, Malik M, Zhao C, He J, Kwon Y, Giusti MM, et al. Anthocyanin-rich extracts inhibit multiple biomarkers of colon cancer in rats. *Nutr Cancer* 2006;54:84-93.
- Harris GK, Gupta A, Nines RG, Kresty LA, Habib SG, Frankel WL, et al. Effects of lyophilized black raspberries on azoxymethane-induced colon cancer and 8-hydroxy-2'-deoxyguanosine levels in the Fischer 344 rat. *Nutr Cancer* 2001;40:125-33.
- Kim JM, Kim JS, Yoo H, Choung MG, Sung MK. Effects of black soybean [*Glycine max* (L.) Merr.] seed coats and its anthocyanidins on colonic inflammation and cell proliferation in vitro and in vivo. *J Agric Food Chem* 2008;56:8427-33.
- Galvano F, La Fauci L, Lazzarino G, Fogliano V, Ritieni A, Ciappellano S, et al. Cyanidins: metabolism and biological properties. *J Nutr Biochem* 2004;15:2-11.
- Balboa MA, Balsinde J. Oxidative stress and arachidonic acid mobilization. *Biochim Biophys Acta* 2006;1761:385-91.
- Nakanishi M, Rosenberg DW. Roles of cPLA2 α and arachidonic acid in cancer. *Biochim Biophys Acta* 2006;1761:1335-43.
- Cummings BS, McHowat J, Schnellmann RG. Phospholipase A(2)s in cell injury and death. *J Pharmacol Exp Ther* 2000;294:793-9.
- Su LK, Kinzler KW, Vogelstein B, Preisinger AC, Moser AR, Luongo C, et al. Multiple intestinal neoplasia caused by a mutation in the murine homolog of the APC gene. *Science* 1992;256:668-70.
- Powell SM, Zilz N, Beazer-Barclay Y, Bryan TM, Hamilton SR, Thibodeau SN, et al. APC mutations occur early during colorectal tumorigenesis. *Nature* 1992;359:235-7.
- Pajari AM, Rajakangas J, Päiväranta E, Kosma VM, Rafter J, Mutanen M. Promotion of intestinal tumor formation by inulin is associated with an accumulation of cytosolic beta-catenin in Min mice. *Int J Cancer* 2003;106:653-60.
- Liu Z, Brooks RS, Ciappio ED, Kim SJ, Crott JW, Bennett G, et al. Diet-induced obesity elevates colonic TNF- α in mice and is accompanied by an activation of Wnt signaling: a mechanism for obesity-associated colorectal cancer. *J Nutr Biochem* 2012;23:1207-13.
- Chiou YS, Tsai ML, Nagabhushanam K, Wang YJ, Wu CH, Ho CT, et al. Pterostilbene is more potent than resveratrol in preventing azoxymethane (AOM)-induced colon tumorigenesis via activation of the NF-E2-related factor 2 (Nrf2)-mediated antioxidant signal-

- ing pathway. *J Agric Food Chem* 2011;59:2725-33.
22. Seidelin JB, Nielsen OH. Continuous cytokine exposure of colonic epithelial cells induces DNA damage. *Eur J Gastroenterol Hepatol* 2005;17:363-9.
 23. Korbecki J, Baranowska-Bosiacka I, Gutowska I, Chlubek D. The effect of reactive oxygen species on the synthesis of prostanoids from arachidonic acid. *J Physiol Pharmacol* 2013;64:409-21.
 24. Hugh TJ, Dillon SA, O'Dowd G, Getty B, Pignatelli M, Poston GJ, et al. beta-catenin expression in primary and metastatic colorectal carcinoma. *Int J Cancer* 1999;82:504-11.
 25. Herter P, Kuhnen C, Müller KM, Wittinghofer A, Müller O. Intracellular distribution of beta-catenin in colorectal adenomas, carcinomas and Peutz-Jeghers polyps. *J Cancer Res Clin Oncol* 1999;125:297-304.
 26. Hao XP, Pretlow TG, Rao JS, Pretlow TP. Beta-catenin expression is altered in human colonic aberrant crypt foci. *Cancer Res* 2001;61:8085-8.
 27. Wang LS, Kuo CT, Cho SJ, Seguin C, Siddiqui J, Stoner K, et al. Black raspberry-derived anthocyanins demethylate tumor suppressor genes through the inhibition of DNMT1 and DNMT3B in colon cancer cells. *Nutr Cancer* 2013;65:118-25.
 28. Lu JN, Lee WS, Yun JW, Kim MJ, Kim HJ, Kim DC, et al. Anthocyanins from *Vitis coignetiae* Pulliat inhibit cancer invasion and epithelial-mesenchymal transition, but these effects can be attenuated by tumor necrosis factor in human uterine cervical cancer hela cells. *Evid Based Complement Alternat Med* 2013; 2013:503043.
 29. D'Orazi G, Sciulli MG, Di Stefano V, Riccioni S, Frattini M, Falcioni R, et al. Homeodomain-interacting protein kinase-2 restrains cytosolic phospholipase A2-dependent prostaglandin E2 generation in human colorectal cancer cells. *Clin Cancer Res* 2006;12:735-41.
 30. Eberhart CE, Coffey RJ, Radhika A, Giardiello FM, Ferrenbach S, DuBois RN. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* 1994;107:1183-8.
 31. Seeram NP, Momin RA, Nair MG, Bourquin LD. Cyclooxygenase inhibitory and antioxidant cyanidin glycosides in cherries and berries. *Phytomedicine* 2001;8:362-9.
 32. Marczylo TH, Cooke D, Brown K, Steward WP, Gescher AJ. Pharmacokinetics and metabolism of the putative cancer chemopreventive agent cyanidin-3-glucoside in mice. *Cancer Chemother Pharmacol* 2009;64:1261-8.
 33. Cooke D, Schwarz M, Boocock D, Winterhalter P, Steward WP, Gescher AJ, et al. Effect of cyanidin-3-glucoside and an anthocyanin mixture from bilberry on adenoma development in the ApcMin mouse model of intestinal carcinogenesis-relationship with tissue anthocyanin levels. *Int J Cancer* 2006;119:2213-20.
 34. Cheah KY, Howarth GS, Bindon KA, Kennedy JA, Bastian SE. Low molecular weight procyanidins from grape seeds enhance the impact of 5-Fluorouracil chemotherapy on Caco-2 human colon cancer cells. *PLoS One* 2014;9:e98921.
 35. Gorlach S, Wagner W, Podsedek A, Szweczyk K, Koziolkiewicz M, Dastych J. Procyanidins from Japanese quince (*Chaenomeles japonica*) fruit induce apoptosis in human colon cancer Caco-2 cells in a degree of polymerization-dependent manner. *Nutr Cancer* 2011;63:1348-60.
 36. Gerhauser C. Cancer chemopreventive potential of apples, apple juice, and apple components. *Planta Med* 2008;74:1608-24.
 37. Kahle K, Huemmer W, Kempf M, Scheppach W, Erk T, Richling E. Polyphenols are intensively metabolized in the human gastrointestinal tract after apple juice consumption. *J Agric Food Chem* 2007;55:10605-14.
 38. Barth SW, Fährndrich C, Bub A, Dietrich H, Watzl B, Will F, et al. Cloudy apple juice decreases DNA damage, hyperproliferation and aberrant crypt foci development in the distal colon of DMH-initiated rats. *Carcinogenesis* 2005;26:1414-21.
 39. Maldonado ME, Bousserouel S, Gossé F, Lobstein A, Raul F. Implication of NF- κ B and p53 in the expression of TRAIL-death receptors and apoptosis by apple procyanidins in human metastatic SW620 cells. *Biomedica* 2010;30:577-86.
 40. Lepri SR, Zanelatto LC, da Silva PB, Sartori D, Ribeiro LR, Mantovani MS. Effects of genistein and daidzein on cell proliferation kinetics in HT29 colon cancer cells: the expression of CTNNBIP1 (β -catenin), APC (adenomatous polyposis coli) and BIRC5 (survivin). *Hum Cell* 2014;27:78-84.
 41. Park HJ, Jeon YK, You DH, Nam MJ. Daidzein causes cytochrome c-mediated apoptosis via the Bcl-2 family in human hepatic cancer cells. *Food Chem Toxicol* 2013;60:542-9.
 42. Min WK, Sung HY, Choi YS. Suppression of colonic aberrant crypt foci by soy isoflavones is dose-independent in dimethylhydrazine-treated rats. *J Med Food* 2010;13:495-502.
 43. Bhagat M, Sharma V, Saxena AK. Anti-proliferative effect of leaf extracts of *Eucalyptus citriodora* against human cancer cells in vitro and in vivo. *Indian J Biochem Biophys* 2012;49:451-7.
 44. Senthilkumar R, Parimelazhagan T, Chaurasia OP, Srivastava RB. Free radical scavenging property and antiproliferative activity of *Rhodiola imbricata* Edgew extracts in HT-29 human colon cancer cells. *Asian Pac J Trop Med* 2013;6:11-9.
 45. Salimi M, Majd A, Sepahdar Z, Azadmanesh K, Irian S, Ardestaniyan MH, et al. Cytotoxicity effects of various *Juglans regia* (walnut) leaf extracts in human cancer cell lines. *Pharm Biol* 2012;50:1416-22.
 46. Choung MG, Baek IY, Kang ST, Han WY, Shin DC, Moon HP, et al. Isolation and determination of anthocyanins in seed coats of black soybean (*Glycine max* (L.) Merr.). *J Agric Food Chem* 2001;49:5848-51.
 47. Xu B, Chang SK. Antioxidant capacity of seed coat, dehulled bean, and whole black soybeans in relation to their distributions of total phenolics, phenolic acids, anthocyanins, and isoflavones. *J Agric Food Chem* 2008;56:8365-73.