

Benefits of Emergency Departments' Contribution to Stroke Prophylaxis in Atrial Fibrillation

The EMERG-AF Study (Emergency Department Stroke Prophylaxis and Guidelines Implementation in Atrial Fibrillation)

Blanca Coll-Vinent, MD, PhD; Alfonso Martín, MD, PhD; Juan Sánchez, MD, PhD; Juan Tamargo, MD, PhD; Coral Suero, MD; Francisco Malagón, MD; Mercedes Varona, MD; Manuel Cancio, MD; Susana Sánchez, MD, PhD; José Carbajosa, MD; José Ríos, BSc, MSc; Georgina Casanovas, BSc; Carles Ràfols, MD; Carmen del Arco, MD, PhD; on behalf of the EMERG-AF Investigators*

Background and Purpose—Long-term benefits of initiating stroke prophylaxis in the emergency department (ED) are unknown. We analyzed the long-term safety and benefits of ED prescription of anticoagulation in atrial fibrillation patients.

Methods—Prospective, multicenter, observational cohort of consecutive atrial fibrillation patients was performed in 62 Spanish EDs. Clinical variables and thromboprophylaxis prescribed at discharge were collected at inclusion. Follow-up at 1 year post-discharge included data about thromboprophylaxis and its complications, major bleeding, and death; risk was assessed with univariate and bivariate logistic regression models.

Results—We enrolled 1162 patients, 1024 (88.1%) at high risk according to CHA₂DS₂-VASc score. At ED discharge, 935 patients (80.5%) were receiving anticoagulant therapy, de novo in 237 patients (55.2% of 429 not previously treated). At 1 year, 48 (4.1%) patients presented major bleeding events, and 151 (12.9%) had died. Anticoagulation first prescribed in the ED was not related to major bleeding (hazard ratio, 0.976; 95% confidence interval, 0.294–3.236) and was associated with a decrease in mortality (hazard ratio, 0.398; 95% confidence interval, 0.231–0.686). Adjusting by the main clinical and sociodemographic characteristics, concomitant antiplatelet treatment, or destination (discharge or admission) did not affect the results.

Conclusions—Prescription of anticoagulation in the ED does not increase bleeding risk in atrial fibrillation patients at high risk of stroke and contributes to decreased mortality. (*Stroke*. 2017;48:1344-1352. DOI: 10.1161/STROKEAHA.116.014855.)

Key Words: anticoagulants ■ atrial fibrillation ■ hemorrhage ■ mortality stroke

Atrial fibrillation (AF) is the most prevalent arrhythmia treated in emergency departments (EDs),^{1–3} and the most serious complication is stroke or systemic embolism.⁴ The high effectiveness of oral anticoagulants in preventing these complications⁵ has produced general agreement that stroke prophylaxis is the mainstay of AF management in all health-care settings.^{6–8}

Given the number of these patients attended in EDs who are at high risk of stroke but not receiving anticoagulants,^{3,9,10} the ED may be an underused resource for starting anticoagulation,

although this remains controversial. Some authors have postulated that ED physicians should only inform patients about the need for stroke prophylaxis, and oral anticoagulation should be prescribed in other settings.¹¹ On the other hand, as substantial underuse has been reported in them,^{12,13} numerous authors have called for coordinated efforts, including ED involvement, to improve treatment.¹⁴

No study to date has prospectively addressed the long-term outcomes of ED prescription and its contribution to stroke prophylaxis. Analysis of the feasibility, benefits, and

Received July 21, 2016; final revision received February 6, 2017; accepted February 9, 2017.

From the Arrhythmia Division, Spanish Society of Emergency Medicine (SEMES), Madrid, Spain (B.C.-V., A.M., J.S., C.S., F.M., M.V., M.C., S.S., J.C., C.d.A.); Atrial Fibrillation Unit (UFA) (B.C.-V.), and Department of Pharmacology, CIBERCV, Universidad Complutense, Madrid, Spain (J.T.); Laboratory of Biostatistics and Epidemiology, Universitat Autònoma de Barcelona; Medical Statistics Core Facility, IDIBAPS (Hospital Clinic), Barcelona, Spain (J.R., G.C.); and Medical Department, Bayer Hispania-sl, Barcelona, Spain (C.R.).

*A list of all EMERG-AF Investigators is given in the Appendix.

The online-only Data Supplement is available with this article at <http://stroke.ahajournals.org/lookup/suppl/doi:10.1161/STROKEAHA.116.014855/-/DC1>.

Correspondence to Blanca Coll-Vinent, MD, PhD, Emergency Department, Hospital Clínic de Barcelona, Carrer de Villarroel, 170, 08036 Barcelona, Spain. E-mail bcvinent@clinic.cat

© 2017 The Authors. *Stroke* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution Non-Commercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.116.014855

long-term safety of ED prescription of anticoagulants could contribute to categorical recommendations for stroke prophylaxis in this setting.

The objectives of the EMERG-AF study (Emergency Department Stroke Prophylaxis and Guidelines Implementation in Atrial Fibrillation) were to analyze thromboprophylaxis prescription in patients with AF attended in the ED and long-term outcomes: major bleeding, stroke or systemic embolism, and death.

Methods

Study Design and Setting

The EMERG-AF was a prospective, multicenter, observational cohort study performed in 62 Spanish EDs from April 2013 to June 2014. Data were collected prospectively at the first ED visit (April 1–30, 2013) on study inclusion and at a 1-year (± 4 weeks) follow-up visit.

To guarantee a hospital sample representative of the Spanish healthcare system, study centers were selected from the national catalogue published by Spain's Ministry of Health and Social Policy. Centers were categorized by size, level of services, and location; the scientific committee recruited 25% of the eligible centers, weighted by category, to participate. The study was approved by the Scientific Ethics Committee of each participating hospital.

Selection of Participants

We included consecutive patients older than 18 years attended in the ED for any reason during the study period with AF documented in clinical records or demonstrated in an ECG obtained when the treating ED physician considered it necessary during clinical evaluation. Exclusion criteria were atrial flutter and clinical trial participation.

A total of 1138 patients were considered necessary for a representative sample, assuming an estimated anticoagulation ratio of 60%, 2-sided 95% confidence interval (CI), $\pm 3\%$ sensitivity and 10% losses to follow-up.

Informed consent was obtained in all cases. To guarantee confidentiality, access to patient identity was restricted to the treating ED physician. The central database contained identification numbers for individual study participants and was password protected.

Initial Data Collection and Processing

Copies of the EMERG-AF study protocol (including definitions and codes) were provided to all collaborating researchers, who were trained by the lead investigator in data recording. The information was collected in a centralized electronic file after interviewing patients or their relatives and later checked and completed by chart review. The study's scientific committee made no therapeutic recommendations and provided no specific instructions about AF and its management during the data collection.

The following information was included on the initial data collection sheet: demographic data, comorbidities, disability (according to the Katz scale),¹⁵ type of AF, duration of the current AF episode, risk factors for stroke (CHA₂DS₂-VASc score),¹⁶ bleeding risk (according to the HAS-BLED score [Hypertension, Abnormal Renal and Liver Function, Stroke, Bleeding, Labile INRs, Elderly, Drugs or Alcohol]),¹⁷ current stroke prophylaxis (anticoagulation, antiplatelet therapy, or both), symptoms that caused ED consultation, clinical presentation, and ED evaluation, patients' final outcome (discharge, admission, or death), and stroke prophylaxis prescribed in the ED.

If anticoagulation was not prescribed, the treating physician was asked to select a reason from a set of multiple-choice possibilities. We included the most frequent causes of a lack of anticoagulant prescription, as stated in daily practice studies,^{2,3,10} with the aim of capturing the main reasons for this behavior (Table I in the [online-only Data Supplement](#)).

Major bleeding was defined as potential risk of death because of bleed location (ie, intracranial, intraspinal, pericardial), potential risk

of severe sequelae (intraocular, intramuscular with compartment syndrome), or required admission, surgery, or blood transfusion.¹⁸

Follow-Up

Follow-up in both groups included a phone call and a review of each patient's electronic medical record at 1 year (± 4 weeks) postinclusion to address long-term outcomes. The follow-up analysis included the following variables: bleeding events, systemic embolism and stroke events, stroke prophylaxis at the time of follow-up, and mortality (date and cause of death). If bleeding or a stroke or systemic embolism had occurred, type of bleeding (minor or major) or type of complication, respectively, were recorded along with date of presentation, location, and outcome.

The study's scientific committee independently reviewed all data-sheets to detect inconsistencies. As needed, a query was made to the principal investigator at the corresponding center to resolve any questions.

Outcome Measures

As the main objective of the present article was patient safety at 1 year post-discharge, incidence of major bleeding was the primary outcome. Mortality and stroke/systemic embolism were secondary outcomes.

Statistical Analysis

All analysis was performed with SPSS 20.0 and SAS statistical software packages, accepting a risk threshold for a 2-sided Type I error of 5%. Differences in the profiles of patients with or without anticoagulation treatment at discharge were assessed by Fisher exact test or *t* test for independent groups for quantitative or qualitative variables, respectively.

Univariate logistic regression models were built to assess the risk of major bleeding or death. The models estimated the hazard ratio (HR) and 95% CI using the most relevant clinical and demographic factors: age, sex, hypertension, diabetes mellitus, renal failure, acute heart failure, structural heart disease, cerebrovascular disease, previous AF, CHA₂DS₂-VASc, HAS-BLED, concomitant antiplatelet treatment, previous major bleeding, and relationship between the reason for ED consultation and AF (yes/no), and destination. Because the prescription of anticoagulant therapy was the main variable of interest, we conducted a number of multivariate analyses, adjusting by these factors to assess its relevance as a potential independent predictor for hemorrhagic events or death. All analyses were also performed for patients without previous stroke prophylaxis treatment.

Results

Characteristics of Study Participants

During the study period, 1162 patients were included (mean age, 74.7 years [SD 11.2], 590 [50.8%] women). In 178 patients (15.3%), de novo AF diagnosis was made during the ED inclusion visit. Comorbid conditions were common. The majority (88.1%) of the patients included were at high risk of stroke according to the CHA₂DS₂-VASc score. Demographic and clinical characteristics of all patients and of patients without previous anticoagulation are listed in Table 1. Two patients (0.2%) died during the ED stay. After evaluation, 512 patients (44.0%) were admitted, and 648 patients (55.8%) were discharged home.

Main Results

Anticoagulation Prescription

At the time of the ED visit, 733 patients (63.1%) were taking anticoagulants. At ED discharge, 935 patients (80.5%), including those diagnosed de novo, were receiving antithrombotic therapy with anticoagulants (146 with concomitant antiplatelets).

Table 2 details ED anticoagulation prescription according to stroke risk. Considering both previous and de novo anticoagulant therapies, 166 high-risk patients according to the CHA₂DS₂-VASc score (14.3%) did not receive anticoagulation therapy at discharge from the ED. The main reasons given by the treating physicians for not prescribing anticoagulation in these high-risk patients were high risk of bleeding (81 patients [48.8%]) and no indication for anticoagulation (56 patients [33.7%]).

Of the 935 patients anticoagulated at ED discharge, 764 (81.7%) received vitamin K antagonists, 97 (10.4%)

low-molecular-weight heparin, 73 (7.8%) direct oral anticoagulants, and 1 patient (0.1%) other anticoagulation. Prescriptions were de novo in 237 patients (55.2% of the 429 not previously anticoagulated). At discharge, all participants were referred to another healthcare facility for follow-up and long-term monitoring.

Follow-Up

After 1 year, 58 patients (4.9%) were deceased, and 52 patients (4.4%) were lost to follow-up. The reasons for losses to follow-up

Table 1. Demographic and Clinical Characteristics of All Patients and Patients Without Previous Anticoagulation

Variable	Total, n=1162, n (%)	Not Previously Anticoagulated			P Value
		Total, n=429, n (%)	Not Anticoagulated at Discharge, (n=192/429), n (%)	Anticoagulated at Discharge, (n=237/429), n (%)	
Age, mean (SD)	74.7±11.2	72.4±13.0	71.5±15.0	73.1±11.0	0.22
Sex (female)	590 (50.8)	204 (47.6)	95 (49.5)	109 (46.0)	0.47
Structural heart disease	585 (50.3)	129 (30.1)	56 (29.2)	73 (30.8)	0.71
Acute heart failure	406 (34.9)	87 (20.3)	33 (17.2)	54 (22.8)	0.18
Left ventricular dysfunction	209 (18)	41 (9.6)	17 (8.9)	24 (10.1)	0.65
Hypertension	868 (74.7)	286 (66.7)	113 (58.9)	173 (73.0)	0.002
Previous stroke	172 (14.8)	47 (11.0)	21 (11.0)	26 (11.0)	1.00
Previous systemic embolism	19 (1.6)	1 (0.2)	1 (0.5)	0 (0.0)	0.26
Diabetes mellitus	316 (27.2)	92 (21.5)	34 (17.8)	58 (22.5)	0.09
Renal failure	191 (16.4)	55 (12.8)	25 (13.1)	30 (12.6)	0.91
CHA ₂ DS ₂ -VASc					
0	49 (4.2)	39 (9.0)	28 (14.6)	11 (4.6)	<0.001
1	89 (7.7)	55 (12.8)	31 (16.1)	24 (10.1)	
≥2	1024 (88.1)	335 (78.1)	133 (69.3)	202 (85.2)	
HAS-BLED					
<3	568 (48.9)	242 (56.5)	99 (51.8)	143 (60.3)	0.08
≥3	594 (51.1)	187 (43.6)	93 (48.4)	94 (39.7)	
Disability	126 (10.8)	44 (10.0)	24 (12.0)	20 (8.4)	0.17
Total	8 (0.7)	3 (0.7)	1 (0.5)	2 (0.8)	0.44
Types of AF					
First episode	178 (15.3)	164 (38.1)	49 (25.1)	115 (48.5)	<0.001
Paroxysmal	312 (26.9)	154 (36.0)	81 (42.4)	73 (30.8)	
Persistent	131 (11.3)	44 (10.3)	15 (7.8)	29 (12.2)	
Permanent	541 (46.6)	67 (15.6)	47 (24.6)	20 (8.4)	
Main complaint					
Related to AF*	681 (58.6)	318 (74.1)	111 (57.8)	207 (87.3)	<0.001
Other	481 (41.4)	111 (25.9)	81 (42.2)	30 (12.7)	
Duration of episode					
<48 h	210 (18.1)	141 (32.7)	73 (37.7)	68 (28.7)	0.010
>48 h	539 (46.4)	101 (23.6)	51 (26.7)	50 (21.1)	
Unknown	413 (35.5)	187 (43.7)	68 (35.6)	119 (50.2)	

AF indicates atrial fibrillation.

*Related to AF or its treatment.

Table 2. Anticoagulation at Discharge From the Emergency Department According to Stroke Risk, Previous Treatment, Anticoagulation at Discharge and Final Outcome

CHA ₂ DS ₂ -VASc	Previous Anticoagulation, n (%)		Anticoagulation at Discharge From ED, n (%)		Admitted, n (%)	Discharged, n (%)
<2, n=138	Yes	44 (31.9)	Yes	42 (95.5)	16 (38.1)	26 (61.9)
			No	2 (4.5)	0 (0.0)	2 (100)
	No	94 (68.1)	Yes	35/94 (37.2)*	8 (22.9)	27 (77.1)
			No	59 (62.8)	6 (10.2)	53 (89.8)
≥2, n=1024	Yes	688 (67.3)	Yes	656 (95.3)	298 (45.4)	358 (54.6)
			No	33 (4.7)	27 (84.4)	5 (15.6)
	No	334 (32.7)	Yes	202 (60.5)	92 (45.5)	110 (54.5)
			No	133 (39.5)	65 (49.2)	67 (50.8)

ED indicates emergency department.

*Eighteen of these patients were anticoagulated for subsequent elective cardioversion after ED discharge.

were change of address or other factors making contact impossible (49 patients) and decision to withdraw (3 patients). One-year follow-up was completed by 956 patients (94.6% of survivors and 82.3% of all participants). At the end of the year, 699 (73.1%) patients were taking anticoagulants; the great majority of them (670 patients, 95.8%) received de novo or continued anticoagulant treatment at ED discharge. Anticoagulation persisted in 81.2% of the patients anticoagulated de novo who survived and completed follow-up. The 29 patients prescribed an anticoagulant during follow-up constituted 12.8% of the 227 patients discharged without anticoagulation.

Primary Outcome: Safety

At the end of 1 year, 184 bleeding events had occurred in 150 patients (12.9%), including 55 (29.8%) major bleeding episodes that occurred in 48 patients (4.1%), with 6 fatalities. The main bleeding sites were gastrointestinal (23.9%), otorhinolaryngologic (22.7%), and urogenital (14.1%). Twelve patients had intracranial hemorrhage (6.5% of all bleeding events).

Anticoagulant treatment was not related to major bleeding (HR, 1.376; 95% CI, 0.581–3.260; Figure 1A). These results were similar when only patients who were not receiving anticoagulation before the index visit were taken into account (HR, 0.976; 95% CI, 0.294–3.236; Figure 1B), even when adjusted for possible confounders: final outcome in the first visit (discharge versus admission), antiplatelet treatment, or other relevant clinical or sociodemographic characteristics (Table 3). A separate analysis of discharged, not previously anticoagulated patients also found no significant relationship between starting anticoagulant treatment and major bleeding (HR, 2.597; 95% CI, 0.270–24.966).

Secondary Outcomes

Mortality: During the 1-year follow-up period, 151 patients (12.8%) died, 62 (41.6%) of them because of cardiovascular causes. Anticoagulant treatment was associated with decreased mortality (HR, 0.505; 95% CI, 0.357–0.715; Figure 2A). This association also persisted when only patients who were not receiving anticoagulation before the index visit were taken into account (HR, 0.398; 95% CI, 0.231–0.686; Figure 2B), even when adjusted for the final outcome in the first visit (discharge versus admission), antiplatelet treatment,

or any relevant clinical or sociodemographic characteristic (Table 4). When a separate analysis of the 257 discharged, not previously anticoagulated patients was performed, anticoagulated patients (n=137) had lower mortality than those discharged without anticoagulation (n=120) but the difference (5.8% versus 9.1% mortality, respectively) was nonsignificant (HR, 0.668; 95% CI, 0.264–1.694).

Stroke and systemic embolism: Thromboembolic complications were present in 31 patients (2.7%): 13 strokes, 8 transient ischemic attacks, and 13 systemic embolisms. Four of these patients died, and 15 were left with sequelae.

Anticoagulation was not associated with the presence of stroke events in the univariate analysis (HR, 0.929; 95% CI, 0.381–2.264). When only patients not previously anticoagulated were considered, those who received anticoagulants in the ED visit tended to have a lower incidence of embolic events, although this was not significant (2.1% versus 3.2%; HR, 0.614; 95% CI, 0.187–2.011), even when adjusted for final outcome, antiplatelet treatment, or any relevant clinical or sociodemographic characteristic (Table II in the [online-only Data Supplement](#)). A separate analysis of discharged, not previously anticoagulated patients also found a nonsignificantly lower incidence of embolic complications (1.4% versus 2.5%; HR, 0.556; 95% CI, 0.093–3.330).

Limitations

Participation in an investigative study may have had an impact on the clinicians. To minimize this limitation, risk stratification scores were not made available in the electronic record. The large number of researchers involved could have generated variability in interpreting the questions on the data collection sheet. To avoid this bias, multiple-choice responses were specified. In addition, verbal information obtained from patients at 1-year follow-up could have been affected by recall bias. However, all serious complications would likely have been attended in a hospital and reflected in the official records reviewed to ensure accuracy of outcome information.

The EMERG-AF study was not designed to compare differences in effectiveness between stroke prophylaxis prescribed in EDs and that prescribed in other clinical settings. No such conclusions should be inferred from the study results.




Figure 1. Kaplan–Meier curves showing time to first major bleeding according to anticoagulant treatment. **A**, All patients; **B**, patients who were not receiving oral anticoagulation when they arrived to the emergency department.

Finally, our results can be generalized to the Spanish National Health System and other closely related systems. Further studies are needed to confirm the external validity of these results.

Discussion

The EMERG-AF is the first study to date to prospectively analyze the long-term effects of the ED physician’s contribution to stroke prophylaxis in AF patients, both in prescribing

anticoagulation de novo in high-risk patients and in reinforcing the therapeutic message and encouraging long-term adherence in those patients already taking anticoagulants. In this representative sample of EDs in Spain, the clinical features and stroke risk of the patients included were comparable to those in the global AF population, especially those attended in EDs, as described in the largest studies to date.^{2,3,19} Therefore, the results obtained may be widely applicable to ED daily practice.

Table 3. Factors Associated With Major Bleeding in Patients Not Previously Anticoagulated

Variable	Nonbleeding Patients, n=418, n (%)	Bleeding Patients, n=11, n (%)	Univariate		Multivariate (Anticoagulation at Discharge, Adjusted by Each Variable)	
			HR	95% CI	HR	95% CI
Anticoagulation at discharge	231 (55.26)	6 (54.55)	0.976	0.294–3.236	Nonestimable	
Destination (discharged home)	253 (60.53)	4 (36.36)	0.369	0.107–1.277	0.868	0.261–2.882
Age ≥ 75 y	203 (48.56)	8 (72.73)	2.915	0.769–11.050	0.908	0.275–2.996
Sex (female)	198 (47.37)	6 (54.55)	1.286	0.389–4.244	0.997	0.298–3.339
Hypertension	280 (66.99)	6 (54.55)	0.701	0.210–2.343	1.013	0.305–3.368
Diabetes mellitus	89 (21.29)	3 (27.27)	1.554	0.402–6.009	0.936	0.280–3.132
Renal failure	50 (11.96)	5 (45.45)	6.020	1.794–20.197	0.814	0.245–2.704
Acute heart failure	84 (20.10)	3 (27.27)	1.462	0.385–5.553	0.913	0.270–3.082
Structural heart disease	121 (28.95)	8 (72.73)	6.055	1.585–23.131	0.872	0.265–2.874
Cerebrovascular disease	44 (10.53)	3 (27.27)	2.703	0.694–10.520	0.899	0.273–2.960
Disability	42 (10.05)	2 (18.18)	2.580	0.547–12.157	1.028	0.309–3.424
Consultation related to AF	314 (75.15)	4 (36.36)	7.231	1.869–27.973	2.078	0.592–7.293
Previous AF	257 (61.48)	8 (72.73)	1.699	0.450–6.410	1.064	0.318–3.557
CHA ₂ DS ₂ -VASc score of ≥ 2	326 (77.99)	9 (81.82)	1.432	0.308–6.647	0.903	0.264–3.086
HAS-BLED score of ≥ 3	179 (42.82)	8 (72.73)	3.931	1.041–14.841	0.939	0.285–3.086
Concomitant antiplatelet treatment	179 (42.82)	4 (36.36)	0.723	0.209–2.506	0.822	0.220–3.069
Previous major bleeding	67 (16.03)	3 (27.27)	2.513	0.649–9.729	1.103	0.327–3.724

Definitions detailed in the Statistical Analysis. AF indicates atrial fibrillation; CI, confidence interval; and HR, hazard ratio.

This study yielded 3 main conclusions. First, prescribing anticoagulants in the ED to AF patients at a high risk of stroke is not only feasible but also beneficial in the long term, with a favorable safety profile. Anticoagulation was prescribed by the ED physicians without requiring other medical assessments, and instructions for therapy and referral to other healthcare settings were provided exclusively by ED physicians and nurses. In addition, the prescription followed current guidelines in the vast majority of cases. Nevertheless, and despite the high level of anticoagulation found in our series, almost 15% of eligible patients were not taking anticoagulants at discharge. This highlights the need for an additional effort to provide this prescription to all high-risk patients.

Second, the safety profile of anticoagulation prescribed in EDs, following the guidelines' recommendations and providing systematic referral for follow-up at discharge, is at least as safe in the long term as are prescriptions made in other healthcare settings.^{6-9,20} After adjusting for clinical and sociodemographic variables, the major bleeding rate in patients anticoagulated de novo at the ED was similar to that in nonanticoagulated patients. Moreover, these good results reflect not only an adequate indication for anticoagulation, but also appropriate referrals to INR monitoring. Together with the finding that the majority of patients in whom anticoagulation was prescribed de novo at the ED were adhering to treatment at 1-year follow-up, these results strongly support an assertion that accurate prescription of anticoagulants is feasible in the daily ED practice and suggest that decisions made in the ED influence longitudinal care, as recently reported by Atzema et al.²¹

The third main result of our study was that the good patient safety outcomes were not achieved at the expense of effectiveness. Although there were no significant differences in the benefits of anticoagulant prescription in the ED with respect to embolic events, likely because of the overall paucity of such events, the impact on mortality was overwhelmingly positive. Mortality decreased overall and in patients without previous anticoagulation, even after adjusting for all confounding variables. When a separate analysis was performed on discharged, not previously anticoagulated patients, although the mortality was lower in anticoagulated patients, the difference was not significant. The lack of significance in this subpopulation is likely because of the lower global mortality that would be expected in discharged patients. Although the study was not sufficiently powered for the analysis of subpopulations, this does not lessen the implications of the results because the aim was to analyze the long-term effects of the ED physician's contribution to overall stroke prophylaxis in AF patients. Moreover, although the management of patients with AF is guided by a common protocol and is highly consistent among EDs in Spain, the management of admissions varies greatly between the participating hospitals; thus, many of the included patients were admitted to an Observation Unit or Short Stay Unit, in most cases headed by a physician from the ED staff.

It can be argued that these long-term good results could also be attributed to patient management by other healthcare specialists after ED discharge. We certainly agree with this because the objectives of the EMERG-AF study were to analyze the contribution of ED physicians to the multidisciplinary team approach to stroke prophylaxis in patients with AF,⁶ not to assess in isolation the benefits of prescribing anticoagulants




Figure 2. Kaplan–Meier curves showing mortality according to anticoagulant treatment. **A**, All patients; **B**, patients who were not receiving oral anticoagulation when they arrived to the emergency department.

in the acute setting. Indeed, as a general rule and as one of the pillars of starting anticoagulation, all AF patients discharged from EDs in our setting are referred to another level of health care (internal medicine, cardiology, neurology, geriatrics, and primary care) for follow-up and to specialized clinics or primary care to monitor anticoagulation and reinforce patient education. Thus, our results are an accurate reflection of the potential benefits of a coordinated and multidisciplinary

strategy of stroke prophylaxis involving ED physicians and nurses, and specifically highlight the potential ED contribution to maximize appropriate prophylaxis in these patients.^{22–24}

Conclusions

Prescription in the ED of anticoagulation in AF patients at a high risk of stroke is not only feasible but also beneficial in the long term to decrease mortality. Therefore, the active

Table 4. Factors Associated With 1-Year Mortality in Patients Not Previously Anticoagulated

Variable	Alive at 1 y n=371, n(%)	Deceased, at 1 y, n=58, n(%)	Univariate		Multivariate, (Anticoagulation at Discharge Adjusted by Each Variable)	
			HR	95% CI	HR	95% CI
Anticoagulation at discharge	217 (58.49)	20 (34.48)	0.398	0.231–0.686	Nonestimable	
Destination (discharged home)	238 (64.15)	19 (32.76)	0.284	0.162–0.498	0.363	0.209–0.628
Age ≥75 y	164 (44.20)	47 (81.03)	4.839	2.506–9.343	0.380	0.220–0.655
Sex (female)	169 (45.55)	35 (60.34)	1.714	1.009	0.401	0.233–0.691
Hypertension	241 (64.96)	45 (77.59)	1.724	0.929–3.201	0.365	0.211–0.632
Diabetes mellitus	75 (20.22)	17 (29.31)	1.578	0.895–2.784	0.376	0.217–0.650
Renal failure	37 (9.97)	18 (31.03)	3.634	2.078–6.354	0.399	0.232–0.688
Acute heart failure	66 (17.79)	21 (36.21)	2.253	1.307–3.881	0.370	0.214–0.638
Structural heart disease	100 (26.95)	29 (50.00)	2.364	1.406–3.974	0.388	0.225–0.669
Cerebrovascular disease	34 (9.16)	13 (22.41)	2.440	1.314–4.530	0.401	0.233–0.691
Disability	26 (7.01)	18 (31.03)	4.622	2.643–8.085	0.447	0.258–0.773
Consultation related to AF	287 (77.36)	31 (53.45)	2.749	1.634–4.624	0.519	0.289–0.931
Previous AF	221 (59.57)	44 (75.86)	1.973	1.079–3.606	0.443	0.252–0.775
CHA ₂ DS ₂ -VASc score of ≥2	282 (76.01)	53 (91.38)	3.082	1.231–7.715	0.337	0.195–0.583
HAS-BLED score of ≥3	142 (38.27)	45 (77.59)	4.898	2.638–9.094	0.435	0.253–0.751
Concomitant antiplatelet treatment	217 (58.49)	20 (34.48)	1.368	0.814–2.300	0.395	0.221–0.706
Previous major bleeding	48 (12.94)	22 (37.93)	3.390	1.979–5.810	0.468	0.269–0.815

Definitions detailed in the text. AF indicates atrial fibrillation; CI, confidence interval; and HR, hazard ratio.

involvement of ED physicians could help to improve the global results of stroke prophylaxis in AF and thus help to improve the prognosis and quality of life of this increasing population of patients attended in the acute setting.

Appendix

María Agud; José Aguilar; Alfons Aguirre; Amparo de Simón Almela; Mercè Almirall; Oscar Álvarez; Luis Amador; Juan Antonio Andueza; Francisco José Aramburu; Ignacio Ayala; Ángel Bajo; Carlos Bilbaíno; Ricardo Calvo; María Elena Díaz; María Jesús Estévez; Cristina Flaño; Carolina Fuenzalida; Cristina Garcés; María Teresa García; Pedro García; Luis García-Castrillo; Carmen Gargallo; José Manuel Garrido; Juan González; Pablo Herrero; Eduardo Jiménez; Gregorio Jiménez; José Lázaro; Pedro Lopetegui; Ignacio López; José María Lubillo; María José Marchena; Francisco Moya; Julián Mozota; Francisco José Navarro; Xavier Palom; Javier Ochoa; Ana María Peiró; Juan Manuel Parra; Álvaro Perea; Pascual Piñera; Pere Riambau; Fernando Richard; Belén Rodríguez; Roberto Rodríguez; Eva Ruiz; Francisco Ruiz; Ana María Segarra; Carmen Seijas; Javier Sesma; Wilfredo Soler; María del Mar Sousa; José Manuel Torres; Olga María Trejo; José Vicente.

Acknowledgments

We thank Elaine Lilly, PhD, for her careful English language revision of the article.

Sources of Funding

The study was supported by a grant from Bayer Hispania, which had no influence on the content of the article.

Disclosures

Dr Ràfols is an employee of Bayer Hispania SL. Dr Suero has perceived modest honoraria from Bayer, Boehringer-Ingelheim, and

Bristol Myers-Squibb for specific speech and advisory board. The other authors report no conflicts.

References

- McDonald AJ, Pelletier AJ, Ellinor PT, Camargo CA Jr. Increasing US emergency department visit rates and subsequent hospital admissions for atrial fibrillation from 1993 to 2004. *Ann Emerg Med.* 2008;51:58–65. doi: 10.1016/j.annemergmed.2007.03.007.
- del Arco C, Martín A, Laguna P, Gargantilla P; Investigators in the Spanish Atrial Fibrillation in Emergency Medicine Study Group (GEFAUR). Analysis of current management of atrial fibrillation in the acute setting: GEFAUR-1 study. *Ann Emerg Med.* 2005;46:424–430. doi: 10.1016/j.annemergmed.2005.03.002.
- Coll-Vinent B, Martín A, Malagón F, Suero C, Sánchez J, Varona M, et al; HERMES-AF Investigators; HERMES-AF Investigators. Stroke prophylaxis in atrial fibrillation: searching for management improvement opportunities in the emergency department: the HERMES-AF study. *Ann Emerg Med.* 2015;65:1–12. doi: 10.1016/j.annemergmed.2014.07.016.
- Hughes M, Lip GY; Guideline Development Group, National Clinical Guideline for Management of Atrial Fibrillation in Primary and Secondary Care, National Institute for Health and Clinical Excellence. Stroke and thromboembolism in atrial fibrillation: a systematic review of stroke risk factors, risk stratification schema and cost effectiveness data. *Thromb Haemost.* 2008;99:295–304. doi: 10.1160/TH07-08-0508.
- Hart RG, Pearce LA, Aguilar MI. Meta-analysis: antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med.* 2007;146:857–867.
- Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Europace.* 2016;18:1609–1678. doi: 10.1093/europace/euw295.
- January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, et al; ACC/AHA Task Force Members. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task

- Force on practice guidelines and the Heart Rhythm Society. *Circulation*. 2014;130:e199–e267. doi: 10.1161/CIR.0000000000000041.
8. Martín A, Fernández-Lozano I, Coll-Vinent B, Tercedor L, del Arco C, Arribas F, et al. Management of patients with atrial fibrillation in hospital emergency departments (update 2012). A consensus document of the arrhythmia division of the Spanish society of emergency medicine and the electrophysiology and arrhythmia division of the Spanish Society of Cardiology. *Emergencias*. 2012;24:300–324.
 9. Scheuermeyer FX, Innes G, Pourvali R, Dewitt C, Grafstein E, Heslop C, et al. Missed opportunities for appropriate anticoagulation among emergency department patients with uncomplicated atrial fibrillation or flutter. *Ann Emerg Med*. 2013;62:557.e2–565.e2. doi: 10.1016/j.annemergmed.2013.04.004.
 10. Laguna P, Martín A, del Arco C, Gargantilla P; Investigators in the Spanish Atrial Fibrillation in Emergency Medicine Study Group (GEFAUR). Risk factors for stroke and thromboprophylaxis in atrial fibrillation: what happens in daily clinical practice? The GEFAUR-1 study. *Ann Emerg Med*. 2004;44:3–11. doi: 10.1016/S0196064404000587.
 11. Barrett TW, Marill KA. Anticoagulation for emergency department patients with atrial fibrillation: is our duty to inform or prescribe? *Ann Emerg Med*. 2013;62:566–568. doi: 10.1016/j.annemergmed.2013.05.027.
 12. Shantsila E, Wolff A, Lip GY, Lane DA. Optimising stroke prevention in patients with atrial fibrillation: application of the GRASP-AF audit tool in a UK general practice cohort. *Br J Gen Pract*. 2015;65:e16–e23. doi: 10.3399/bjgp15X683113.
 13. Blanch P, Freixa R, Ibernón M, Delso J, Salas E, Sobrepera JL, et al. [Use of oral anticoagulants in patients discharged with atrial fibrillation in 2000]. *Rev Esp Cardiol*. 2003;56:1057–1063.
 14. Scott PA, Pancioli AM, Davis LA, Frederiksen SM, Eckman J. Prevalence of atrial fibrillation and antithrombotic prophylaxis in emergency department patients. *Stroke*. 2002;33:2664–2669.
 15. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of illness in the aged. the index of ADL: a standardized measure of biological and psychosocial function. *JAMA*. 1963;185:914–919.
 16. Lip GY, Nieuwlaat R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the Euro Heart Survey on atrial fibrillation. *Chest*. 2010;137:263–272. doi: 10.1378/chest.09-1584.
 17. Pisters R, Lane DA, Nieuwlaat R, de Vos CB, Crijns HJ, Lip GY. A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: the Euro Heart Survey. *Chest*. 2010;138:1093–1100. doi: 10.1378/chest.10-0134.
 18. Schulman S, Angerås U, Bergqvist D, Eriksson B, Lassen MR, Fisher W; Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in surgical patients. *J Thromb Haemost*. 2010;8:202–204. doi: 10.1111/j.1538-7836.2009.03678.x.
 19. Oldgren J, Healey JS, Ezekowitz M, Commerford P, Avezum A, Pais P, et al; RE-LY Atrial Fibrillation Registry Investigators. Variations in cause and management of atrial fibrillation in a prospective registry of 15,400 emergency department patients in 46 countries: the RE-LY Atrial Fibrillation Registry. *Circulation*. 2014;129:1568–1576. doi: 10.1161/CIRCULATIONAHA.113.005451.
 20. Mant J, Hobbs FD, Fletcher K, Roalfo A, Fitzmaurice D, Lip GY, et al; BAFTA Investigators; Midland Research Practices Network (MidReC). Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial. *Lancet*. 2007;370:493–503. doi: 10.1016/S0140-6736(07)61233-1.
 21. Atzema CL, Austin PC, Chong AS, Dorian P, Jackevicius CA. The long-term use of warfarin among atrial fibrillation patients discharged from an emergency department with a warfarin prescription. *Ann Emerg Med*. 2015;66:347.e2–354.e2. doi: 10.1016/j.annemergmed.2015.03.024.
 22. Fuenzalida C, Ferró I, Siches C, Ambrós A, Sánchez M, Cabrera J, et al. Nurse-led educational intervention in patients with atrial fibrillation discharged from the emergency department reduces complications and short term admissions (article in Spanish). *Emergencias*. 2015;27:75–81.
 23. Coll-Vinent B, Pacheco G, Junyent M, Benito L, Hoyo J, García A, et al. [Impact of implementing common guidelines at different care levels in a healthcare area on the improvement of atrial fibrillation treatment]. *Rev Esp Cardiol*. 2007;60:392–403.
 24. Hendriks JM, de Wit R, Crijns HJ, Vrijhoef HJ, Prins MH, Pisters R, et al. Nurse-led care vs. usual care for patients with atrial fibrillation: results of a randomized trial of integrated chronic care vs. routine clinical care in ambulatory patients with atrial fibrillation. *Eur Heart J*. 2012;33:2692–2699. doi: 10.1093/eurheartj/ehs071.