



POSTER PRESENTATION

Open Access

Increased expression of A-kinase anchoring proteins in T cells from systemic lupus erythematosus patients

M J Pérez-Lorenzo, M Galindo, A J García-González, G Criado *

From 5th European Workshop on Immune-Mediated Inflammatory Diseases
Sitges-Barcelona, Spain. 1-3 December 2010

Introduction

Deficient activation of protein kinase A (PKA) is a characteristic of T cells in systemic lupus erythematosus (SLE). A-kinase Anchoring Proteins (AKAPs) associate to and regulate the activity of PKA [1]. Furthermore, some AKAPs are expressed in T cells and influence their function [2]. Therefore, altered expression and/or function of AKAPs can play a role in the deregulated activity of PKA observed in SLE T cells.

Aims

To analyse the expression of different AKAPs in T cells isolated from SLE patients.

Patients and methods

T cells were isolated by negative selection using magnetic beads from SLE patients (n = 12) and healthy controls (HC, n = 12). RNA was purified and levels of AKAP79, AKAP95 and AKAP450 mRNA were quantified by RT-qPCR. Subsequently, the analysis of AKAP79 expression was extended to include a total of 24 SLE patients and 19 HC. In addition, AKAP79 protein was detected by Western Blot and quantified with Quantity One software.

Results

Levels of AKAP450 mRNA were comparable in HC and SLE T cells (10.73 +/- 0.73 (HC, n = 12) vs 12.69 +/- 1.09 (SLE, n = 12), P = 0.15). However, T Cells from SLE patients had significantly higher levels of AKAP79 and AKAP95 mRNA than HC (AKAP79: 1.99 +/- 0.29 (HC, n = 19) vs 3.61 +/- 0.72 (SLE, n = 24), P = 0.04;

AKAP95: 2.58 +/- 0.18 (HC, n = 12) vs 4.13 +/- 0.31 (SLE, n = 12), P = 0.0005). Analysis of AKAP79 protein levels showed increased levels of AKAP79 in SLE T cells compared to HC T cells (0.38 ± 0.05 (HC) vs 0.75 ± 0.18 (SLE), P = 0.06).

Conclusions

Increased levels of AKAP79 and AKAP95 in T cells from SLE patients can contribute to the deregulation of PKA activity in these cells.

Published: 25 November 2010

References

1. Jarnaess E, Taskén K: Spatiotemporal control of cAMP signalling processes by anchored signalling complexes. *Biochem Soc Trans.* 2007, **35**:931-7.
2. Schillace RV, Carr SD: The role of Protein Kinase A and A Kinase Anchoring proteins in modulating T cell activation: Progress and future directions. *Crit Rev Immunol* 2006, **26**:113-31.

doi:10.1186/1479-5876-8-S1-P49

Cite this article as: Pérez-Lorenzo et al: Increased expression of A-kinase anchoring proteins in T cells from systemic lupus erythematosus patients. *Journal of Translational Medicine* 2010 **8**(Suppl 1):P49.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

