



Lack of molecular evidence of fecal-borne viruses in capybaras from São Paulo state, Brazil, 2018–2020: a minor public health issue

Lais Sampaio de Azevedo¹ · Yasmin França¹ · Ellen Viana¹ · Roberta Salzone Medeiros¹ · Simone Guadagnucci Morillo¹ · Raquel Guiducci¹ · Cibele Daniel Ribeiro¹ · Heloisa Rosa Vieira¹ · Karolina Morales Barrio-Nuevo² · Mariana Sequetin Cunha² · Juliana Mariotti Guerra³ · Dulcilena de Matos Castro e Silva⁴ · Valter Batista Duo Filho⁴ · Emerson Luiz Lima Araújo⁵ · Sérgio Roberto Santos Ferreira⁶ · Camila Freitas Batista⁶ · Gislaíne Celestino Dutra da Silva⁷ · Maurício Lacerda Nogueira⁷ · Cintia Mayumi Ahagon⁸ · Regina Célia Moreira⁹ · Lia Cunha⁹ · Vanessa Santos Morais¹⁰ · Antonio Charlys da Costa¹⁰ · Adriana Luchs^{1,11}

Received: 14 June 2022 / Accepted: 24 October 2022

© The Author(s) under exclusive licence to Sociedade Brasileira de Microbiologia 2022

Abstract

Capybara (*Hydrochoerus hydrochaeris*) is the world's largest rodent species distributed throughout South America. These animals are incredibly tolerant to anthropogenic environments and are occupying large urban centers. Capybaras are known to carry potentially zoonotic agents, including *R. rickettsia*, *Leishmania* spp., *Leptospira* spp., *Trypanosoma* spp., *Salmonella* spp., *Toxoplasma gondii*, and rabies virus. Focusing on the importance of monitoring potential sources of emerging zoonotic viruses and new viral reservoirs, the aim of the present study was to assess the presence of fecal-borne viruses in the feces of capybaras living in urban parks in São Paulo state, Brazil. A total of 337 fecal samples were collected between 2018 and 2020 and screened for the following: (i) Rotavirus group A (RVA) by ELISA; (ii) non-RVA species and Picobirnavirus (PBV) using PAGE; (iii) Human Bocaparvovirus (HBoV), Bufavirus (BuV), Tusavirus (TuV), and Cutavirus (CuV) qPCR; (iv) Human Enterovirus (EV), Norovirus GII (NoV), and Hantavirus by in houses RT-qPCR; (v) SARS-CoV-2 via commercial RT-qPCR kit assay; and (vi) Astrovirus (AstV) and Adenovirus (AdV) using conventional nested (RT)-PCRs. All fecal samples tested were negative for fecal-borne viruses. This study adds further evidence that the fecal-borne viruses is a minor public health issue in Brazilian capybaras, at least during the surveillance period and surveyed areas. Continuous monitoring of sylvatic animals is essential to prevent and control the emergence or re-emergence of newly discovered virus as well as viruses with known zoonotic potential.

Keywords Capybaras · Zoonosis · Molecular surveillance · Fecal-borne viruses · Virus reservoir

Zoonosis is an infectious disease that is transmitted between animals and humans and can be caused by bacteria, viruses, parasites, and even prions [1]. In recent years, there has been an emergence of zoonotic infections, especially viral-borne diseases, including the recently Zika virus outbreak and SARS-CoV-2 pandemic [2, 3]. Wild animal species in close contact with humans are often involved in viral-borne infections [1] and should be carefully considered when

assessing zoonotic disease emergence. Continuing wildlife surveillance is crucial for mitigating the consequences of emerging zoonotic viruses.

Rodents are the most species-rich order of mammals and well known recognized as reservoirs associated with new virus emergence [1]. Capybara (*Hydrochoerus hydrochaeris*) is the world's largest rodent species, displaying semi-aquatic habit and distributed throughout South America. Their high reproductive capacity, generalist feeding habits, and minimal habitat quality requirements have contributed to overpopulation in numerous regions of Brazil. In addition, capybaras are regularly found in close proximity to human population in urban parks across the country [4–6]. *R. rickettsii* is transmitted to humans mainly by the tick *Amblyomma sculptum*,

Responsible Editor: Luis Nero.

✉ Adriana Luchs
driluchs@gmail.com

Extended author information available on the last page of the article

which uses the capybara as its main host [7]. Capybaras are also known to carry other potentially zoonotic agents, including *Leishmania* spp., *Leptospira* spp., *Trypanosoma* spp., *Salmonella* spp., *Toxoplasma gondii*, and rabies virus [8, 9]. Therefore, monitoring these animals in areas cohabited by humans is vital to help outline measures in order to decrease the risk of a zoonotic emergence. The role of these rodents as asymptomatic hosts for potential zoonotic fecal-borne viruses has been poorly investigated in the country [4–6].

Enteric viruses can be transmitted by the fecal–oral route directly or by indirect contact via contaminated fluids, including surface water, food, and carriers such as fomites. In addition, enteric viruses are usually very stable in the environment and can survive under different conditions once these pathogens must tolerate the hostile conditions of the gastrointestinal tract [10]. Recently, there have been a growing number of reports describing the interspecies transmission of enteric viruses between animal and humans [1, 11]. Rotavirus, Picobirnavirus, Norovirus, Astrovirus, Bocaparvovirus, Enterovirus, and Adenovirus infect a wide range of vertebrates including human, domestic and wild mammals, especially rodents [1, 11, 12]. Moreover, rodents are also a source of Coronaviruses and Hantaviruses as well as newly identified viruses [12]. It is important to highlight that virus fecal shedding do not necessarily require replication in the intestine, like hepatitis A and E viruses that is shed in the feces despite their liver tropism [13]. In the same way, not all viruses that are shed in feces infect new hosts via the fecal–oral route, standing out the respiratory viruses [2], but in both cases, viral DNA/RNA can be detected in stool samples.

Focusing on the importance of monitoring potential sources of emerging zoonotic viruses and new viral reservoirs, the aim of the present study was to assess the presence of fecal-borne viruses in the feces of capybaras living in urban parks in São Paulo state, Brazil.

A total of 337 fecal samples were collected between 2018 and 2020 from 4 urban parks located in 2 municipalities of São Paulo state, Brazil. The (i) Yacht Club Santo Amaro (YCSA) is placed on the shores of the Guarapiranga Reservoir (Socorro District); the (ii) Novo Rio Pinheiros bike lane is placed on the banks of the Pinheiros River, both of them located at south region of São Paulo City (Santo Amaro District); the (iii) Tietê Ecological Park (PET) is placed on the east region of São Paulo City (Cangaíba District); and the (iv) Municipal Dam Park is placed on the east region of São José do Rio Preto City (Fig. 1). The 4 parks inhabited by capybaras present low water quality conditions, and the pollutants are associated with anthropogenic environments, mostly emerging contaminants resultant from incomplete degradation in sewage treatment plants (<https://cetesb.sp.gov.br/infoaguas/>). Stool samples were obtained *in natura*

and immediately stored at $-20\text{ }^{\circ}\text{C}$. All frozen specimens were taken to the Enteric Disease Laboratory, Virology Center, Adolfo Lutz Institute, São Paulo, Brazil. All animal procedures in this study were approved by the Ethics Committee on Animal Use (CEUA) guidelines of the Adolfo Lutz Institute, IAL, São Paulo, SP, Brazil (n^o 04/2018) and by Chico Mendes Institute for Biodiversity Conservation (ICMBio), Ministry of the Environment (No 77659–1/2021).

Controls for (RT-) PCR inhibitors or for DNA/RNA extraction efficiency are not available for animal origin specimens. In order to improve the reliability of negative PCR test result eventually obtained (i.e., assure that a negative PCR test result represent a true negative in the sample), we proposed to verify the viability of the capybaras stools screened in the present study using MALDI-TOF MS (matrix-assisted laser desorption ionization–time of flight mass spectrometry) and seeded experimental specimens with rotavirus and adenovirus reference strains (representing enteric RNA and DNA viruses, respectively). MALDI-TOF MS is a rapid and simple technique for microbial identification [14]. The detection of bacteria and fungi in capybara feces ensures an adequate logistics applied in the collection, packaging, and transport of feces samples from the field to the laboratory for viral testing. In addition, the capacity of recovering viruses artificially seeded in stools specimens offer the opportunity to evaluate the presence of potential PCR inhibitory factors. Therefore, based on collection site (representative sampling), a total of 35 stool samples were selected for MALDI-TOF MS investigation and 20 fecal specimens for seeded experiments.

One positive Rotavirus group A (RVA) G2P[4] specimen (GenBank accession No. JQ693560) and one positive Human Adenovirus F-40 (HAdV-F40) (GenBank accession No. MH201117) stool sample representing enteric RNA viruses and DNA viruses, respectively, were used to spike experimental specimens. The reference RVA and HAdV-F40 samples used here belong to the Enteric Diseases Laboratory collection, a regional reference center for viral gastroenteritis surveillance. The titers of positive RVA and HAdV-F samples used for the spiking experiments were not established; hence, detection limits were not evaluated. The aim of the seeded experiments was exclusively tested the sample matrices (stool) for containing potential PCR inhibitory factors. A total of 20 capybaras stool samples was spiked with the reference RVA G2P[4] and HAdV-F40 specimens. Briefly, 1 g of experimental stool and 1 g of each reference sample were added to 1 mL of TRIS/HCL in 1.5-mL microtubes. The suspensions were vigorously vortexed and centrifuged at $5000\times g$ for 30 min. The clarified supernatants were transferred to 2-mL cryotubes and stored at $-20\text{ }^{\circ}\text{C}$ for subsequent direct DNA/RNA extraction using *Bio Gene DNA/RNA Viral* (Quibasa – Química Básica Ltda, Belo Horizonte, BH, Brazil) according to the




Fig. 1 Map of São Paulo state, Southwestern region, Brazil; highlighting municipalities (in red and blue) and São Paulo municipalities' districts (in red, green, and yellow) from which samples were collected from free-living capybaras in urban parks between 2018 and 2020. Up left: Map of Brazil stressing São Paulo state. Down:

Municipalities surveyed: São Paulo and São José do Rio Preto. Up right: Districts surveyed at São Paulo municipality. Map was generated with QGIS software v2.14.9 (https://www.qgis.org/pt_BR/site/about/index.html)

manufacturer's instructions. Qualitative (RT-) PCRs were used for screening RVA and HAdV in seeded specimens. For RVA detection, the extracted dsRNA was subjected to G and P typing by semi-nested multiplex reverse transcription–polymerase chain reaction (RT-PCR) with specific primers for G and P types following protocol previously described by Gouvea et al. [15]. The presence of HAdV was detected by amplifying the Hexon gene, using primers AdVFhex(+)/ (–) (292 bp fragment) according to the protocol reported by Primo et al. [16]. All PCR products were visualized by Gel-Red™ (Biotium, Fremont, CA) staining and UV transillumination, following electrophoretic separation (1 h, 140 V) on 1.5% agarose gels. The G2 and P[4] genotypes were assigned using agarose gel analysis of second-round PCR products, 652 pb and 483 pb, respectively.

The 337 fecal specimens were screened for 12 different fecal-borne viruses: Rotavirus, Picobirnavirus, Human Bocaparvovirus, Bufavirus, Tusavirus, Cutavirus, Enterovirus, Norovirus genogroup GII, Hantavirus, SARS-CoV-2, Astrovirus, and Adenovirus (Supplement 1). All specimens were screened for RVA by a commercial enzyme-linked immunosorbent assay (ELISA) (RIDASCREEN® Rotavirus,

R-biopharm, Darmstadt, Germany), according to the manufacturer's protocol. Fecal samples were also tested for non-RVA species and Picobirnavirus (PBV) using polyacrylamide gel electrophoresis (PAGE) followed by silver staining of gels, according to standard procedure previously describe [17].

Viral nucleic acids were extracted from 10% fecal samples using *Bio Gene DNA/RNA Viral* (Quibasa – Química Básica Ltda, Belo Horizonte, BH, Brazil) according to the manufacturer's instructions. Human Bocaparvovirus (HBoV), Bufavirus (BuV), Tusavirus (TuV), and Cutavirus (CuV) were screened by quantitative real-time polymerase chain reaction (qPCR) following protocols previously reported [18, 19]. Human Enterovirus (EV), Norovirus GII (NoV), and Hantavirus were tested by reverse transcription quantitative real-time polymerase chain reaction (RT-qPCR) using procedures formerly defined by Zhang et al. [20], Kageyama et al. [21], and Nunes et al. [22], respectively. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was evaluated using commercial *Allplex™ 2019-nCoV* (Seegene Inc., Seoul, South Korea) RT-qPCR kit assay. Reverse transcription was carried out with random primers [23], and cDNA was screened for the presence of Astrovirus (AstV) using

heminested PCR targeting the RdRp gene following primers and protocol described by Chu et al. [24]. Nested-PCR amplification of a partial sequence of the adenoviral DNA polymerase gene was performed for Adenovirus (AdV) testing following primers and protocol described by Li et al. [12].

Of the 35 stool samples tested using MALDI-TOF MS, a total of 25 bacterial and 5 fungi species were identified (Table 1). MALDI-TOF MS is a rapid, simple, inexpensive, and high-throughput proteomic technique for identification of both bacteria and fungi, but it is important to highlight that its application is dependent on the reference strains

included in the database [25]. Therefore, for some fungi or bacteria species, their identification by MALDI-TOF MS may be failed during the sample screening conducted here. *Enterococcus hirae* (20%, 7/35), *Escherichia coli* (14.3%, 5/35), and *Escherichia hermannii* (5.7%, 2/35) were identified in the feces (Table 1). The presence of the enterobacterias in capybaras stool has been previously investigated in Brazil [8], and a high inter-individual bacterial variation was observed. This data was also consistent with a previous study conducted in Venezuela [26]. The successfully microbial identification in these 35 selected samples assures that the biological samples were of consistent quality for the intent analysis and study goals.

The qualitative PCR was able to detect RVA G2P[4] and HAdV-F40 in all 20 seeded capybaras stool samples. Virus detection in stool samples could be hampered by PCR inhibitors that may lead to false-negative results [27], especially considering feces from animal origin that can be a very heterogeneous biological material. Several PCR inhibitors have been characterized including phenolic compounds, glycogen, fats, cellulose, constituents of bacterial cells, heavy metals [28], and, more recently, dietary factor and gut microbiota [29]. All DNA/RNA extracts in seeded specimens amplified efficiently and did not differ from results of DNA/RNA extracted from the original reference strains used. Although viral titer was not determined prior to seeding experiments, the qualitative (RT-) PCRs protocols applied here successfully recover virus from the capybaras specimens. Thus, the stool samples employed in the present study were suitable for viral testing. This investigation emphasizes the need for PCR inhibitor evaluation when clinical samples obtained from animal origin are analyzed.

All samples tested negative for RVA, non-RVA species, HBoV, NoV GII, AstV, and AdV. These viruses are well known recognized as important cause associated with acute gastroenteritis in human and animals worldwide, as well as involved in interspecies transmission [30–34]. RVA, non-RVA species, HBoV, NoV GII, AstV, and AdV infections in rodents have been reported in the past, especially in rats [1, 11, 35, 36]. However, no capybara has tested positive for these viruses so far [4–6], in agreement with the data obtained here. There is also no serological evidence of viral gastroenteric infections [8, 37]. Collectively, capybaras' role in the epidemiological chain of enteric viral infections remains unknown. It is important to mentioning that besides anthropogenic influences, local water bodies are also contaminated with sewage, sustaining the continuous exposing of capybaras to enteric viruses and to a potentially zoonotic-transmission route as these animals are living at the human-animal interface environments.

PBV is ubiquitous in feces and gut contents of humans and animals with or without diarrhea but also detected in invertebrates and environmental samples [38]. The true host(s) and the role of these viruses as causative agents of

Table 1 Frequency of bacterial and fungi species identified in the fecal samples of 35 free-ranging capybaras (*Hydrochoerus hydrochaeris*), São Paulo state, Brazil, 2018–2020

Bacterial	Positive individuals	% of strains
<i>Acinetobacter lwoffii</i>	2	5.7%
<i>Acinetobacter schindleri</i>	1	2.8%
<i>Acinetobacter seifertii</i>	1	2.8%
<i>Arthrobacter gandavensis</i>	1	2.8%
<i>Bacillus altitudinis</i>	1	2.8%
<i>Bacillus cereus</i>	7	20%
<i>Bacillus koreensis</i>	1	2.8%
<i>Bacillus licheniformis</i>	1	2.8%
<i>Bacillus marisflavi</i>	10	28.6%
<i>Bacillus megaterium</i>	5	14.3%
<i>Bacillus pumilus</i>	10	28.6%
<i>Bacillus</i> spp.	2	5.7%
<i>Bacillus thuringiensis</i>	1	2.8%
<i>Cellulosimicrobium cellulans</i>	1	2.8%
<i>Enterobacter bugandensis</i>	2	5.7%
<i>Enterobacter kobei</i>	2	5.7%
<i>Enterococcus hirae</i>	7	20%
<i>Enterococcus mundtii</i>	1	2.8%
<i>Escherichia coli</i>	5	14.3%
<i>Escherichia hermannii</i>	2	5.7%
<i>Klebsiella variicola</i>	1	2.8%
<i>Kurthia gibsonii</i>	1	2.8%
<i>Microbacterium arborescens</i>	1	2.8%
<i>Pantoea séptica</i>	1	2.8%
<i>Serratia marcescens</i>	1	2.8%
Total	35 samples	100%
Fungi	Positive individuals	% of strains
<i>Cryptococcus diffluentis</i>	2	18%
<i>Cryptococcus laurentii</i>	3	27%
<i>Cryptococcus liquefaciens</i>	1	9%
<i>Rhodotorula mucilaginosa</i>	4	36%
<i>Torulaspora delbrueckii</i> (<i>Saccharomyces delbrueckii</i>)	1	9%
Total	11 samples	100%

Bold means the total of samples and the percentage obtained

gastroenteritis and respiratory infections or as opportunistic enteric pathogens remain unclear [39]. Although PBV has been frequently detected in a wide range of animal feces [40], there is no evidence of PBV infecting capybaras in viral surveillance studies [4–6] as also demonstrated in the present investigation.

Hantaviruses are considered infectious agents of great importance for public health and has been known to circulate throughout the Americas in the rodents [41]. The growing number of reports on hantaviruses in other vertebrate animal species, including bats and shrews, has stimulated the interest in search for discovering potential new reservoirs, and Brazil is not an exception [42, 43]. Most rodent hosts shed Hantavirus in their saliva, urine, and feces [44]. In a controlled experiment conducted with bank voles, Hardestam et al. [45] demonstrated the excretion of Puumala hantavirus in these three clinical samples, corroborating that these transmission routes may occur in nature and eventually able to reach humans. None of the capybaras tested here were positive for viral hantavirus RNA in feces. Nevertheless, one limitation of the present study is the lack of serum samples collected from the animals in order to verify serological evidence of previous infections by hantavirus.

BuV, TuV, and CuV are among the newest parvoviruses described in humans [46]. These viruses have been also reported in wild and domestic animals, including in rodents, but information on their epidemiology is rather limited [47–49]. BuV, CuV, and TuV have been described circulating in feces of human and animal origin in Brazil [50, 51]. Here we investigated the capybaras potential to play a role as harboring new and poorly surveyed viruses in the country. No positivity for BuV, CuV, and TuV were detected in capybaras feces, excluding them as potential hosts or reservoirs of these new parvoviruses in the investigated areas.

In the same way, we aimed to determine whether capybaras could be sheltering EV strains. EV are classified into twelve species EV-A to EV-L (<https://talk.ictvonline.org/taxonomy/>), where species EV-A to EV-D commonly infects humans, species EV-E to EV-G and EV-K causes diseases affecting livestock industry, including cattle and swine's, species EV-H and EV-J was identified in monkeys, and species EV-I and EV-L were discovered in fecal samples collected from dromedaries and goats, respectively [52–55]. Given the evidence that EV naturally circulate in animals and that human EV have been detected in distinct mammalian species, including in Brazil [56–58], it must be considered its potential for interspecies transmission [59]. Our molecular tests detected no evidence of EV infection among capybaras populations living in the Sao Paulo state, reducing concerns about rodent reservoirs as a potential source of emerging EV infections.

Humans have infected a wide range of animals with SARS-CoV-2 [60–62]. Potential establishment of new

reservoirs, like occurred in white-tailed deer populations in USA, rises critical concerns about SARS-CoV-2 evolution and epidemiology but mostly brings serious implications for public health issues, especially considering the risk of spillback of novel variants to humans [61]. Continuous monitoring of SARS-CoV-2 in animals is essential for risk management, particularly considering capybaras as they are water-loving rodents found in areas with abundant water sources, incredibly tolerant to anthropogenic environments and present in large urban centers such as Sao Paulo and São José do Rio Preto cities where water bodies are contaminated, mainly with domestic sewage [63–65]. Several studies have demonstrated the presence of the SARS-CoV-2 in human stool samples [66, 67]. Therefore, viral RNA is shed and disposed in wastewaters, highlighting a potential indirect infection pathway to humans or other susceptible mammals' species [68]. No specimen tested positive for SARS-CoV-2 in the present study, indicating that although the capybaras are in close contact with contaminated effluents, they are not being infected, and they probably do not constitute a reservoir for SARS-CoV-2 in the studied regions. It is important to mention that enteric coronavirus has been previously detected in capybaras feces from Sao Paulo state using transmission electron microscopy technique [4].

The knowledge about the presence and diversity of fecal-borne viruses in capybaras is so far limited, and a better understanding of a potential role of enteric infections is that animals are essential to develop preventive measures. The present study reports a molecular surveillance of various viral pathogens in Brazilian capybaras and adds further evidence that the fecal-borne viruses is a minor public health issue, at least during the surveillance period and surveyed areas. Continuous monitoring of sylvatic animals is essential to prevent and control the emergence or re-emergence of newly discovered virus as well as viruses with known zoonotic potential, especially in anthropogenic environments. Capybaras are extremely tolerant to environmental changes and exhibit high vagility along waterways, thus configuring a potential to trigger emerging zoonosis.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s42770-022-00859-2>.

Acknowledgements We thank the staff of the Hepatitis Laboratory of Adolfo Lutz Institute Marcilio Figueiredo Lemos, Adriana Parise Compri, and Vanessa Cristina Martins Silva for laboratory assistance. We thank Graduate Program in Science, Coordinator for Disease Control, Ministry of Health-PPG-CCD-SES/SP, and Coordination for the Improvement of Higher Education Personnel (CAPES).

Author contribution AL conceived and designed the study protocol; LSA, YF, EV, RSM, SGM, RG, CDR, HRV, KMBN, MSC, JMG, DMCS, VBDF, ELLA, SRSF, CFB, GCDS, MLN, CMA, RCM, LC, VSM, and ACC participated in the conduct of the study; LSA, DMCS, and VBDF conduct the MALDI-TOF MS assays; LSA conducted the PCR inhibition evaluations; LSA, EV, KMBN, MSC, SRSF, CFB,

GCDS, and AL acquired samples and data; LSA and RG performed the ELISA tests and conduct PAGE screening; LSA, YF, EV, RSM, SGM, RG, CDR, HRV, MSC, JMG, CMA, LC, VSM, ACC, and AL performed the RNA/DNA extraction and the PCR tests; LSA, MSC, JMG, ELLA, MLN, RCM, ACC, and AL analyzed, interpreted the data, and drafted the manuscript; all authors critically revised the manuscript for intellectual content and approved the final version. AL is guarantor of the paper.

Funding This study was supported by the Sao Paulo Research Foundation (FAPESP) through the grant numbers #2015/12944–9 and #2020/14786–0 to AL, #2020/02469–0 to YF, #2020/11182–6 to RSM, and #2019/21706–5 to VSM; by Brazilian National Council for Scientific and Technological Development (CNPq) through the grant #404510/2021–3 to JMG; by Fundo Especial de Saúde para Imunização em Massa e Controle de Doenças (FESIMA) CAF N° #001/2021 and #060/2021 to AL, and #076/2019 to MSC. ACC is supported by a scholarship from Hospital das Clínicas, Faculty of Medicine, University of São Paulo (HCFMUSP), with funds donated by NUBANK under the #HCCOMVIDA scheme. MLN is partly funded by the Centers for Research in Emerging Infectious Diseases [CREID], “The Coordinating Research on Emerging Arboviral Threats Encompassing the Neotropics [CREATE-NEO]” grant #U01-AI151807 by the National Institutes of Health [NIH/USA].

Declarations

Competing interest The authors declare no competing interests.

References


- Niendorf S, Harms D, Hellendahl KF, Heuser E, Böttcher S, Bock CT, Ulrich RG (2021) Presence and diversity of different enteric viruses in wild Norway rats (*Rattus norvegicus*). *Viruses* 13(6):992. <https://doi.org/10.3390/v13060992>
- Jones DL, Baluja MQ, Graham DW, Corbishley A, McDonald JE, Malham SK, Hillary LS, Connor TR, Gaze WH, Moura IB, Wilcox MH, Farkas K (2020) Shedding of SARS-CoV-2 in feces and urine and its potential role in person-to-person transmission and the environment-based spread of COVID-19. *Sci Total Environ* 749:141364. <https://doi.org/10.1016/j.scitotenv.2020.141364>
- Costa ACDS, Hasan MM, Xenophontos E, Mohanan P, Bassey EE, Hashim HT, Ahmad S, Essar MY (2021) COVID-19 and Zika: an emerging dilemma for Brazil. *J Med Virol* 93(7):4124–4126. <https://doi.org/10.1002/jmv.27006>
- Catrox MHB, Miranda LB, Lavorenti A, Petrella S, Melo NA, Martins AMCPRF (2010) Detection of coronavirus in capybaras (*Hydrochoerus hydrochaeris*) by transmission electron microscopy in São Paulo. *Brazil Int J Morphol* 28(2):549–555
- Dutra LA, de Freitas Almeida GM, Oliveira GP, Abrahão JS, Kroon EG, Trindade GS (2017) Molecular evidence of Orthopoxvirus DNA in capybara (*Hydrochoerus hydrochaeris*) stool samples. *Arch Virol* 162(2):439–448. <https://doi.org/10.1007/s00705-016-3121-3>
- Fontenele RS, Lacorte C, Lamas NS, Schmidlin K, Varsani A, Ribeiro SG (2019) Single stranded DNA viruses associated with capybara faeces sampled in Brazil. *Viruses* 11(8):710. <https://doi.org/10.3390/v11080710>
- Luz HR, Costa FB, Benatti HR, Ramos VN, de A Serpa MC, Martins TF, Acosta ICL, Ramirez DG, Muñoz-Leal S, Ramirez-Hernandez A, Binder LC, Carvalho MP, Rocha V, Dias TC, Simeoni CL, Brites-Neto J, Brasil J, Nieves AM, Monticelli PF, Moro MEG, Lopes B, Aguiar DM, Pacheco RC, Souza CE, Piovezan U, Juliano R, Ferraz KMPMB, Szabó MPJ, Labruna MB (2019) Epidemiology of capybara-associated Brazilian spotted fever. *PLoS Negl Trop Dis* 13(9):e0007734. <https://doi.org/10.1371/journal.pntd.0007734>
- Chiacchio RG, Prioste FE, Vanstreels RE, Knöbl T, Kolber M, Miyashiro SI, Matsushima ER (2014) Health evaluation and survey of zoonotic pathogens in free-ranging capybaras (*Hydrochoerus hydrochaeris*). *J Wildl Dis* 50(3):496–504. <https://doi.org/10.7589/2013-05-109>
- Barbosa AV, Medaglia ML, Soares HS, Quixabeira-Santos JC, Gennari SM, Damaso CR (2014) Presence of neutralizing antibodies to Orthopoxvirus in capybaras (*Hydrochoerus hydrochaeris*) in Brazil. *J Infect Dev Ctries* 15:8(12):1646–9. <https://doi.org/10.3855/jidc.5216>
- de Graaf M, Beck R, Caccio SM, Duim B, Fraaij P, Le Guyader FS, Lecuit M, Le Pendu J, de Wit E, Schultsz C (2017) Sustained fecal-oral human-to-human transmission following a zoonotic event. *Curr Opin Virol* 22:1–6. <https://doi.org/10.1016/j.coviro.2016.11.001>
- Sachsenröder J, Braun A, Machnowska P, Ng TFF, Deng X, Guenther S, Bernstein S, Ulrich RG, Delwart E, Johne R (2014) Metagenomic identification of novel enteric viruses in urban wild rats and genome characterization of a group A rotavirus. *J Gen Virol* 95(Pt 12):2734–2747. <https://doi.org/10.1099/vir.0.070029-0>
- Li Y, Ge X, Zhang H, Zhou P, Zhu Y, Zhang Y et al (2010) Host range, prevalence, and genetic diversity of adenoviruses in bats. *J Virol* 84(8):3889–3897. <https://doi.org/10.1128/JVI.02497-09>
- Cuthbert JA (2001) Hepatitis A: old and new. *Clin Microbiol Rev* 14(1):38–58. <https://doi.org/10.1128/CMR.14.1.38-58.2001>
- van Veen SQ, Claas EC, Kuijper EJ (2010) High-throughput identification of bacteria and yeast by matrix-assisted laser desorption ionization-time of flight mass spectrometry in conventional medical microbiology laboratories. *J Clin Microbiol* 48(3):900–907. <https://doi.org/10.1128/JCM.02071-09>
- Gouvea V, Glass RI, Woods P, Taniguchi K, Clark HF, Forrester B, Fang ZY (1999) Polymerase chain reaction amplification and typing of rotavirus nucleic acid from stool specimens. *J Clin Microbiol* 28(2):276–282. <https://doi.org/10.1128/jcm.28.2.276-282.1990>
- Primo D, Pacheco GT, Timenetsky MDCST, Luchs A (2018) Surveillance and molecular characterization of human adenovirus in patients with acute gastroenteritis in the era of rotavirus vaccine, Brazil, 2012–2017. *J Clin Virol* 109:35–40. <https://doi.org/10.1016/j.jcv.2018.10.010>
- Herring AJ, Inglis NF, Ojeh CK, Snodgrass DR, Menzies JD (1982) Rapid diagnosis of rotavirus infection by direct detection of viral nucleic acid in silver-stained polyacrylamide gels. *J Clin Microbiol* 16:473–477
- Lu X, Chittaganpitch M, Olsen SJ, Mackay IM, Sloots TP, Fry AM, Erdman DD (2006) Real-time PCR assays for detection of bocavirus in human specimens. *J Clin Microbiol* 44(9):3231–3235. <https://doi.org/10.1128/JCM.00889-06>
- Väisänen E, Fu Y, Koskenmies S, Fyhrquist N, Wang Y, Keinonen A, Mäkisalo H, Väkevä L, Pitkänen S, Ranki A, Hedman K, Söderlund-Venermo M (2019) Cutaneous DNA in malignant and nonmalignant skin of cutaneous T-cell lymphoma and organ transplant patients but not of healthy adults. *Clin Infect Dis* 68(11):1904–1910. <https://doi.org/10.1093/cid/ciy806>
- Zhang S, Wang J, Yan Q, He S, Zhou W, Ge S, Xia N (2014) A one-step, triplex, real-time RT-PCR assay for the simultaneous detection of enterovirus 71, coxsackie A16 and pan-enterovirus in a single tube. *PLoS ONE* 9(7):e102724. <https://doi.org/10.1371/journal.pone.0102724>
- Kageyama T, Kojima S, Shinohara M, Uchida K, Fukushi S, Hoshino FB, Takeda N, Katayama K (2003) Broadly reactive

- and highly sensitive assay for Norwalk-like viruses based on real-time quantitative reverse transcription-PCR. *J Clin Microbiol* 41(4):1548–1557. <https://doi.org/10.1128/JCM.41.4.1548-1557.2003>
22. Nunes BTD, de Mendonça MHR, Simith DB et al (2019) Development of RT-qPCR and semi-nested RT-PCR assays for molecular diagnosis of hantavirus pulmonary syndrome. *PLoS Negl Trop Dis* 13(12):e0007884. <https://doi.org/10.1371/journal.pntd.0007884>
 23. Iturriza-Gomara M, Green J, Brown DWG, Desselberger U, Gray JJ (1999) Comparison of specific and random priming in the reverse transcriptase polymerase chain reaction for genotyping group A rotaviruses. *J Virol Meth* 78:93–103
 24. Chu DK, Poon LL, Guan Y, Peiris JS (2008) Novel astroviruses in insectivorous bats. *J Virol* 82(18):9107–9114. <https://doi.org/10.1128/JVI.00857-08>
 25. Samb-Ba B, Mazonot C, Gassama-Sow A, Dubourg G, Richet H, Hugon P, Lagier JC, Raoult D, Fenollar F (2014) MALDI-TOF identification of the human gut microbiome in people with and without diarrhea in Senegal. *PLoS ONE* 9(5):e87419. <https://doi.org/10.1371/journal.pone.0087419>
 26. García-Amado MA, Godoy-Vitorino F, Piceno YM, Tom LM, Andersen GL, Herrera EA, Domínguez-Bello MG (2012) Bacterial diversity in the cecum of the world's largest living rodent (*Hydrochoerus hydrochaeris*). *Microb Ecol* 63(4):719–725. <https://doi.org/10.1007/s00248-011-9963-z>
 27. Valentine-Thon E (2002) Quality control in nucleic acid testing—where do we stand? *J Clin Virol* 25(Suppl 3):S13-21. [https://doi.org/10.1016/s1386-6532\(02\)00196-8](https://doi.org/10.1016/s1386-6532(02)00196-8)
 28. Monteiro L, Bonnemaison D, Vekris A, Petry KG, Bonnet J, Vidal R, Cabrita J, Mégraud F (1997) Complex polysaccharides as PCR inhibitors in feces: *Helicobacter pylori* model. *J Clin Microbiol* 35(4):995–998. <https://doi.org/10.1128/jcm.35.4.995-998.1997>
 29. Oikarinen S, Tauriainen S, Viskari H, Simell O, Knip M, Virtanen S, Hyöty H (2009) PCR inhibition in stool samples in relation to age of infants. *J Clin Virol* 44(3):211–214. <https://doi.org/10.1016/j.jcv.2008.12.017>
 30. Manteufel J, Truyen U (2008) Animal bocaviruses: a brief review. *Intervirology* 51(5):328–334. <https://doi.org/10.1159/000173734>
 31. Martella V, Bányai K, Matthijssens J, Buonavoglia C, Ciarlet M (2010) Zoonotic aspects of rotaviruses. *Vet Microbiol* 140(3–4):246–255. <https://doi.org/10.1016/j.vetmic.2009.08.028>
 32. De Benedictis P, Schultz-Cherry S, Burnham A, Cattoli G (2011) Astrovirus infections in humans and animals - molecular biology, genetic diversity, and interspecies transmissions. *Infect Genet Evol* 11(7):1529–1544. <https://doi.org/10.1016/j.meegid.2011.07.024>
 33. Borkenhagen LK, Fieldhouse JK, Seto D, Gray GC (2019) Are adenoviruses zoonotic? A systematic review of the evidence. *Emerg Microbes Infect* 8(1):1679–1687. <https://doi.org/10.1080/22221751.2019.1690953>
 34. Villabruna N, Koopmans MPG, de Graaf M (2019) Animals as reservoir for human norovirus. *Viruses* 11(5):478. <https://doi.org/10.3390/v11050478>
 35. Kumakamba C, N'Kawa F, Kingebeni PM, Losoma JA, Lukusa IN, Muyembe F, Mulembakani P, Makuwa M, LeBreton M, Gillis A, Rimoin AW, Hoff NA, Schneider BS, Monagin C, Joly DO, Wolfe ND, Rubin EM, Tamfum JJM, Lange CE (2019) Analysis of adenovirus DNA detected in rodent species from the Democratic Republic of the Congo indicates potentially novel adenovirus types. *New Microbes New Infect* 34:100640. <https://doi.org/10.1016/j.nmni.2019.100640>
 36. You FF, Zhang MY, Wu F, Li QS, Chen Q (2022) Human bocavirus 2 detected in *Rattus norvegicus* feces in China. *Arch Virol* 167(1):171–175. <https://doi.org/10.1007/s00705-021-05274-w>
 37. Almeida MF, Massad E, Aguiar EA, Martorelli LF, Joppert AM (2001) Neutralizing antirabies antibodies in urban terrestrial wildlife in Brazil. *J Wildl Dis* 37(2):394–398. <https://doi.org/10.7589/0090-3558-37.2.394>
 38. Ghosh S, Malik YS (2021) The true host/s of picobirnaviruses. *Front Vet Sci* 7:615293. <https://doi.org/10.3389/fvets.2020.615293>
 39. Malik YS, Kumar N, Sharma K, Dhama K, Shabbir MZ, Ganesh B, Kobayashi N, Banyai K (2014) Epidemiology, phylogeny, and evolution of emerging enteric Picobirnaviruses of animal origin and their relationship to human strains. *Biomed Res Int* 2014:780752. <https://doi.org/10.1155/2014/780752>
 40. Ganesh B, Masachessi G, Mladenova Z (2014) Animal picobirnavirus. *Virusdisease* 25(2):223–238. <https://doi.org/10.1007/s13337-014-0207-y>
 41. de Oliveira RC, Guterres A, Fernandes J, D'Andrea PS, Bonvicino CR (2014) de Lemos ER (2014) Hantavirus reservoirs: current status with an emphasis on data from Brazil. *Viruses* 6(5):1929–1973. <https://doi.org/10.3390/v6051929>
 42. Sabino-Santos G Jr, Maia FGM, Martins RB, Gagliardi TB, Souza WM, Muylaert RL, Luna LKS, Melo DM, Cardoso RS, Barbosa NDS, Pontelli MC, Mamani-Zapana PR, Vieira TM, Melo NM, Jonsson CB, Goodin D, Salazar-Bravo J, daSilva LLP, Arruda E, Figueiredo LTM (2018) Natural infection of neotropical bats with hantavirus in Brazil. *Sci Rep* 8(1):9018. <https://doi.org/10.1038/s41598-018-27442-w>
 43. Lwande OW, Mohamed N, Bucht G, Ahlm C, Olsson G, Evander M (2020) Seewis hantavirus in common shrew (*Sorex araneus*) in Sweden. *Virol J* 17(1):198. <https://doi.org/10.1186/s12985-020-01477-w>
 44. Forbes KM, Sironen T, Plyusnin A (2018) Hantavirus maintenance and transmission in reservoir host populations. *Curr Opin Virol* 28:1–6. <https://doi.org/10.1016/j.coviro.2017.09.003>
 45. Hardestam J, Karlsson M, Falk KI, Olsson G, Klingström J, Lundkvist A (2008) Puumala hantavirus excretion kinetics in bank voles (*Myodes glareolus*). *Emerg Infect Dis* 14(8):1209–1215. <https://doi.org/10.3201/eid1408.080221>
 46. Väisänen E, Fu Y, Hedman K, Söderlund-Venermo M (2017) Human protoparvoviruses. *Viruses* 9(11):354. <https://doi.org/10.3390/v9110354>
 47. Yang S, Liu D, Wang Y, Qu F, He Y, Sun Z, Shen Q, Li W, Fu X, Deng X, Zhang W, Delwart E (2016) Bufavirus Protoparvovirus in feces of wild rats in China. *Virus Genes* 52(1):130–133. <https://doi.org/10.1007/s11262-015-1262-1>
 48. Ganji VK, Buddala B, Yella NR, Putty K (2022) First report of canine bufavirus in India. *Arch Virol* 167(4):1145–1149. <https://doi.org/10.1007/s00705-022-05398-7>
 49. Reuter G, Pankovics P, László Z, Gáspár G, Hui A, Delwart E, Boros Á (2022) Human-stool-associated tusavirus (Parvoviridae) in domestic goats and sheep. *Arch Virol* 167(5):1307–1310. <https://doi.org/10.1007/s00705-022-05424-8>
 50. Kluge M, Campos FS, Tavares M, de Amorim DB, Valdez FP, Giongo A, Roehle PM, Franco AC (2016) Metagenomic survey of viral diversity obtained from feces of Subantarctic and South American fur seals. *PLoS ONE* 11(3):e0151921. <https://doi.org/10.1371/journal.pone.0151921>
 51. Phan TG, Dreno B, da Costa AC, Li L, Orlandi P, Deng X, Kapusinszky B, Siqueira J, Knol AC, Halary F, Dantal J, Alexander KA, Pesavento PA, Delwart E (2016) A new protoparvovirus in human fecal samples and cutaneous T cell lymphomas (mycosis fungoides). *Virology* 496:299–305. <https://doi.org/10.1016/j.virol.2016.06.013>
 52. Van Nguyen D, Harvala H, Ngole EM, Delaporte E, Woolhouse ME, Peeters M, Simmonds P (2014) High rates of infection with

- novel enterovirus variants in wild populations of mandrills and other old world monkey species. *J Virol* 88(11):5967–5976. <https://doi.org/10.1128/JVI.00088-14>
53. Woo PC, Lau SK, Li T, Jose S, Yip CC, Huang Y, Wong EY, Fan RY, Cai JP, Wernery U, Yuen KY (2015) A novel dromedary camel enterovirus in the family Picornaviridae from dromedaries in the Middle East. *J Gen Virol* 96(Pt 7):1723–1731. <https://doi.org/10.1099/vir.0.000131>
 54. Tsuchiaka S, Rahpaya SS, Otomaru K, Aoki H, Kishimoto M, Naoi Y, Omatsu T, Sano K, Okazaki-Terashima S, Katayama Y, Oba M, Nagai M, Mizutani T (2017) Identification of a novel bovine enterovirus possessing highly divergent amino acid sequences in capsid protein. *BMC Microbiol* 17(1):18. <https://doi.org/10.1186/s12866-016-0923-0>
 55. Wang M, He J, Lu H, Liu Y, Deng Y, Zhu L, Guo C, Tu C, Wang X (2017) A novel enterovirus species identified from severe diarrheal goats. *PLoS ONE* 12(4):e0174600. <https://doi.org/10.1371/journal.pone.0174600>
 56. Waldman EA, Moreira RC, Saez SG, Souza DF, Carmona Rde C, Takimoto S, de A Cortes V (1996) Human enterovirus infection in stray dogs Some aspects of interest to public health. *Rev Inst Med Trop Sao Paulo* 38(2):157–61. <https://doi.org/10.1590/s0036-46651996000200012>
 57. Mombo IM, Lukashev AN, Bleicker T, Brünink S, Berthet N, Maganga GD, Durand P, Arnathau C, Boundenga L, Ngoubangoye B, Boué V, Liégeois F, Ollomo B, Prugnolle F, Drexler JF, Drosten C, Renaud F, Rougeron V, Leroy E (2017) African non-human primates host diverse enteroviruses. *PLoS ONE* 12(1):e0169067. <https://doi.org/10.1371/journal.pone.0169067>
 58. Yang S, He Y, Chen X, Kalim U, Wang Y, Yang S, Qi H, Cheng H, Lu X, Wang X, Shen Q, Zhang W (2021) Viral metagenomics reveals diverse viruses in the feces samples of raccoon dogs. *Front Vet Sci* 8:693564. <https://doi.org/10.3389/fvets.2021.693564>
 59. Fieldhouse JK, Wang X, Mallinson KA, Tsao RW, Gray GC (2018) A systematic review of evidence that enteroviruses may be zoonotic. *Emerg Microbes Infect* 7(1):164. <https://doi.org/10.1038/s41426-018-0159-1>
 60. Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E, Molenkamp R, van der Spek A, Tolsma P, Rietveld A, Brouwer M, Bouwmeester-Vincken N, Harders F, Hakze-van der Honing R, Wegdam-Blans MCA, Bouwstra RJ, GeurtsvanKessel C, van der Eijk AA, Velkers FC, Smit LAM, Stegeman A, van der Poel WHM, Koopmans MPG (2021) Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. *Science* 371(6525):172–177. <https://doi.org/10.1126/science.abe5901>
 61. Hale VL, Dennis PM, McBride DS, Nolting JM, Madden C, Huey D, Ehrlich M, Grieser J, Winston J, Lombardi D, Gibson S, Saif L, Killian ML, Lantz K, Tell RM, Torchetti M, Robbe-Austerman S, Nelson MI, Faith SA, Bowman AS (2022) SARS-CoV-2 infection in free-ranging white-tailed deer. *Nature* 602(7897):481–486. <https://doi.org/10.1038/s41586-021-04353-x>
 62. Jairak W, Chamsai E, Udom K, Charoenkul K, Chaiyawong S, Techakriengkrai N, Tangwangvivat R, Suwannakarn K, Amonsin A (2022) SARS-CoV-2 delta variant infection in domestic dogs and cats. *Thailand Sci Rep* 12(1):8403. <https://doi.org/10.1038/s41598-022-12468-y>
 63. Melo CA, Moreira AB, Bisinoti MC (2009) Seasonal and spatial trend of pollutants in the waters of the São José do Rio Preto municipal dam, São Paulo state. *Brazil Quim Nova* 32(6):1436–1441
 64. Dias TC, Stabach JA, Huang Q, Labruna MB, Leimgruber P, Ferraz KMPMB, Lopes B, Luz HR, Costa FB, Benatti HR, Correia LR, Nievas AM, Monticelli PF, Piovezan U, Szabó MPJ, Aguiar DM, Brites-Neto J, Port-Carvalho M, Rocha VJ (2020) Habitat selection in natural and human-modified landscapes by capybaras (*Hydrochoerus hydrochaeris*), an important host for *Amblyomma sculptum* ticks. *PLoS ONE* 15(8):e0229277. <https://doi.org/10.1371/journal.pone.0229277>
 65. Gadda TMC, Souza JMT, Paula GAR, van Kaick TS, Gervásio JHDB (2021) The international biodiversity agenda at the local level: the case of Capybaras in Curitiba, Brazil. *Ambient Soc* 24:1–21
 66. Xu Y, Li X, Zhu B, Liang H, Fang C, Gong Y, Guo Q, Sun X, Zhao D, Shen J, Zhang H, Liu H, Xia H, Tang J, Zhang K, Gong S (2020) Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med* 26(4):502–505. <https://doi.org/10.1038/s41591-020-0817-4>
 67. Fumian TM, Malta FC, Dos Santos DRL, Pauvolid-Corrêa A, Fialho AM, Leite JPG, Miagostovich MP (2021) SARS-CoV-2 RNA detection in stool samples from acute gastroenteritis cases. *Brazil J Med Virol* 93(4):2543–2547. <https://doi.org/10.1002/jmv.26786>
 68. Kitajima M, Ahmed W, Bibby K, Carducci A, Gerba CP, Hamilton KA, Haramoto E, Rose JB (2020) SARS-CoV-2 in wastewater: state of the knowledge and research needs. *Sci Total Environ* 739:139076. <https://doi.org/10.1016/j.scitotenv.2020.139076>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Lais Sampaio de Azevedo¹ · Yasmin França¹ · Ellen Viana¹ · Roberta Salzone Medeiros¹ ·
Simone Guadagnucci Morillo¹ · Raquel Guiducci¹ · Cibele Daniel Ribeiro¹ · Heloisa Rosa Vieira¹ ·
Karolina Morales Barrio-Nuevo² · Mariana Sequetin Cunha² · Juliana Mariotti Guerra³ ·
Dulcilena de Matos Castro e Silva⁴ · Valter Batista Duo Filho⁴ · Emerson Luiz Lima Araújo⁵ ·
Sérgio Roberto Santos Ferreira⁶ · Camila Freitas Batista⁶ · Gislaine Celestino Dutra da Silva⁷ ·
Maurício Lacerda Nogueira⁷ · Cintia Mayumi Ahagon⁸ · Regina Célia Moreira⁹ · Lia Cunha⁹ ·
Vanessa Santos Morais¹⁰ · Antonio Charlys da Costa¹⁰ · Adriana Luchs^{1,11} 

¹ Enteric Disease Laboratory, Virology Center, Adolfo Lutz Institute, Sao Paulo, Brazil

² Vector Borne Diseases Laboratory, Virology Center, Adolfo Lutz Institute, Sao Paulo, Brazil

³ Quantitative Pathology Laboratory, Pathology Center, Adolfo Lutz Institute, Sao Paulo, Brazil

⁴ Micology Laboratory, Parasitology Center, Adolfo Lutz Institute, Sao Paulo, Brazil

⁵ General Coordination of Public Health, Laboratories of the Strategic Articulation, Department of the Health, Surveillance Secretariat, Ministry of Health (CGLAB/DAEVS/SVS-MS), Brasília, Brazil

⁶ Cruzeiro Do Sul University, Sao Paulo, Brazil

⁷ São José Do Rio Preto School of Medicine (FAMERP), São Paulo, Brazil

⁸ Blood and Sexual Diseases – Retrovirus Laboratory, Virology Center, Adolfo Lutz Institute, Sao Paulo, Brazil

⁹ Hepatitis Laboratory, Virology Center, Adolfo Lutz Institute, Sao Paulo, Brazil

¹⁰ Medical Parasitology Laboratory (LIM/46), Institute of Tropical Medicine, University of Sao Paulo, Sao Paulo, Brazil

¹¹ Centro de Virologia, Núcleo de Doenças Entéricas, Instituto Adolfo Lutz, Av. Dr Arnaldo, nº 355, São Paulo, SP 01246-902, Brasil