

PRIMARY RESEARCH

Open Access

Target delivery of MYCN siRNA by folate-nanoliposomes delivery system in a metastatic neuroblastoma model

Qiqi Zhu, Chen Feng, Weiwei Liao, Yan Zhang and Suoqin Tang*

Abstract

Background: Folate-nanoliposomes delivery system has emerged recently as a specific and safety delivery method and gradually used as the carrier of a variety kinds of drugs including compounds, plasmids and siRNAs.

Methods: In this study, we established a bone marrow and bone metastasis xenograft mouse model by injecting the LA-N-5 cell into the bone marrow cavity. Fluorescence microscopy, TUNEL Assay, Quantitative RT-PCR and western blot were conducted to analysis the distribution of folate-nanoliposomes entrapped MYCN (V-myc myelocytomatosis viral related oncogene) siRNA in mice and the relevant suppression effect.

Results: The folate-nanoliposomes entrapped MYCN siRNA can be specifically distributed in tumor tissues. Further study shows that folate-nanoliposomes entrapped MYCN siRNA lead to MYCN mRNA expression significantly down-regulated (>50%, and $p < 0.05$) compared with negative control siRNA treatment. MYCN protein expression was inhibited about 60% in vivo, thus induced tumor cell apoptosis markedly.

Conclusion: This study point to a new way for treatment of metastatic neuroblastoma and could widen the application of folate-nanoliposomes delivery system in tumor therapy.

Keywords: Folate-nanoliposomes, Metastatic neuroblastoma, MYCN, siRNA

Introduction

Neuroblastoma is the most common extracranial solid tumor occurring in children, which accounts for about 15% of all childhood cancer deaths [1,2]. It is generally accepted that the amplification of the MYCN proto-oncogene is highly relevant to the genesis and prognosis of neuroblastoma [3,4]. MYCN is an important member of the myc family that includes 3 strong oncogenes, MYC, MYCN, and MYCL [5]. Targeting MYCN gene has developed into a promising therapeutic strategy [6-8]. Small interference RNA (siRNA) is synthetic short double-stranded RNA, which can induce the destruction of homologous mRNA when introduced into a cell [9]. siRNA has been developed into an effective tool for suppressing target proteins expression by specifically digesting their mRNAs [10]. siRNAs have been widely investigated

as therapeutic agents to treat a wide range of human diseases including neuroblastoma [7,11].

Folate, a nontoxic low-weight compound, is vital for tumor cell proliferation and survival. Studies show that tumor cells can express 200-folds more folate receptor on the membrane than normal cells, which is accordance to the high intake of folate in tumor cells. Folate receptor is overexpressed in a wide range of human cancer cells, including ovarian cancer, endometrial cancer, head and neck cancers, myeloid leukaemia, and some pediatric cancers [12]. Folate conjugating on liposomes with entrapped drugs [13-16] or plasmids [17-20] has been used successfully to deliver drugs to folate-receptor-expressing cancer cells. Our previous study had revealed that the folate receptor was highly expressed on the membrane of LA-N-5 neuroblastoma cells. In this study, we want to further investigate whether folate receptor-targeted liposome can act as a good delivery tool of MYCN siRNA and exert a satisfying therapeutic effect on neuroblastoma.

* Correspondence: suoqin@yahoo.com

Department of Pediatrics, Hainan Branch of Chinese PLA General Hospital, Haitang Bay, Sanya, Hainan Province 572013, China

In this study, we established a bone marrow and bone metastasis xenograft mouse model by injecting the LA-N-5 cell into the bone marrow cavity. Then we examined whether folate-nanoliposomes entrapped MYCN siRNA can specifically distribute to tumor tissues and suppress MYCN gene expression as well as induce apoptosis in neuroblastoma cells.

Materials and methods

Animals

BALB/c nude mice (4 wk) were used for all experiments. All animals were housed in the Experimental Animal Centre of PLA General Hospital, and all procedures were approved by the Institutional Animal Care and Use Committee of PLA General Hospital.

Cell culture

LA-N-5 neuroblastoma cell was the generous gift of Dr. Robert Seeger from Children's Hospital Los Angeles, and the cells were grown in RPMI 1640 medium (GIBCO BRL, USA) containing 15% fetal bovine serum (Hyclone, USA), 2 mmol/L l-glutamine, 100 IU/ml penicillin G and 100 µg/mL streptomycin (all were from Huamei Company, China) in a humidified 5% CO₂ and 95% air atmosphere at 37°C.

Establishment of the bone marrow and bone metastasis xenograft mouse model

Previously described procedure was introduced in the current study [21,22]. Briefly, LA-N-5 neuroblastoma cells were harvested and washed by centrifugation and resuspended in cold serum-free medium prior to injection. Mice were immobilized in a restraining device, and cells (10⁵ cells in 5 µl) were injected into the bone marrow cavity of the femoral proximal metaphysis. About five weeks later, the diameters of tumors can be up to 5 mm and the bone metastasis model is successfully established.

Preparation of the folate-nanoliposome entrapped MYCN siRNA

The folate-nanoliposome entrapped MYCN siRNA with Cy-3 fluorescence tag was the generous gift from Dr. Robert J. Lee (Ohio State University, College of Pharmacy). siRNA sequences were as follows: MYCN siRNA sense: 5'-CGGAGAUGCUGCUUGAGAA dtdt-3', anti-sense: 5'-UUCUCAAGCAGCAUCUCCG dtdt-3'; the negative control siRNA: sense: 5'-UUCUCCGAACGU GUACGU dtdt-3', anti-sense: 5'-ACGUGACAC GUU CGGAGAA dtdt-3'.

siRNA distribution analysis and gene therapy

For siRNA distribution analysis, LA-N-5 neuroblastoma cells were injected into mice bone marrow cavity to

establish the bone marrow and bone metastasis xenograft models. Then the folate-nanoliposome entrapped MYCN siRNA with Cy3 tag (3 mg/kg) was injected into mice via lateral tail vein, and mice were sacrificed 8 hours after injection. Tumors of femur and other indicated organs (heart, lung, liver, kidney) were harvested, embedded in optimum cutting temperature compound (OCT) and sectioned into 4 µm frozen sections. Tissues sections were observed under fluorescence microscopy (stimulating wavelength, 554 nm). Three views of each organ were chosen randomly and subjected to calculate the average integrated optical density (IOD) with Image pro plus 6.0 Software. In the gene therapy experiment, mice were divided into two groups (6 mice for each group). LA-N-5 neuroblastoma cells were injected into mice bone marrow cavity to establish the bone marrow and bone metastasis xenograft model. Then the folate-nanoliposome entrapped MYCN siRNA with Cy3 tag (3 mg/kg/d) were injected into one group of mice by lateral tail vein. The other group of mice were injected with folate-nanoliposome entrapped the control siRNA (3 mg/kg/d). Mice were sacrificed at the sixth day after 5 injections of folate-nanoliposome entrapped siRNAs. Tumors frozen sections were analyzed by TUNEL Assay.

Quantitative RT-PCR analysis

The cells were harvested and RNA was extracted by using the Ultrapure RNA Kit (CWbio.Co.Ltd). Briefly, cells were lysed in RLT buffer and RNA was purified according to the manufacturer's instructions. RNase-free DNase set (CWbio.Co.Ltd) was used to remove any contaminating genomic DNA. The cDNA was synthesized using HiFi-MMLV cDNA Kit (CWbio.Co.Ltd). The primer sequences for MYCN were: 5'-CTCAGTACCTCC GGAGAG-3' (sense) and 5'-GGCATCGTIGAGGATC-3' (antisense). The primer sequences for GAPDH were: 5'-TGCACCACCAACTGCTTAGC-3' (sense) and 5'-GGCATGGACTGTGGTCATGAG-3' (antisense). Real-time PCR was performed on an Roche LightCycler 480II system, using LightCycler 480 SYBR Green I Master (Roche). Each sample was determined in triplicate. MYCN mRNA expression was calculated as MYCN copies/GAPDH copies. Relative standard curve method was used to determine the relative mRNA expression of MYCN gene.

Western blot

Tumor frozen samples from each mouse in a same group were mixed and total protein was extracted. Proteins were separated by a precast 4-12% SDS-PAGE gel (Sigma) and electrophoretically transferred to a PVDF membrane, followed by incubation with anti-β-actin antibody (1/2000 ab3280 Abcam, Cambridge, MA, USA) and anti-n-Myc antibody [NCM II 100] (1:1000 ab16898 Abcam, Cambridge, MA, USA). Protein bands were

detected with ECL reagents (Amersham Biosciences, Buckinghamshire, UK) and visualized by autoradiography.

TUNEL assay

Briefly, frozen tumor samples were cut on a cryostat and got 4 μm -thick sections. Slices were fixed with acetone and soaked in 3% hydrogen peroxide-methanol solution for 20 min, then rinsed in PBS for 3 times. Fixed sections were incubated with proteinase K (20 $\mu\text{g}/\text{ml}$) for 20 min at 37°C and washed with TBS for 3 times. Then they were incubated with TdT buffer containing digoxigenin labelled desoxyuridinetriphosphate (DIG-UTP) and dATP for 2 h at 37°C, and immersed in stop solution for 30 min at room temperature. After washing, the sections were incubated with strept avidin-biotin-peroxidase complex (SABC) antibody solution for 60 min at 37°C, followed by incubation with DAB for 20 min. Slices were counterstained with hematoxylin and examined with a fluorescence microscope.

Statistical analysis

All experiments were performed at least in triplicate and data are expressed as means \pm SEM. Statistical differences were analyzed by two-tailed *t* test. $P < 0.05$ was considered statistically significant.


Results

Folate-nanoliposome entrapped MYCN siRNA is specifically distributed in tumor tissues

LA-N-5 neuroblastoma cells were injected into mice bone marrow cavity to establish the bone marrow and bone metastasis xenograft model. Then mice were injected with the folate-nanoliposome entrapped MYCN siRNA with Cy3 tag (Figure 1) by tail vein, and sacrificed 8 hours after injection. Tissues sections were observed under fluorescence microscopy and showed that Cy3 fluorescence was very bright in tumors in bones (Figure 2A, B). Liver, kidney, heart and lung tissues have weak or no visible fluorescence (Figure 2A, B). This indicates that folate-nanoliposome entrapped MYCN siRNA is specifically distributed in tumor tissues.

In vivo gene silencing effect of folate-nanoliposome entrapped MYCN siRNA in LA-N-5 neuroblastoma

Mice were injected (i.v.) with folate-nanoliposome entrapped MYCN siRNA or control siRNA for 5 days and gene silencing effect of MYCN siRNA in LA-N-5 neuroblastoma were examined by real-time PCR and Western blot. As shown in Figure 3A, MYCN mRNA expression was significant down-regulated (>50%) by folate-nanoliposome entrapped MYCN siRNA compared to negative control siRNA treatment ($p < 0.01$). Western blot result showed MYCN protein expression was about 60% inhibited by






Figure 2 Distribution of folate-nanoliposome entrapped MYCN siRNA: **A**, Cy3 fluorescence of different tissues. **a**, tumors tissue; **b**, heart; **c**, lung; **d**, liver; **e**, kidney. This graph is one representative result of three views; **B**, Quantitative analysis of fluorescence intensity of each tissue. Fluorescence intensity was expressed as integrated optical density (IOD). Results are expressed as means \pm SE.




Figure 3 The in vivo gene silencing effect of folate-nanoliposome entrapped MYCN siRNA in LA-N-5 neuroblastoma: **A**, Quantitative RT-PCR analysis of MYCN gene expression. Results are normalized to GAPDH expression and are presented as fold increases over the control. 6 mice each group, * $P < 0.05$, ** $P < 0.01$; **B**, Western blot analysis of MYCN gene expression. Frozen tumor samples of each mouse in a same group were mixed and analyzed by Western blot. The right graph is a quantitative analysis of the left one.




Figure 4 TUNEL assay of LA-N-5 neuroblastoma: **A**, folate-nanoliposome entrapped control siRNA group; **B**, folate-nanoliposome entrapped MYCN siRNA group.

folate-nanoliposome entrapped MYCN siRNA (Figure 3B). These data demonstrate that systematic injection of folate-nanoliposome entrapped MYCN siRNA can silence MYCN gene expression of tumor tissues efficiently.

The pro-apoptotic effect of folate-nanoliposome entrapped MYCN siRNA in neuroblastoma

Gene therapeutic effect of folate-nanoliposome entrapped MYCN siRNA was evaluated by TUNEL assay of neuroblastoma. As shown in Figure 4, MYCN siRNA treatment group caused much more TUNEL positive cells than control siRNA treatment group. This indicates that folate-nanoliposome entrapped MYCN siRNA treatment can lead to apoptosis of neuroblastoma in vivo.

Discussion

Our results showed MYCN knockdown induces apoptosis of metastatic neuroblastoma in vivo, MYCN gene, located on the short arm of chromosome 2, is only expressed during the stages of embryonic development of the nervous system, kidney, lung and spleen [23]. MYCN gene is considered a proto-oncogene associated with the malignant tumors growth [24-26]. Since MYCN gene amplification plays a key role in the neuroblastoma progression, inhibition the MYCN gene expression becomes a probable strategy for high-risk neuroblastoma therapy. Studies by other groups [27-29] have demonstrated that inhibition of MYCN gene expression in neuroblastoma cells makes cell cycle arrest at G1 phase, and promotes tumor cell apoptosis or differentiation into neural cells. The MYCN gene is a promising drug target for treating peripheral and central nervous system tumors [30].

Although the therapeutic strategy targeting MYCN in vivo by siRNA is attractive, the traditional drugs delivery methods seem to be nonspecific and toxic [31]. The folate receptor-targeted liposome delivery system has emerged recently as a specific and safety delivery method and become the frontier of drugs delivery. This study was designed to examine the specificity and efficacy of folate-nanoliposome entrapped MYCN siRNA in

treatment of metastatic neuroblastoma, showing that MYCN siRNA was distributed specifically in tumor tissues under the direction of folate-folate receptor interaction and further silences MYCN gene expression both in mRNA and protein levels and finally induces apoptosis of neuroblastoma cells. This indicates that folate-nanoliposome entrapped MYCN siRNA could be used in future gene therapy of neuroblastoma. Of note, the in vivo gene knockdown efficiency is up to 60% in protein level, which is much higher than that of traditional methods such as lipofection or calcium-phosphate transfection [31].

This study points to a new way for treatment of metastatic neuroblastoma and widen the application of folate-nanoliposomes delivery system in tumor therapy. The high specificity, efficiency and safety make folate receptor-targeted liposome delivery system a killer mace in siRNAs-mediated cancer gene therapy.

Abbreviations

siRNA: Small interference RNA; IOD: Integrated optical density; DIG-UTP: Digoxigenin labelled desoxyuridinetriphosphate; SABC: Strept avidin-biotin-peroxidase complex.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The work presented here was carried out in collaboration between all authors. ST defined the research theme. QZ designed methods and experiments, carried out the laboratory experiments, CF and WL analyzed the data, interpreted the results. YZ co-worked on associated data collection and their interpretation. All authors read and approved the final manuscript.

Acknowledgements

The work was supported by the National High Technology Research and Development Program of China (863 Program). Fund number: 2012AA020804

Received: 16 March 2013 Accepted: 21 June 2013

Published: 27 June 2013

References

1. Schor NF: Neuroblastoma as a neurobiological disease. *J Neurooncol* 1999, **41**:159-166.
2. Hogarty MD, Maris JM: PI3King On MYCN to improve neuroblastoma therapeutics. *Cancer Cell* 2012, **21**:145-147.

3. Schwab M, Westermann F, Hero B, Berthold F: **Neuroblastoma: biology and molecular and chromosomal pathology.** *Lancet Oncol* 2003, **4**:472–480.
4. Suenaga Y, Kaneko Y, Matsumoto D, Hossain MS, Ozaki T, Nakagawara A: **Positive auto-regulation of MYCN in human neuroblastoma.** *Biochem Biophys Res Commun* 2009, **390**:21–26.
5. Cole MD, McMahon SB: **The Myc oncoprotein: a critical evaluation of transactivation and target gene regulation.** *Oncogene* 1999, **18**:2916.
6. Aagaard L, Rossi JJ: **RNAi therapeutics: principles, prospects and challenges.** *Adv Drug Deliv Rev* 2007, **59**:75–86.
7. Jiang R, Xue S, Jin Z: **Stable knockdown of MYCN by lentivirus-based RNAi inhibits human neuroblastoma cells growth in vitro and in vivo.** *Biochem Biophys Res Commun* 2011, **410**:364–370.
8. Lovén J, Zinin N, Wahlström T, Müller I, Brodin P, Fredlund E, Ribacke U, Pivarcsi A, Pählman S, Henriksson M: **MYCN-regulated microRNAs repress estrogen receptor- α (ESR1) expression and neuronal differentiation in human neuroblastoma.** *Proc Natl Acad Sci* 2010, **107**:1553–1558.
9. Heo I, Kim VN: **Regulating the regulators: posttranslational modifications of RNA silencing factors.** *Cell* 2009, **139**:28–31.
10. Fire A, Xu SQ, Montgomery MK, Kostas SA, Driver SE, Mello CC: **Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*.** *Nature* 1998, **391**:806–811.
11. Talmadge JE, Singh RK, Fidler IJ, Raz A: **Murine models to evaluate novel and conventional therapeutic strategies for cancer.** *Am J Pathol* 2007, **170**:793.
12. Daldrup-Link HE, Hawkins RA, Meier R, Goldsby RE, Artemov D: **Receptor imaging of pediatric tumors: clinical practice and new developments.** *Pediatr Radiol* 2008, **38**:1154–1161.
13. Lee RJ, Low PS: **Delivery of liposomes into cultured KB cells via folate receptor-mediated endocytosis.** *J Biol Chem* 1994, **269**:3198–3204.
14. Rui Y, Wang S, Philip S, Thompson DH: **Diplasmenylcholine-folate liposomes: an efficient vehicle for intracellular drug delivery.** *J Am Chem Soc* 1998, **120**:11213–11218.
15. Gabizon A, Horowitz AT, Goren D, Tzemach D, Mandelbaum-Shavit F, Qazen MM, Zalipsky S: **Targeting folate receptor with folate linked to extremities of poly (ethylene glycol)-grafted liposomes: in vitro studies.** *Bioconjug Chem* 1999, **10**:289–298.
16. Tyagi N, Ghosh PC: **Folate receptor mediated targeted delivery of ricin entrapped into sterically stabilized liposomes to human epidermoid carcinoma (KB) cells: Effect of monensin intercalated into folate-tagged liposomes.** *Eur J Pharm Sci* 2011, **43**:343–353.
17. Leamon CP, Weigl D, Hendren RW: **Folate copolymer-mediated transfection of cultured cells.** *Bioconjug Chem* 1999, **10**:947–957.
18. Reddy JA, Dean D, Kennedy MD, Low PS: **Optimization of folate-conjugated liposomal vectors for folate receptor-mediated gene therapy.** *J Pharm Sci* 1999, **88**:1112–1118.
19. Reddy JA, Low PS: **Enhanced folate receptor mediated gene therapy using a novel pH-sensitive lipid formulation.** *J Control Release* 2000, **64**:27–37.
20. Reddy JA, Low PS: **Enhanced folate receptor mediated gene therapy using a novel pH-sensitive lipid formulation.** *J Control Release* 2000, **64**:27–37.
21. Sohara Y, Shimada H, Scadeng M, Pollack H, Yamada S, Ye W, Reynolds CP, DeClerck YA: **Lytic bone lesions in human neuroblastoma xenograft involve osteoclast recruitment and are inhibited by bisphosphonate.** *Cancer Res* 2003, **63**:3026–3031.
22. Zhao H, Cai W, Li S, Da Z, Sun H, Ma L, Lin Y, Zhi D: **Establishment and characterization of xenograft models of human neuroblastoma bone metastasis.** *Childs Nerv Syst* 2012, **28**:2047–2054.
23. Galderisi U, Di Bernardo G, Cipollaro M, Peluso G, Cascino A, Cotrufo R, Melone MAB: **Differentiation and apoptosis of neuroblastoma cells: role of N-myc gene product.** *J Cell Biochem* 1999, **73**:97–105.
24. Phoenix TN, Gilbertson RJ: **There's a time and a place for MYCN.** *Cancer Cell* 2012, **21**:593–595.
25. Huang R, Cheung NKV, Vider J, Cheung IY, Gerald WL, Tickoo SK, Holland EC, Blasberg RG: **MYCN and MYC regulate tumor proliferation and tumorigenesis directly through BMI1 in human neuroblastomas.** *FASEB J* 2011, **25**:4138–4149.
26. Corvi R, Savelyeva L, Breit S, Wenzel A, Handgretinger R, Barak J, Oren M, Amler L, Schwab M: **Non-syntenic amplification of MDM2 and MYCN in human neuroblastoma.** *Oncogene* 1995, **10**:1081.
27. Kang JH, Rychahou PG, Ishola TA, Qiao J, Evers BM, Chung DH: **MYCN silencing induces differentiation and apoptosis in human neuroblastoma cells.** *Biochem Biophys Res Commun* 2006, **351**:192–197.
28. Nara K, Kusafuka T, Yoneda A, Oue T, Sangkhathat S, Fukuzawa M: **Silencing of MYCN by RNA interference induces growth inhibition, apoptotic activity and cell differentiation in a neuroblastoma cell line with MYCN amplification.** *Int J Oncol* 2007, **30**:1189–1196.
29. Woo CW, Tan F, Cassano H, Lee JH, Lee KC, Thiele CJ: **Use of RNA interference to elucidate the effect of MYCN on cell cycle in neuroblastoma.** *Pediatr Blood Cancer* 2008, **50**:208–212.
30. Bell E, Premkumar R, Carr J, Lu X, Lovat PE, Kees UR, Lunec J, Tweddle DA: **The role of MYCN in the failure of MYCN amplified neuroblastoma cell lines to G.** *Cell Cycle* 2006, **5**:2639–2647.
31. Biewenga JE, Destrée OHJ, Schrama LH: **Plasmid-mediated gene transfer in neurons using the biolistics technique.** *J Neurosci Methods* 1997, **71**:67–75.

doi:10.1186/1475-2867-13-65

Cite this article as: Zhu et al.: Target delivery of MYCN siRNA by folate-nanoliposomes delivery system in a metastatic neuroblastoma model. *Cancer Cell International* 2013 **13**:65.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

