

Broad, broader, broadest

L.X. van Nunen · S.A.W.G. Dello · L.R.C. Dekker

Published online: 8 April 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

Answer

At presentation, the 12-lead electrocardiogram (ECG) showed third-degree AV block with an escape rhythm originating from the anterior fascicle of the left bundle branch (Fig. 1). The subsequent electrocardiograms showed second-degree heart block type Wenckebach, only confined to the left bundle branch (Figs. 2, 3). Within one day, all the conduction disturbances disappeared and the ECG was identical to previous ECGs at regular control visits.

Flecainide is a class IC antiarrhythmic drug that blocks the sodium channel and is known for its ability to prolong the right atrial, atrioventricular, and His-Purkinje conduction time, although it does not slow AV conduction [1]. Nortriptyline has a 'quinidine-like' antiarrhythmic effect and can cause conduction disturbances, amongst which third-degree AV block [2]. Its class IA and IC antiarrhythmic effects may block the cardiac sodium channel and can, especially in combination with other causes of cardiac sodium channel blockage (such as flecainide), lead to a relevant reduction of intracellular sodium.

Moreover, the antiarrhythmic effects of both flecainide and nortriptyline are use-dependent, i.e. requiring repetitive opening and closing of the sodium channels to act as a sodium channel blocker, thus more effective at fast heart rates [3].

In this case, the underlying respiratory tract infection most likely enhanced the antiarrhythmic effect of nortriptyline by causing faster heart rates (up to 130 beats/min in the emergency department), thereby enabling the sodium channel blocking potential of nortriptyline, enhanced by flecainide, which ultimately resulted in third-degree heart block.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Aliot E, Capucci A, Crijns HJ, et al. Twenty-five years in the making: flecainide is safe and effective for the management of atrial fibrillation. *Europace*. 2011;13:161–73.
2. Thanacoody HK, Thomas SH. Tricyclic antidepressant poisoning: cardiovascular toxicity. *Toxicol Rev*. 2005;24:205–14.
3. Bardai A, Amin AS, Blom MT, et al. Sudden cardiac arrest associated with use of a non-cardiac drug that reduces cardiac excitability: evidence from bench, bedside, and community. *Eur Heart J*. 2013;34:1506–16.

L.X. van Nunen (✉) · S.A.W.G. Dello · L.R.C. Dekker
Department of Cardiology, Catharina Hospital Eindhoven,
Michelangelolaan 2,
5623 EJ Eindhoven, The Netherlands
e-mail: lokien.v.nunen@catharinaziekenhuis.nl