

# A sharp end to sugary Wingless travels

Ilia A. Droujinine,<sup>1</sup> Dong Yan,<sup>1</sup> and Norbert Perrimon<sup>1,2</sup>

<sup>1</sup>Department of Genetics and <sup>2</sup>Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115

*Drosophila melanogaster* follicle stem cells are controlled by Wingless (Wg) ligands secreted 50  $\mu\text{m}$  away, raising the question of how long-distance Wg spreading occurs. In this issue of *JCB*, Wang and Page-McCaw (2014. *J. Cell Biol.* <http://dx.doi.org/10.1083/jcb.201403084>) demonstrate a potential mechanism by which the heparan sulfate proteoglycan Dally-like (Dlp) promotes Wg travel, whereas matrix Mmp2 (Metalloproteinase 2) impedes it by inactivating Dlp.

Tissues are maintained and patterned by stem cells that are controlled in part by signals derived from their niches (Losick et al., 2011). Follicle stem cells (FSCs), located in the germaria of each ovariole in *Drosophila melanogaster* ovaries, give rise to the epithelium that surrounds the egg chambers (Losick et al., 2011). FSCs are regulated by several signaling pathways, including Wingless (Wg), derived from the distal ( $\leq 50 \mu\text{m}$ ) terminal filaments (TFs) and cap niche cells (Fig. 1; Losick et al., 2011). Because this signaling is long range, an unresolved issue is how Wg molecules spread. In this issue of *JCB*, Wang and Page-McCaw provide new insights into this process by identifying the heparan sulfate proteoglycan (HSPG) Dally-like (Dlp) and the matrix metalloproteinase Mmp2 as positive and negative regulators of long-range Wg signaling in the germarium, respectively.

In the *Drosophila* wing imaginal disc, Wg has been proposed to act as a morphogen, and a Wg gradient can be detected 50  $\mu\text{m}$  from the source (Strigini and Cohen, 2000). The spreading of Wg in the wing disc requires the glypican Dlp that binds Wg and promotes Wg signaling in distal cells (Baeg et al., 2001, 2004; Kirkpatrick et al., 2004; Kreuger et al., 2004; Franch-Marro et al., 2005; Han et al., 2005; Yan et al., 2009). In the germarium, Wang and Page-McCaw (2014) find that Wg forms a gradient with highest concentrations at the cap/TF cells, whereas Dlp forms an inverse pattern with higher levels closer to the FSCs. They show that *Dlp* loss of function led to a reduction in extracellular Wg level, Wg signaling activity, and FSC proliferation, suggesting that, in the germarium as in the wing disc, Dlp is involved in retaining Wg at the cell surface and preventing its degradation.

In contrast, the authors found that extracellular Wg level and signaling and FSC proliferation (number of stalk cells between follicles, phospho-histone H3 staining, and mitotic clone frequency) are increased in *Mmp2* mutant germaria. Matrix

metalloproteinases (MMPs) are extracellular  $\text{Zn}^{2+}$ -dependent endopeptidases that play pivotal roles in normal tissue remodeling and disease. MMPs have been shown to act on ECM proteins, including collagen, HSPGs, surface molecules, and signaling proteins (Kessenbrock et al., 2010). Mmp2, like Wg, is produced in germarium apical cells. The function of Mmp2 in Wg signaling is likely caused by its regulation of Dlp because Dlp accumulates in *Mmp2* mutant germaria at the TF and mutations in *dlp* suppress the *Mmp2* mutant phenotype.

Previous studies have suggested that Dlp is regulated at multiple layers. For example, in the wing disc, Dlp transcription is modulated by Wg and Hippo signaling (Han et al., 2005; Baena-Lopez et al., 2008), and Notum, a secreted member of  $\alpha/\beta$  hydrolase family, has been shown to cleave Dlp at the level of its glycosylphosphatidylinositol anchor (Kreuger et al., 2004). Wang and Page-McCaw (2014) demonstrate a novel mechanism of Dlp regulation, whereby cleavage of Dlp at its N-terminal domain by Mmp2 causes Dlp to relocate from the cell surface to intracellular vesicles, preventing its interaction with Wg. This finding is of particular interest because the core protein of glypicans, rather than their attached GAG chains, interacts directly with various signaling molecules. For example, the Dlp core protein interacts with Wg and Hedgehog (Hh), whereas the core protein of mammalian glypican-3 binds with high affinity to Sonic Hh (Capurro et al., 2008; Yan et al., 2009, 2010). Moreover, both *Drosophila* and mammalian glypicans are involved in Wnt, Hh, bone morphogenetic protein, FGF, and JAK/STAT (Janus kinase/signal transducer and activator of transcription) pathways (Filmus et al., 2008). Thus, uncovering the regulation of glypicans has a major impact on our understanding of signaling transduction in normal development and tumor progression.

In mammals, as in the fly ovary, important production sites for MMPs are the niche cells (Kessenbrock et al., 2010). Reminiscent of the study by Wang and Page-McCaw (2014), the HSPG syndecan-1 sequesters the chemokine CXCL1; upon lung injury, MMP7 is up-regulated, cleaving syndecan-1 and activating CXCL1, thereby inducing neutrophil migration (Li et al., 2002). MMPs can also cleave insulin growth factor (IGF) binding proteins (Fowlkes et al., 1995) and latent TGF- $\beta$  binding protein (Dallas et al., 2002), releasing active IGF and TGF- $\beta$ , respectively. In addition, MMP3 binds or cleaves

© 2014 Droujinine et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

Correspondence to Norbert Perrimon: [perrimon@receptor.med.harvard.edu](mailto:perrimon@receptor.med.harvard.edu)




Figure 1. **Regulation of FSCs by Mmp2 and the glypican Dlp in *Drosophila* germarium.** Cap cells produce a long-range signal Wg to regulate the behavior of FSCs. Dlp mediates the transport of Wg from the cap cells to the FSCs to promote their proliferation. Dlp and Wg form opposing gradients in the germarium. Mmp2, expressed in the cap and TF cells, cleaves Dlp in its N-terminal domain and relocates Dlp from the cell surface to intracellular vesicles, preventing its interaction with Wg. It remains to be determined what signals regulate Mmp2 activity and what other factors mediate Wg spreading in the germarium. Also, Wg may be locally generated by escort cells (Sahai-Hernandez and Nystul, 2013). GSC, germline stem cell.

Wnt5b, a Wnt signaling inhibitor, increasing mammary stem cell function (Kessenbrock et al., 2013). Therefore, the work by Wang and Page-McCaw (2014) is relevant to mammalian systems in which HSPGs and MMPs act on multiple signaling pathways (Filmus et al., 2008; Kessenbrock et al., 2010).

The study by Wang and Page-McCaw (2014) raises several questions. First, is Dlp cleavage by Mmp2 required in vivo (only in vitro data were shown)? Second, given the evidence from mammals and *Drosophila* that HSPGs and/or MMPs affect numerous secreted factors (Filmus et al., 2008; Kessenbrock et al., 2010; Wang et al., 2010), does Mmp2 or Dlp act on other signaling pathways to affect FSCs or other cells or do they primarily act through Wg? Third, MMP activity is known to be regulated by proteinases, inhibitors, reactive oxygen species, localization, ECM stiffness, and signaling pathways (NF- $\kappa$ B, FGF, and leptin; Kessenbrock et al., 2010; Wang et al., 2010). Is Mmp2 activated by these or other signals (e.g., nutrition and systemic factors)? Fourth, what are the roles of Mmp2–Dlp interactions in other tissues? Fifth, is Wg spreading in the ovary dependent on other Wg binding factors, such as Swim, Wntless, Lipophorin, or others (Mulligan et al., 2012)? Sixth, it has been suggested that Wg may be produced by FSC-neighboring escort cells (Sahai-Hernandez and Nystul, 2013). As a membrane-tethered form of Wg can replace the endogenous Wg protein in the wing disc (Alexandre et al., 2014), it will be interesting to assess long-range Wg signaling in the ovaries of these flies.

In conclusion, Wang and Page-McCaw (2014) demonstrate beautifully the regulation of a signaling factor through proteinase–HSPG interactions. MMPs (Kessenbrock et al., 2010) and HSPGs (Blackhall et al., 2001) are altered in mammalian tumors, raising the question whether they act through similar mechanisms to influence tumor progression.

Submitted: 27 August 2014  
Accepted: 9 September 2014

## References

- Alexandre, C., A. Baena-Lopez, and J.P. Vincent. 2014. Patterning and growth control by membrane-tethered Wingless. *Nature*. 505:180–185. <http://dx.doi.org/10.1038/nature12879>
- Baeg, G.H., X. Lin, N. Khare, S. Baumgartner, and N. Perrimon. 2001. Heparan sulfate proteoglycans are critical for the organization of the extracellular distribution of Wingless. *Development*. 128:87–94.
- Baeg, G.H., E.M. Selva, R.M. Goodman, R. Dasgupta, and N. Perrimon. 2004. The Wingless morphogen gradient is established by the cooperative action of Frizzled and Heparan Sulfate Proteoglycan receptors. *Dev. Biol.* 276:89–100. <http://dx.doi.org/10.1016/j.ydbio.2004.08.023>
- Baena-Lopez, L.A., I. Rodríguez, and A. Baonza. 2008. The tumor suppressor genes dachsous and fat modulate different signalling pathways by regulating dally and dally-like. *Proc. Natl. Acad. Sci. USA*. 105:9645–9650. <http://dx.doi.org/10.1073/pnas.0803747105>
- Blackhall, F.H., C.L. Merry, E.J. Davies, and G.C. Jayson. 2001. Heparan sulfate proteoglycans and cancer. *Br. J. Cancer*. 85:1094–1098. <http://dx.doi.org/10.1054/bjoc.2001.2054>
- Capurro, M.I., P. Xu, W. Shi, F. Li, A. Jia, and J. Filmus. 2008. Glypican-3 inhibits Hedgehog signaling during development by competing with patched for Hedgehog binding. *Dev. Cell*. 14:700–711. <http://dx.doi.org/10.1016/j.devcel.2008.03.006>

- Dallas, S.L., J.L. Rosser, G.R. Mundy, and L.F. Bonewald. 2002. Proteolysis of latent transforming growth factor-beta (TGF-beta)-binding protein-1 by osteoclasts. A cellular mechanism for release of TGF-beta from bone matrix. *J. Biol. Chem.* 277:21352–21360. <http://dx.doi.org/10.1074/jbc.M111663200>
- Filmus, J., M. Capurro, and J. Rast. 2008. Glypicans. *Genome Biol.* 9:224. <http://dx.doi.org/10.1186/gb-2008-9-5-224>
- Fowlkes, J.L., K.M. Thraikill, D.M. Serra, K. Suzuki, and H. Nagase. 1995. Matrix metalloproteinases as insulin-like growth factor binding protein-degrading proteinases. *Prog. Growth Factor Res.* 6:255–263. [http://dx.doi.org/10.1016/0955-2235\(95\)00017-8](http://dx.doi.org/10.1016/0955-2235(95)00017-8)
- Franch-Marro, X., O. Marchand, E. Piddini, S. Ricardo, C. Alexandre, and J.P. Vincent. 2005. Glypicans shunt the Wingless signal between local signaling and further transport. *Development.* 132:659–666. <http://dx.doi.org/10.1242/dev.01639>
- Han, C., D. Yan, T.Y. Belenkaya, and X. Lin. 2005. *Drosophila* glypicans Dally and Dally-like shape the extracellular Wingless morphogen gradient in the wing disc. *Development.* 132:667–679. <http://dx.doi.org/10.1242/dev.01636>
- Kessenbrock, K., V. Plaks, and Z. Werb. 2010. Matrix metalloproteinases: regulators of the tumor microenvironment. *Cell.* 141:52–67. <http://dx.doi.org/10.1016/j.cell.2010.03.015>
- Kessenbrock, K., G.J. Dijkgraaf, D.A. Lawson, L.E. Littlepage, P. Shahi, U. Pieper, and Z. Werb. 2013. A role for matrix metalloproteinases in regulating mammary stem cell function via the Wnt signaling pathway. *Cell Stem Cell.* 13:300–313. <http://dx.doi.org/10.1016/j.stem.2013.06.005>
- Kirkpatrick, C.A., B.D. Dimitroff, J.M. Rawson, and S.B. Selleck. 2004. Spatial regulation of Wingless morphogen distribution and signaling by Dally-like protein. *Dev. Cell.* 7:513–523. <http://dx.doi.org/10.1016/j.devcel.2004.08.004>
- Kreuger, J., L. Perez, A.J. Giraldez, and S.M. Cohen. 2004. Opposing activities of Dally-like glypican at high and low levels of Wingless morphogen activity. *Dev. Cell.* 7:503–512. <http://dx.doi.org/10.1016/j.devcel.2004.08.005>
- Li, Q., P.W. Park, C.L. Wilson, and W.C. Parks. 2002. Matrilysin shedding of syndecan-1 regulates chemokine mobilization and transepithelial efflux of neutrophils in acute lung injury. *Cell.* 111:635–646. [http://dx.doi.org/10.1016/S0092-8674\(02\)01079-6](http://dx.doi.org/10.1016/S0092-8674(02)01079-6)
- Losick, V.P., L.X. Morris, D.T. Fox, and A. Spradling. 2011. *Drosophila* stem cell niches: a decade of discovery suggests a unified view of stem cell regulation. *Dev. Cell.* 21:159–171. <http://dx.doi.org/10.1016/j.devcel.2011.06.018>
- Mulligan, K.A., C. Fuerer, W. Ching, M. Fish, K. Willert, and R. Nusse. 2012. Secreted Wingless-interacting molecule (Swim) promotes long-range signaling by maintaining Wingless solubility. *Proc. Natl. Acad. Sci. USA.* 109:370–377. <http://dx.doi.org/10.1073/pnas.1119197109>
- Sahai-Hernandez, P., and T.G. Nystul. 2013. A dynamic population of stromal cells contributes to the follicle stem cell niche in the *Drosophila* ovary. *Development.* 140:4490–4498. <http://dx.doi.org/10.1242/dev.098558>
- Strigini, M., and S.M. Cohen. 2000. Wingless gradient formation in the *Drosophila* wing. *Curr. Biol.* 10:293–300. [http://dx.doi.org/10.1016/S0960-9822\(00\)00378-X](http://dx.doi.org/10.1016/S0960-9822(00)00378-X)
- Wang, Q., M. Uhlirova, and D. Bohmann. 2010. Spatial restriction of FGF signaling by a matrix metalloprotease controls branching morphogenesis. *Dev. Cell.* 18:157–164. <http://dx.doi.org/10.1016/j.devcel.2009.11.004>
- Wang, X., and A. Page-McCaw. 2014. A matrix metalloproteinase mediates long-distance attenuation of stem cell proliferation. *J. Cell Biol.* 206:923–936.
- Yan, D., Y. Wu, Y. Feng, S.C. Lin, and X. Lin. 2009. The core protein of glypican Dally-like determines its biphasic activity in wingless morphogen signaling. *Dev. Cell.* 17:470–481. <http://dx.doi.org/10.1016/j.devcel.2009.09.001>
- Yan, D., Y. Wu, Y. Yang, T.Y. Belenkaya, X. Tang, and X. Lin. 2010. The cell-surface proteins Dally-like and Ihog differentially regulate Hedgehog signaling strength and range during development. *Development.* 137:2033–2044. <http://dx.doi.org/10.1242/dev.045740>