


# Use of gene expression profiling to identify candidate genes for pretherapeutic patient classification in acute appendicitis

N. Kiss<sup>1</sup>, M. Minderjahn<sup>1</sup>, J. Reismann<sup>1</sup>, J. Svensson<sup>2</sup>, T. Wester<sup>2</sup>, K. Hauptmann<sup>3</sup>, M. Schad<sup>4</sup>, J. Kallarackal<sup>4</sup>, H. von Bernuth<sup>5</sup> and M. Reismann <sup>1,\*</sup>

<sup>1</sup>Department of Paediatric Surgery, Charité – Universitätsmedizin Berlin, Berlin, Germany

<sup>2</sup>Department of Paediatric Surgery, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden

<sup>3</sup>Institute of Pathology, Charité – Universitätsmedizin Berlin, Berlin, Germany

<sup>4</sup>OakLabs, Hennigsdorf, Germany

<sup>5</sup>Department of Paediatric Pulmonology and Immunology, Charité – Universitätsmedizin Berlin, Berlin, Germany

\*Correspondence to: Department of Paediatric Surgery, Charité – Universitätsmedizin Berlin, Augustenburger Platz 1, 13353 Berlin, Germany (e-mail: marc.reismann@charite.de)

## Abstract

**Background:** Phlegmonous and gangrenous appendicitis represent independent pathophysiological entities with different clinical courses ranging from spontaneous resolution to septic disease. However, reliable predictive methods for these clinical phenotypes have not yet been established. In an attempt to provide pathophysiological insights into the matter, a genomewide gene expression analysis was undertaken in patients with acute appendicitis.

**Methods:** Peripheral blood mononuclear cells were isolated and, after histological confirmation of PA or GA, analysed for genomewide gene expression profiling using RNA microarray technology and subsequent pathway analysis.

**Results:** Samples from 29 patients aged 7–17 years were included. Genomewide gene expression analysis was performed on 13 samples of phlegmonous and 16 of gangrenous appendicitis. From a total of 56 666 genes, 3594 were significantly differently expressed. Distinct interaction between T and B cells in the phlegmonous appendicitis group was suggested by overexpression of T cell receptor  $\alpha$  and  $\beta$  subunits, CD2, CD3, MHC II, CD40L, and the B cell markers CD72 and CD79, indicating an antiviral mechanism. In the gangrenous appendicitis group, expression of genes delineating antibacterial mechanisms was found.

**Conclusion:** These results provide evidence for different and independent gene expression in phlegmonous and gangrenous appendicitis in general, but also suggest distinct immunological patterns for the respective entities. In particular, the findings are compatible with previous evidence of spontaneous resolution in phlegmonous and progressive disease in gangrenous appendicitis.

## Introduction

Surgeons know how to treat appendicitis: either surgically with appendectomy and supporting measures like abscess drainage, or conservatively without operation<sup>1</sup>. Nevertheless, the optimal treatment depends on the stage or type of inflammation of the vermiform appendix.

The presence of necrosis seems to be the determining factor: necrotizing gangrenous appendicitis (GA) is associated with a complicated clinical course including increased rates of peritonitis and abscess formation, even if perforation is not present, whereas non-necrotizing, phlegmonous appendicitis (PA) usually seems to be self-limiting<sup>1–3</sup>. Further insights into epidemiological aspects have shown that perforated and non-perforated appendicitis are characterized by independent frequencies within defined populations<sup>4</sup>. Histologically, perforated GA, referred to clinically as complicated appendicitis, seems to evolve much faster than previously thought, with increased risk of abscess formation up

to septic disease, and can rarely be prevented. In contrast, uncomplicated, non-perforated inflammation appears to be self-limiting, with spontaneously decreasing inflammatory values in a large proportion of those affected<sup>5</sup>. These epidemiological data are supported by evidence from an RCT<sup>5</sup> comparing treatment results for patients who had been diagnosed with uncomplicated appendicitis, and were treated conservatively with or without antibiotics. Rates of treatment failure were comparable in the two arms (79.3 versus 76.6 per cent). Further support for the hypothesis of independent inflammatory entities is provided by immunological evidence. PA and GA differ significantly in terms of distinct underlying immunological patterns; there are primarily Th2-helper cell-dependent mechanisms in those with PA and Th1/Th17-dependent mechanisms in patients with gangrenous inflammation<sup>6,7</sup>.

As the inflammatory entity seemingly determines the clinical course, these types of appendicitis require different therapeutic

Received: August 18, 2020. Revised: October 6, 2020. Accepted: October 24, 2020

© The Author(s) 2021. Published by Oxford University Press on behalf of BJS Society Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

courses of action. Therefore, decisive diagnostics are mandatory to reliably differentiate complicated from uncomplicated appendicitis.

Clinical investigation and laboratory values are basic methods used in the diagnosis of acute appendicitis<sup>8</sup>. In an analysis of the use of routine laboratory values (full blood count and C-reactive protein) for pretherapeutic prediction of PA and GA, the parameters tested did not provide a decisive distinction between the two, although steady pathophysiological differences between the histopathological entities were demonstrated<sup>9</sup>.

Currently, imaging techniques such as CT seem to provide good capacity for differentiation between complicated and uncomplicated appendicitis<sup>10</sup>. In a study<sup>11</sup> focused on the value of radiation-free sonography for differentiation of the inflammatory types in children, abdominal ultrasonography proved to be a valuable tool for the diagnosis of appendicitis, but was limited in terms of differentiation of the two entities. Application of artificial intelligence and methods of machine learning to the laboratory parameters and sonographically measured appendiceal diameters led to substantially improved capacity for the discrimination of PA and GA, and thereby computer algorithm-based biomarker signatures were established<sup>12</sup>.

This comprehensive prospective whole-genome gene expression study essentially encompasses two objectives. First, the aim was to investigate the general extent of gene expression differences in children with PA and GA. Second, the entities were examined at a basic level of gene expression that would allow description of specific pathophysiological and, particularly, immunological differences in terms of pathways. Interesting differences between PA and GA have already been described at the genomic level<sup>13</sup>. The present description of dynamic gene expression level with subsequent pathway analysis for differential diagnosis of acute appendicitis is a novel approach.

## Methods

This was a single-centre, prospective study of patients aged 7–17 years who had undergone surgery for acute appendicitis at the Department of Paediatric Surgery of Charité – Universitätsmedizin Berlin, between April and August 2019. The specific age group in this paediatric surgical population was determined as a necessary level of ability to consent and reading comprehension was required in order to be able to give informed consent to participate in the study. The study was approved for the limited number of patients (30 + expected exclusion rate of 10 per cent) by the institutional ethics committee (reference number ES2/130/16), with the perspective of a subsequent sample size calculation for a larger study.

Inclusion criteria for the study were: patients aged 7–17 years with a sonographically suspected diagnosis of acute appendicitis and scheduled for appendicectomy<sup>5</sup>. Patients and parents were informed by means of appropriate age-adapted information sheets (children aged 7–14 and 15–17 years; separate leaflets for parents) to obtain informed consent. Patients were specifically asked to report the onset of abdominal pain ( $\pm$  30 min). The interval between the start of symptoms and blood sample collection was calculated accordingly.

Exclusion criteria were: missing consent; concomitant disease; previous or present specific treatment for acute appendicitis (conservative or surgical); antibiotic treatment within the past 2 weeks; interval between blood sample collection and isolation of peripheral blood mononuclear cells (PBMCs) more than 1 h;

insufficient RNA quality (RNA integrity number (RIN) below 7); and appendicectomy not performed (no histopathological specimen).

In strict accordance with the ethics committee's approval, blood samples (at least 5 ml) were drawn during routine medical laboratory investigations in the paediatric emergency department. The isolation of PBMCs was performed within 1 h after blood sampling. The blood from each individual was suspended in phosphate-buffered saline (PBS) at a 1 : 1 ratio. Density gradient centrifugation was performed using Ficoll<sup>®</sup> PM400 (GE Healthcare, Pittsburgh, Pennsylvania, USA) at room temperature for 30 min at 400g. After isolation of the monocyte layer, the cells were resuspended in PBS and centrifuged twice for 5 min at 400g. After final resuspension in 1 ml PBS, centrifugation and removal of the supernatant, the native cells were frozen at  $-20^{\circ}\text{C}$  using a sample freezing container (Mr. Frosty<sup>™</sup>, Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA) and finally stored at  $-80^{\circ}\text{C}$  in liquid nitrogen.

## Histopathological examination

After appendicectomy, the resected appendixes were classified histopathologically according to Carr<sup>2</sup>. PA is characterized by transmural infiltration of the appendix by neutrophilic granulocytes, serositis, microabscesses, and oedema, without gangrene or perforation. GA is especially characterized by ischaemic areas in the appendix with transmural myonecrosis. Perforated appendicitis is defined by gangrenous alterations with a transmural defect of the appendix wall. Fig. 1 shows the major histopathological features for group assignment (phlegmonous versus gangrenous) in comparison to normal appendix.


After the primary histopathological examination, a second blinded evaluation was undertaken by a specialized paediatric pathologist.

## Laboratory workflow

Total RNA was isolated from PBMCs using a NucleoSpin<sup>™</sup> RNA Plus kit (Macherey-Nagel, Germany). RNA quality control was achieved using a 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) and the RNA 6000 Pico Kit for total RNA samples. For quantitative control, photometric measurement was performed using a NanoDrop<sup>™</sup> 2000 spectrophotometer (Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA). The quality of each sample was evaluated based on the Bioanalyzer RIN. Samples with a RIN above 7 fulfilled quality control and were subjected to labelling.

For labelling, a Low Input QuickAmp Labeling Kit (Agilent Technologies, Santa Clara, CA, USA) was used to generate fluorescent complementary RNA (cRNA). A random primer-oligo-dT primer mixture was used for first-strand synthesis. After second-strand synthesis, *in vitro* transcription for synthesis of cRNA labelled with cyanine 3-CTP was performed.

Some 600 ng of each Cy3-labelled cRNA was hybridized at  $65^{\circ}\text{C}$  for 17 h on an OakLabs 8x60K ArrayXS Human Agilent microarray (design ID 79407) containing probes for 56 666 genes, using an Agilent Gene Expression Hybridization Kit (Agilent Technologies, Santa Clara, CA, USA), followed by microarray wash and scanning on a SureScan Microarray Scanner (Agilent Technologies, Santa Clara, CA, USA) at a resolution of 3  $\mu\text{m}$  to generate 20-bit TIF files according to the manufacturer's protocols.



**Fig. 1 Appendiceal histology**

**a** Uninflamed appendix with normal lymphofollicular structures, **b** phlegmonous appendicitis with oedema and diffuse transmural granulocytic infiltration, and **c** gangrenous appendicitis with almost complete wall necrosis and fibrinous-purulent inflammation of serosa and adjacent fatty tissue

## Microarray data analysis

TIF files were extracted using Feature Extraction Software version 11 (Agilent Technologies, Santa Clara, CA, USA) and the GE1\_1105\_Oct12 protocol. The resulting raw data were analysed using DirectArray Software (OakLabs, Hennigsdorf, Germany). The signal distributions of the raw data were visualized using box plots to identify potential issues for individual samples.

Samples were quantile normalized and subjected to statistical analysis (PA versus GA) by applying Welch's test and calculating  $\log_2$  fold changes for each gene.  $P < 0.050$  was considered statistically significant.

## Pathway analysis

Differentially expressed pathways were identified using the generally applicable gene set enrichment method of Luo and co-workers<sup>14</sup> and the gage package in R (R Foundation for Statistical Computing, Vienna, Austria). The following immunologically relevant pathways are referred to in this analysis: Toll-like receptor signalling pathway; antigen processing and presentation; NOD-like receptor signalling pathway; haematopoietic cell lineage; natural killer cell-mediated cytotoxicity; tumour necrosis factor (TNF) signalling pathway and intestinal immune network for IgA production; complement and coagulation pathways; retinoic acid-inducible gene (RIG) I-like receptor signalling pathway; cytosolic DNA-sensing pathway; C-type lectin receptor signalling pathway; T cell receptor signalling pathway; Th1 and Th2 cell differentiation; Th17 cell differentiation; interleukin (IL) 17 signalling pathway; Fc $\epsilon$ RI signalling pathway; Fc $\gamma$ R-mediated phagocytosis; and chemokine signalling pathway.

The significantly differentially expressed genes from the relevant pathways were visualized in a heatmap. For the colour representation, Z-scores were calculated for each line; Z-scores represent differences between a gene's normalized signal for a sample and the gene's mean signal for all samples divided by the standard deviation. The absolute Z-value represents the distance between a sample's gene signal and the gene's mean signal of all samples in standard deviation units. A sample's gene signal below the mean Z-value indicates a negative (blue) signal, whereas a signal above the mean Z-value indicates a positive (red) signal. Heatmap visualization was combined with hierarchical clustering, whereby samples with the most similar expression profiles—based on normalized expression values—were clustered together. Additionally, clusters were visualised as dendrograms, complementing the heatmaps.

## Results

Thirty-three otherwise healthy patients with sonographic suspicion of acute appendicitis were initially included. RNA quality control led to exclusion of four samples that had signs of degradation: three from patients with GA and one from a patient with PA. After primary routine histopathological examination, 15 specimens were classified as phlegmonous and 14 as gangrenous. After blind histopathological re-evaluation by a specialized paediatric pathologist, two samples from patients with a primary diagnosis of PA were classified as gangrenous inflammation. The PA group eventually consisted of 13 samples and the GA group of 16 (Fig. 2). No statistically significant differences were found with regard to age, sex distribution, and time from onset of symptoms to blood sample collection (Table 1).

## Gene expression analysis

Box plots of raw gene expression signals of the samples finally included and classified showed a similar distribution, with no obviously problematic samples (Fig. S1).

From a total of 56 666 protein-coding and non-coding RNAs analysed, 3594 (6.3 per cent) were significantly differentially expressed between the groups ( $P < 0.050$ ). The distribution of expression patterns ( $\log_2$  fold change versus  $-\log(P)$ ) with respect to significant upregulation and downregulation of genes is illustrated in a volcano plot (Fig. S2).

## Expression patterns within immunological pathways

The following immunological pathways were significantly differentially expressed: Toll and lmg signalling pathway, antigen processing and presentation, NOD-like receptor signalling pathway, haematopoietic cell lineage, natural killer cell-mediated cytotoxicity, TNF signalling pathway and intestinal immune network for IgA production, complement and coagulation pathways, RIG-I-like receptor signalling pathway, Cytosolic DNA-sensing pathway, C-type lectin receptor signalling pathway, T cell receptor signalling pathway, Th1 and Th2 cell differentiation, Th17 cell differentiation, IL-17 signalling pathway, Fc $\epsilon$ RI signalling pathway, Fc $\gamma$ R-mediated phagocytosis, and chemokine signalling pathway. Tables 2 and 3 show the significantly differentially regulated mRNAs. Information about the specific functions of the genes is provided in Table S1. Gene overexpression in the PA group was characterized by lymphoid cell lineage-directed patterns, specifically those involved in T cell/B cell




Fig. 2 Patient flow diagram

Table 1 Distribution of age, sex, and duration of symptoms before sample collection

	Total (n = 29)	Phlegmonous (n = 13)	Gangrenous (n = 16)
Age (years) <sup>†</sup>	11.3(2.6)	11.5(2.7)	11.1(2.6)
Sex ratio (F : M)	14 : 15	4 : 9	10 : 6
Duration of symptoms before sample collection (h) <sup>†</sup>	28.3(16.9)	24.0(13.3)	32.1(19.4)

<sup>†</sup>Values are mean(s.d.).

interactions. In contrast, the expression pattern in the GA group was primarily assigned to an antibacterial function.

Eleven of the significantly increased expressed genes in the PA group concerning  $\alpha$  or  $\beta$  subunits of the T cell receptor were found within the top 100 expressed genes. In the GA group, two significantly increased mRNAs were found among the top 100: SOCS3 and neutrophilic CD64. Based on statistically significant P values within immunological pathways, a heat map for respective differentially expressed genes was created (Fig. 3).

## Discussion

Diagnostic methods provide the framework for the surgical options for treatment of acute appendicitis. The authors' motivation beyond clarification of the pathophysiology was to show what could be expected from gene expression data in the diagnosis of acute appendicitis in the future, especially with regard to one of the most affected age groups. The medium-term goal is to obtain the simplest possible molecular marker providing the most valid information possible for rapid therapeutic decision-making. Such a marker should be safe, objective (without need for interpretation), easy to determine and, in terms of prospective informative value, superior to other diagnostic tools such as imaging, clinical examination, standard laboratory parameters, and combinatorial scores. The latter are especially useful, but they

Table 2 Significantly differentially expressed genes with immunological relevance: relative gene overexpression in patients with phlegmonous appendicitis

Gene	Mean signal PA	Mean signal GA	P <sup>†</sup>
MHC class II	142	65	0.043
HLA-F	2395	1585	0.024
HLA-DOB	185	119	0.017
CD40L <sup>‡</sup>	444	250	0.003
CD2	715	304	0.027
CD3	224	121	0.021
CD24	144	87	0.021
CD23	582	276	0.010
CD72	47	27	0.023
NIK	150	83	0.029
TRAF1	147	93	0.042
CD79 <sup>*</sup>	4384	3148	0.004
Pol-III	66	42	0.037
Interleukin 23	74	30	0.027
TCR $\beta$ variable 2	19	7	0.012
TCR $\beta$ variable 3-1	7	3	0.019
TCR $\beta$ variable 4-2	45	29	0.013
TCR $\beta$ variable 5-5 <sup>*</sup>	7	4	0.002
TCR $\beta$ variable 5-6 <sup>*</sup>	11	5	< 0.001
TCR $\beta$ variable 6	55	27	0.013
TCR $\beta$ variable 6-5	31	11	0.024
TCR $\beta$ variable 6-6	23	9	0.015
TCR $\beta$ variable 6-8 <sup>*</sup>	35	13	0.003
TCR $\beta$ variable 7-4 <sup>*</sup>	17	7	0.008
TCR $\beta$ variable 7-7 <sup>*</sup>	7	2	< 0.001
TCR $\beta$ variable 11-1 <sup>*</sup>	13	6	0.002
TCR $\beta$ variable 11-3 <sup>*</sup>	6	2	0.003
TCR $\beta$ variable 14	23	9	0.045
TCR $\beta$ variable 15	6	2	0.019
TCR $\beta$ variable 18	17	8	0.022
TCR $\beta$ variable 29-1	51	23	0.036
TCR $\alpha$ variable 1-2	19	10	0.016
TCR $\alpha$ variable 2	10	5	0.021
TCR $\alpha$ variable 8-6	15	8	0.042
TCR $\alpha$ variable 10	6	4	0.026
TCR $\alpha$ variable 12-1 <sup>*</sup>	7	3	0.007
TCR $\alpha$ variable 12-2	19	11	0.032
TCR $\alpha$ variable 16	8	3	0.018
TCR $\alpha$ variable 17	17	10	0.044
TCR $\alpha$ variable 19 <sup>†</sup>	23	10	0.004
TCR $\alpha$ variable 22	19	10	0.042
TCR $\alpha$ variable 23	17	8	0.021
TCR $\alpha$ variable 24	9	5	0.019
TCR $\alpha$ variable 26-1	8	6	0.048
TCR $\alpha$ variable 35 <sup>†</sup>	7	3	0.006
TCR $\alpha$ variable 38-2	25	15	0.028
TCR $\alpha$ variable 41 <sup>†</sup>	18	11	0.003

<sup>†</sup>Gene expression within the top 100 differentially expressed genes (based on P values). PA, phlegmonous appendicitis; GA, gangrenous appendicitis.

<sup>‡</sup>Welch's test.

represent complicated constructs with many variables, which have to be interpreted partially<sup>15</sup>.

Overall, 6.3 per cent of the genes analysed in this study were expressed significantly differently between the groups, demonstrating substantial differences among inflammatory entities. For complementary data analysis, the Kyoto Encyclopedia of Genes and Genomes was instrumentalized to map and link the separately expressed genes to relevant molecular pathways. The results of this complex data analysis gave rise to the hypothesis that PA represents an immunological response to viral infections, whereas GA results in an expression pattern related to an antibacterial response.

Most theories generally designate luminal obstruction as the primary origin of acute appendicitis<sup>2</sup>. In contrast, however,



studies have shown faecaliths in up to 27 per cent of all autopsies without inflammatory changes to the appendix, indicating that faecaliths are an incidental rather than a pathophysiologically causal finding in appendicitis specimens<sup>2</sup>. Nevertheless, new insights into the pathophysiology of acute inflammation

**Table 3 Significantly differentially expressed genes with immunological relevance: relative gene overexpression in patients with gangrenous appendicitis**


Gene	Mean signal PA	Mean signal GA	P†
CD11b	325	409	0.029
CD16b	1048	1678	0.032
CD35	437	658	0.022
CD55	1121	1540	0.032
CD36	2159	3046	0.023
TLR5	754	1134	0.011
TLR9	116	134	0.022
NLRP3	135	188	0.034
NLRC4	100	148	0.026
NLRP6	139	167	0.011
Cathelicidin	29	37	0.050
CD64	65	127	0.022
C9	68	87	0.032
SOCS3	1217	2366	0.010
IRAK4	260	327	0.030
Tpl2 kinase	188	324	0.037
IL-17A	196	251	0.048
LCN2	162	276	0.023

PA, phlegmonous appendicitis; GA, gangrenous appendicitis.

†Welch's test.

of the vermiform appendix, which could replace the refuted theories, have been scarce.

In the present study, PA was characterized by lymphoid-directed gene expression patterns. Particularly, overexpression of  $\alpha$  and  $\beta$  subunits of the T cell receptor, MHC class II and CD40L, T cell markers CD2 and CD3, and B-cell markers CD72 and CD79 in patients with PA suggested a specific immunological synapse between T and B cells<sup>16–20</sup>. That seems to be essential for the activation of B cells and the induction of an antiviral humoral response; B cells are reciprocally activated by cytokines for antibody production upon virus internalization, degradation, MHC class II-dependent antigen presentation, and recognition by Th2 cells via the T cell receptor–CD3 complex. Co-stimulation is provided by CD40/CD40L and CD2<sup>19,20,16,17</sup>. Overexpression of  $\alpha$  and  $\beta$  subunits of the T cell receptor in the PA group only represented 11 per cent of the top 100 overexpressed genes. These results confirm previous findings regarding Th2 cell-dependent immunological mechanisms in patients with PA<sup>6,7</sup>. In contrast, patterns associated with myeloid lineage were found in patients with gangrenous inflammation. The monocyte and neutrophil markers CD11b, CD16b, and CD64 point to a primarily antibacterial defence<sup>21</sup>. In particular, the significant upregulation of the bacterial pattern recognition receptors toll-like receptor (TLR) 5 and TLR-9 supports this interpretation<sup>22,23</sup>. Consistent with the authors' theory, low antiviral function is demonstrated by overexpression of suppressor of cytokine signalling (SOCS) 3, which inhibits antiviral cytokine signalling<sup>24</sup>. SOCS3 expression might even explain the relative overexpression of Th17 promoting interleukin 23 in patients with PA; SOCS3 is a potent negative regulator of



**Fig. 3 Heat map visualizing cluster formation of samples with similar immunological expression patterns**

Samples from patients with phlegmonous appendicitis (PA) (GrA) or gangrenous appendicitis (GA) (GrB). Signals above (red) and below (blue) the mean signal are shown.

interleukin 23 expression<sup>25</sup>. Thus, this seeming inconsistency might be explained by downregulation in GA rather than upregulation in PA.

Interestingly, the susceptibility of humans with the functional IL-17A rs2275913 polymorphism for appendicitis has been associated with advanced inflammation<sup>13</sup>. This finding correlates with the present observation of significantly enhanced IL-17A expression in patients with GA. Further to this, the pathway analysis especially contributes to the thesis of independent pathophysiology. The specific functions of genes not mentioned here are tabulated in [Table S1](#).

The study results are in accordance with previous immunological and infectiological investigations regarding acute appendicitis, and the authors' hypothesis that PA and GA represent independent entities. Mean symptom duration and distribution of temporal patterns were comparable in the two groups. These findings are consistent with previous evidence of independent inflammatory courses in patients with PA and GA, without transition from one form to the other<sup>3,4</sup>.

Studies have shown that fetal development of the lymphatic tissue in the appendix is already characterized by persistence of T and B lymphocytes<sup>26</sup>. As a result, maturation of the intestinal immune system does not seem to solely depend on microbial and nutritional antigens. The presence of T and B lymphocytes may be linked to current theories about the basic function of the human vermiform appendix, as it has been suggested to provide a 'safe house' for commensal bacteria in the gut, especially during the course of viral diarrhoea. Subsequently, the commensal bacteria are protected by a biofilm while the composition of the associated lymphatic tissue ensures its antiviral function<sup>27</sup>. Indeed, several viruses, such as coxsackievirus, influenza virus, measles, cytomegalovirus, and adenovirus, have been associated with non-perforating inflammation of the appendix<sup>28,29</sup>.

Microarray-based comparative investigation for pathway analysis has been shown previously to be a highly effective tool<sup>30</sup>. However, despite numerous quality features such as its prospective design, the repeated histopathological assessment, strict quality control of the samples, and ambitious data analysis, the present study has essentially two limitations. Although expectations were met in terms of indicating substantial pathophysiological differences between PA and GA, a larger sample size could have revealed an even more comprehensive picture of the immunological pathways involved, as a large number of nearly statistically significant expression differences were found. Second, the inclusion of healthy patients as a control group could have provided more comprehensive insights.

The results presented here offer valuable information regarding the general description of differential gene expression and identification of specific yet independent immunological gene expression patterns in light of current research. They provide proof of concept for a comprehensive follow-up study with an adequate sample size calculation focusing on gene expression profiling, possibly coupled with high-performance algorithms, to allow successful development of a simple and reliable biomarker to distinguish inflammatory entities within acute appendicitis in childhood. The methods described here could therefore provide a fundamental framework for reliable diagnostics in acute appendicitis at a simple, low-invasive, low-risk, and cost-effective level.

## Funding

HRH Crown Princess Lovisa's Foundation, Stockholm, Sweden.

## Acknowledgements

N.K. and M.M. contributed equally to the manuscript. We acknowledge support from the German Research Foundation (DFG) and the Open Access Publication Fund of Charité – Universitätsmedizin Berlin. The study received funding from HRH Crown Princess Lovisa's Foundation (Stockholm, Sweden). M.S. and J.K. are both owners of OakLabs. They were involved in the study design, gene expression and data analysis, decision to publish, and preparation of the manuscript. Their institution (not them personally) received funding for material costs and employee performance within the study.

**Disclosure.** The authors declare no other conflict of interest.

## Supplementary material

[Supplementary material](#) is available at [BJS Open](#) online.

## References

- Bhangu A, Søreide K, Di Saverio S, Assarsson JH, Drake FT. Acute appendicitis: modern understanding of pathogenesis, diagnosis, and management. *Lancet* 2015;**386**:1278–1287
- Carr NJ. The pathology of acute appendicitis. *Ann Diagn Pathol* 2000;**4**:46–58
- Andersson R, Hugander A, Thulin A, Nyström PO, Olaison G. Indications for operation in suspected appendicitis and incidence of perforation. *BMJ* 1994;**308**:107–110
- Andersson RE. The natural history and traditional management of appendicitis revisited: spontaneous resolution and predominance of prehospital perforations imply that a correct diagnosis is more important than an early diagnosis. *World J Surg* 2007;**31**:86–92
- Park HC, Kim MJ, Lee BH. Randomized clinical trial of antibiotic therapy for uncomplicated appendicitis. *Br J Surg* 2017;**104**:1785–1790
- Rubér M, Berg A, Ekerfelt C, Olaison G, Andersson RE. Different cytokine profiles in patients with a history of gangrenous or phlegmonous appendicitis. *Clin Exp Immunol* 2006;**143**:117–124
- Rubér M, Andersson M, Petersson BF, Olaison G, Andersson RE, Ekerfeldt C. Systemic Th17-like cytokine pattern in gangrenous appendicitis but not in phlegmonous appendicitis. *Surgery* 2010;**147**:366–372
- Andersson RE. Meta-analysis of the clinical and laboratory diagnosis of appendicitis. *Br J Surg* 2004;**91**:28–37
- Minderjahn MI, Schädlich D, Radtke J, Rothe K, Reismann M. Phlegmonous appendicitis in children is characterized by eosinophilia in white blood cell counts. *World J Pediatr* 2018;**14**:504–509
- Kim HY, Park JH, Lee SS, Lee WJ, Ko Y, Andersson RE *et al*. CT in differentiating complicated from uncomplicated appendicitis: presence of any of 10 CT features versus radiologists' gestalt assessment. *AJR Am J Roentgenol* 2019;**213**:W218–W227
- Rawolle T, Reismann M, Minderjahn MI, Bassir C, Hauptmann K, Rothe K *et al*. Sonographic differentiation of complicated from uncomplicated appendicitis. *Br J Radiol* 2019;**92**:20190102
- Reismann J, Romualdi A, Kiss N, Minderjahn MI, Kallarakal J, Schad M *et al*. Diagnosis and classification of pediatric acute appendicitis by artificial intelligence methods: an investigator-independent approach. *PLoS One* 2019;**14**:e0222030
- Dimberg J, Rubér M, Skarstedt M, Andersson M, Andersson RE. Genetic polymorphism patterns suggest a genetic driven

- inflammatory response as pathogenesis in appendicitis. *Int J Colorectal Dis* 2020;**35**:277–284
14. Luo W, Friedman MS, Shedden K, Hankenson KD, Woolf PJ. GAGE: Generally applicable gene set enrichment for pathway analysis. *BMC Bioinformatics* 2009;**10**:161
  15. Atema JJ, van Rossem CC, Leeuwenburgh MM, Stoker J, Boermeester MA. Scoring system to distinguish uncomplicated from complicated acute appendicitis. *Br J Surg* 2015;**102**:979–990
  16. Szomolanyi-Tsuda E, Brien JD, Dorgan JE, Welsh RM, Garcea RL. The role of CD40–CD154 interaction in antiviral T cell-independent IgG responses. *J Immunol* 2000;**164**:5877–5882
  17. Clevers H, Alarcon B, Wileman T, Terhorst C. The T cell receptor/CD3 complex: a dynamic protein ensemble. *Annu Rev Immunol* 1988;**6**:629–662
  18. Oxenius A, Campbell KA, Maliszewski CR, Kishimoto T, Kikutani H, Hengartner H, Zinkernagel RM, Bachmann MF. CD40–CD40 ligand interactions are critical in T-B cooperation but not for other anti-viral CD4+ T cell functions. *J Exp Med* 1996;**183**:2209–18. doi: 10.1084/jem.183.5.2209
  19. Adachi T, Wakabayashi C, Nakayama T, Yakura H, Tsubata T. CD72 negatively regulates signaling through the antigen receptor of B cells. *J Immunol* 2000;**164**:1223–1229
  20. Chu PG, Arber DA. CD79: a review. *Appl Immunohistochem Mol Morphol* 2001;**9**:97–106
  21. Mortaz E, Alipoor SD, Adcock IM, Mumby S, Koenderman L. Update on neutrophil function in severe inflammation. *Front Immunol* 2018;**9**:2171
  22. Yoon SI, Kurnasov O, Natarajan V, Hong M, Gudkov AV, Osterman AL et al. Structural basis of TLR5-flagellin recognition and signaling. *Science* 2012;**335**:859–864
  23. El Kebir D, József L, Pan W, Wang L, Filep JG. Bacterial DNA activates endothelial cells and promotes neutrophil adherence through TLR9 signaling. *J Immunol* 2009;**182**:4386–4394
  24. Song MM, Shuai K. The suppressor of cytokine signaling (SOCS) 1 and SOCS3 but not SOCS2 proteins inhibit interferon-mediated antiviral and antiproliferative activities. *J Biol Chem* 1998;**273**:35 056–35 062
  25. Chen Z, Laurence A, Kanno Y, Pacher-Zavisin M, Zhu B-M, Tato C et al. Selective regulatory function of Socs3 in the formation of IL-17-secreting T cells. *Proc Natl Acad Sci USA* 2006;**103**:8137–8142
  26. Bhide SA, Wadekar KV, Koushik SA. Peyer's patches are precocious to the appendix in human development. *Dev Immunol* 2001;**8**:159–166
  27. Laurin M, Everett M, Lou, Parker W. The cecal appendix: one more immune component with a function disturbed by post-industrial culture. *Anat Rec* 2011;**294**:567–579
  28. Alder AC, Fomby TB, Woodward WA, Haley RW, Sarosi G, Livingston EH. Association of viral infection and appendicitis. *Arch Surg* 2010;**145**:63–71
  29. Lamps LW. Infectious causes of appendicitis. *Infect Dis Clin North Am* 2010;**24**:995–1018
  30. Tang BMP, McLean AS, Dawes IW, Huang SJ, Lin RCY. The use of gene-expression profiling to identify candidate genes in human sepsis. *Am J Respir Crit Care Med* 2007;**176**:676–684