

POSTER PRESENTATION

Open Access

Endothelial dysfunction in a mouse model of human neutral lipid storage disease

Astrid Schrammel^{1*}, Marion Mussbacher¹, Gerald Wölkart¹, Heike Stessel¹, Günter Hämmerle², Wael Al Zoughbi³, Gerald Höfler³, Alois Lametschwandtner⁴, Rudolf Zechner², Bernd Mayer¹

From 6th International Conference on cGMP: Generators, Effectors and Therapeutic Implications Erfurt, Germany. 28-30 June 2013


Background

Systemic knockout of adipose triglyceride lipase (ATGL), the rate-limiting enzyme of triglyceride catabolism, results in a murine phenotype characterized by progressive accumulation of lipids in the heart finally leading to lethal cardiac dysfunction. Since cardiac and vascular dysfunction are closely related we investigated endothelium-dependent and -independent vessel function of ATGL knockout (ATGL^{-/-}) mice. Using mice with cardiomyocyte-restricted overexpression of ATGL (cardiac-rescued phenotype;

ATGL^{-/-}/MHC-A35) we were able to differentiate between heart-related and -unrelated effects.

Results

Aortic relaxation studies and Langendorff perfusion experiments of isolated hearts demonstrated that ATGL^{-/-} mice suffer from pronounced micro- and macrovascular endothelial dysfunction. Experiments with DEA/NO revealed the functional integrity of the smooth muscle cell layer. Since loss in vascular reactivity was restored by ~50 % in cardiac-rescued mice, this phenomenon seems partly a consequence of impaired cardiac contractility. Biochemical analysis revealed that aortic eNOS protein was down-regulated by more than 60% in aortas of ATGL^{-/-} mice. As consequence, phosphorylation of VASP at Ser 239 was almost abolished. Both parameters were



* Correspondence: astrid.schrammel-gorren@uni-graz.at

¹Department of Pharmacology and Toxicology, Institute of Pharmaceutical Sciences, University of Graz, 8010 Graz, Austria

Full list of author information is available at the end of the article

totally reversed in vessels of cardiac rescued mice. Aortic expression of eNOS mRNA was not affected in ATGL deficiency excluding a transcriptional mechanism underlying the observed effect. Total levels of ubiquitinated proteins (a measure of vascular proteasomal activity) were down-regulated in ATGL-deficient aortas (~26% compared to WT controls) and fully recovered upon cardiomyocyte-restricted overexpression of ATGL.

Conclusion

Endothelial dysfunction in murine ATGL deficiency partly arises from impaired cardiac contractility and originates from down-regulation of aortic eNOS presumably due to activation of the vascular proteasome. Potential heart-independent mechanisms contributing to the observed defect are currently investigated.

Authors' details

¹Department of Pharmacology and Toxicology, Institute of Pharmaceutical Sciences, University of Graz, 8010 Graz, Austria. ²Department of Molecular Bioscience, University of Graz, 8010 Graz, Austria. ³Institute of Pathology, Medical University of Graz, Auenbruggerplatz 25, 8010 Graz, Austria. ⁴Department of Cell Biology and Physiology, Vessel and Muscle Research Unit, University of Salzburg, 5020 Salzburg, Austria.

Published: 29 August 2013

doi:10.1186/2050-6511-14-S1-P64

Cite this article as: Schrammel *et al.*: Endothelial dysfunction in a mouse model of human neutral lipid storage disease. *BMC Pharmacology and Toxicology* 2013 **14**(Suppl 1):P64.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

