



## Surgical Technique

## Combating the Opioid Epidemic: Experience with a Single Prescription for Total Joint Arthroplasty

Daniel Sun, MD<sup>a</sup>, Antonio Cusano, MD<sup>b</sup>, Melvin A. Harrington, MD<sup>a</sup>, Mohamad J. Halawi, MD<sup>a,\*</sup><sup>a</sup> Department of Orthopaedic Surgery, Baylor College of Medicine, Houston, TX, USA<sup>b</sup> Department of Orthopaedic Surgery, University of Connecticut Health Center, Farmington, CT, USA

## ARTICLE INFO

## Article history:

Received 13 April 2020

Received in revised form

16 June 2020

Accepted 10 July 2020

Available online xxx

## Keywords:

Opioids

Hip

Knee: primary

Revision

Arthroplasty

## ABSTRACT

**Background:** Despite advances in perioperative total joint arthroplasty (TJA) pain protocols, opiates continue to play a major role in postoperative pain control. This brief communication reports our experience with a restrictive opioid protocol allowing patients only a single prescription of low-dose opioids.

**Methods:** One hundred consecutive elective, primary, and revision TJAs were analyzed. All patients received preoperative counseling and multimodal analgesia. Counseling involved discussion of patient expectations on postoperative pain management, weaning off opioids before surgery, and emphasis that opioid refills were not permitted. Ninety-day outcomes including pain-related phone calls, opioid refill requests, emergency room visits, complications, and readmissions were assessed. Opioid dispensing was tracked using our state prescription monitoring program.

**Results:** There was a high prevalence of preoperative opioid use, depression, and anxiety (25%, 34%, and 39%, respectively). Sixty-eight percent of chronic opioid users were able to wean off opioids before surgery. The average initial prescription of opioids was equivalent to 48 pills of 5 mg oxycodone. There were only 10 pain-related phone calls from 9 patients; all were using opioids preoperatively, with only one patient requesting a refill. All pain-related phone calls occurred in the first week after surgery. There were no emergency room visits, complications, or readmissions related to pain.

**Conclusions:** A single prescription of low-dose opioids was sufficient for patients undergoing TJA when using preoperative patient preparation and multimodal analgesia. Standardized guidelines are needed to guide best practices for patient education and pain management, especially in patients on chronic opioid therapy. This information will help implement evidence-based strategies to accelerate the decline of opioid use and hopefully pave the way for opioid-free TJA.

© 2020 The Authors. Published by Elsevier Inc. on behalf of The American Association of Hip and Knee Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The United States is the largest consumer of opioids worldwide with more than 200 million prescriptions dispensed annually [1–3]. Opioids have traditionally played a major role in the management of postoperative pain for total joint arthroplasty (TJA) [4]. Apart from concerns for potential abuse [5,6], there is mounting evidence

linking prolonged opioid use with detrimental effects after TJA. Patients on chronic preoperative opioid therapy are likely to experience extended hospital length of stay (LOS), persistent pain, higher complication and reoperation rates, and decreased satisfaction [7–10]. Furthermore, patients receiving chronic opioids before TJA are at a higher risk for prolonged opioid use and overdose postoperatively [7–10].

Over the past few years, the perioperative management of patients undergoing TJA has witnessed tremendous improvements. Preoperative preparation including patient education and medical optimization, multimodal analgesia including perioperative infiltration and nerve blocks, blood conservation techniques including tranexamic acid and spinal anesthesia, and minimally invasive

\* Corresponding author. Department of Orthopaedic Surgery, Baylor College of Medicine, 7200 Cambridge Street, Suite 10A, Houston, TX 77030, USA. Tel.: +1 713 986 1606.

E-mail address: [mohamad.halawi@bcm.edu](mailto:mohamad.halawi@bcm.edu)

surgical approaches have collectively led to improved pain control, shorter hospitalization, enhanced recovery, and decreased consumption of opioids postoperatively [11–13]. These practices have been further reinforced by the growing shift toward outpatient surgery and bundled payment models.

The purpose of this study was to report on our experience using a restrictive opioid-prescribing protocol consisting of only 1-week supply with no refills. A consecutive series of primary and revision TJAs was reviewed. Ninety-day outcomes including pain-related phone calls, opioid refill requests, emergency room (ER) visits, complications, and readmissions were assessed.

## Materials and methods

Institutional review board approval was obtained. In fall 2017, the senior author started a new arthroplasty position in a state with restrictive opioid regulations. This study was a retrospective review of the first 100 consecutive patients who (1) underwent elective procedure, (2) underwent total hip arthroplasty (THA) and total knee arthroplasty (TKA) procedures by the senior author, and (3) were admitted to him. Primary and revision cases were included. All patients had 90-day follow-up data.

Before surgical scheduling, all patients received counseling by the surgeon. Specific details of the surgical procedure, risks, benefits, hospitalization, recovery, pain control, rehabilitation, and expectations were presented. Patients on chronic opioids were advised to either wean off or obtain a letter from their chronic opioid prescribers that they would continue to assume opioid stewardship postoperatively. Chronic opioid use was defined as receiving opioid medications for at least 3 months with at least one refill at the time of surgical consultation. Weaning off was confirmed by laboratory testing. The state's restrictive laws regarding opioid prescription were outlined. All patients were instructed that no opioid refills are permitted and were warned about their potential adverse effects. Patients were informed that pain is an expected and normal part of the recovery process. The pain control strategy was outlined. Unless contraindicated, pre-emptive analgesia consisting of 200 mg of celecoxib and 1000 mg of acetaminophen was administered in the preoperative holding unit. Spinal anesthesia with intravenous sedation was the preferred technique, but the final decision was made by the anesthesiologist on the day of surgery. A posterolateral approach was used for THAs and a median parapatellar approach for TKAs. A single-shot adductor canal block (administered by either the surgeon or anesthesiologist) was performed. Additionally, ropivacaine-based periarticular infiltration was administered depending on patient's weight. Postoperatively, analgesia consisted of celecoxib (or naproxen) depending on insurance type, acetaminophen, and oxycodone (5mg every 3–4 hours as needed for severe pain for a total of 60 pills). The number of prescribed oxycodone pills was chosen based on our state laws limiting initial opioid prescriptions to a 7-day supply for adults. Some pharmacies dispensed lower quantities depending on their internal guidelines.

Variables collected were the age, gender, body mass index, American Society of Anesthesiologists (ASA) physical classification score, procedure type (primary or revision), history of substance abuse, depression, anxiety, and opioid use within 3 months from decision to proceed with surgery. Operative variables (eg, anesthesia type and blood loss) and initial opioid dose in morphine milligram equivalents were also collected. The outcomes of the study were hospital LOS, 90-day opioid refills (surgeons and outside providers), pain-related phone calls, opioid refill requests, ER visits, and complications. Categorical variables were presented as the frequency and proportion. Continuous values were presented as the mean and standard deviation.

## Results

One hundred consecutive TJA procedures (51 THAs, 7 revision THAs, 34 TKAs, and 8 revision TKAs) were included in the analysis. Twenty-four of the 85 (28%) primary cases were classified as complex procedures (eg, conversion THA/TKA, subtrochanteric osteotomy, and takedown of hip ankylosis). The mean age was  $62.2 \pm 9.5$  years, and the mean body mass index was  $32.5 \pm 6.2$  kg/m<sup>2</sup>. Fifty-three (53%) patients were females, and 47 (47%) were males. Thirty-four (34%) patients had a diagnosis of depression, and 39 (39%) patients had a diagnosis of anxiety. Ten (10%) patients reported a history of substance abuse (heroin and/or alcohol), and 25 (25%) patients were on chronic opioids before surgery. Seventeen of 25 (68%) patients who were on chronic opioid therapy were able to wean off before surgery. General anesthesia was used in 13 (13%) patients, spinal anesthesia in 16 (16%) patients, and combined spinal and general anesthesia in 71 (71%) patients. The mean estimated blood loss was  $194.5 \pm 173.3$  cc. Table 1 summarizes the characteristics of the study group.

The mean LOS was  $0.9 \pm 0.7$  days. The mean initial opioid prescription was  $357.1 \pm 98.5$  morphine milligram equivalents or the equivalent of  $47.6 \pm 13.1$  oxycodone pills (5 mg). Thirty-eight (38%) patients had less than 60 pills filled as some pharmacies followed internal guidelines when determining the number of pills dispensed. Sixteen patients (5 TKAs, 2 revision TKAs, 4 THAs, and 5 revision THAs) did not fill their initial opioid prescription. In the first 90 days after surgery, only one requested an opioid refill. Eight patients received opioid refills by outside nonorthopaedic providers, all of whom were on chronic opioids preoperatively. There were only 10 pain-related phone calls from 9 patients, all occurring within the first week of surgery. Those patients were on chronic opioids preoperatively, and all underwent primary TJA. Table 2 summarizes the outcomes of the study group.

Four (4%) emergency department visits occurred within the first 90 days after surgery. Two patients presented for medical complications (one for pneumonia and the other for decompensated congestive heart failure). Two presented for evaluation of deep venous thrombosis, which was ruled out by a negative sonographic assessment. Four surgical complications occurred within the first 90 days. One patient (revision TKA) presented with wound dehiscence requiring medial gastrocnemius rotational flap coverage, one patient had failure of the arthrotomy repair after forced knee

**Table 1**  
Baseline demographic characteristics of the study group.

Age (years)	62.2 ( $\pm 9.5$ )
The American Society of Anesthesiologists physical classification score	2.3 ( $\pm 0.5$ )
Body mass index (kg/m <sup>2</sup> )	32.5 ( $\pm 6.2$ )
Sex	
Female	53 (54%)
Male	47 (4%)
Procedure	
Primary total hip	51 (51%)
Primary total knee	34 (34%)
Revision total hip	7 (7%)
Revision total knee	8 (8%)
History of substance abuse	10 (10%)
Chronic opioid use	25 (25%)
Depression	34 (34%)
Anxiety	39 (39%)
Anesthesia type	
Combined spinal and general	71 (71%)
General	13 (13%)
Spinal	16 (16%)
Estimated blood loss (cc)	194.5 ( $\pm 173.3$ )

Values given as the mean and standard deviation or the frequency and proportion.

**Table 2**  
Outcomes of the study group.

Hospital length of stay	0.9 ( $\pm$ 0.7)
Initial opioid dose (morphine equivalents)	357.1 ( $\pm$ 198.5)
Initial opioid dose (oxycodone equivalents)	48.1 ( $\pm$ 26.1)
Ninety-day opioid refill by the surgeon	0 (0%)
Ninety-day opioid refill by the nonorthopaedic providers	8 (8%)
Ninety-day pain-related phone calls	10 (10%)
Ninety-day opioid refill requests	1 (1%)
Ninety-day emergency room visits	4 (4%)
Ninety-day complications	4 (4%)

Values given as the mean and standard deviation or the frequency and proportion.

flexion during physical therapy, one patient had instability after revision THA, and one patient succumbed to complications of metastatic lung cancer. Table 3 summarizes the reasons for emergency room visits and complications.

## Discussion

In this study, we reported on our experience with a restrictive opioid-prescribing protocol in a consecutive series of 100 primary and revision TJAs followed for 90 days. Nearly a third of the primary cases were complex procedures, and there was a common prevalence of chronic opioid use, depression, and anxiety. Despite the wide case mix, only one patient requested refill and there were only 10 pain-related phone calls to the clinic during the study period, all occurring in the first week after surgery. Sixteen percent of patients never filled their opioid prescription, and 68% of patients on chronic opioids were able to wean off before surgery. There were no ER visits, complications, or readmissions related to pain.

The mean number of 5-mg oxycodone pills prescribed in our cohort was 48, which is less than previously reported in the literature. Sabatino et al [14] found that their institution prescribed a median of 90 oxycodone pills (range: 10–200 and 20–330 pills for THA and TKA, respectively). Hernandez et al [15] reported an average of 189 oxycodone pills prescribed after primary TKA, with 83% of patients also receiving a prescription for tramadol. A prospective observational study by Premkumar et al [16] found that, on average, 77.7 oxycodone pills were prescribed after primary TKA. A 2018 survey of members of the American Association of Hip and Knee Surgeons found that the average number of pills prescribed for primary THA and TKA was 44 and 49, respectively, although 48.3% of respondents also prescribed tramadol in addition to oxycodone [4].

The restrictive opioid protocol used in this study heavily relies on patient counseling and multimodal analgesia. Preoperative patient counseling has been shown to positively influence pain ratings, kinesiophobia, and patient satisfaction [17]. Patient reassurance that pain is a normal part of recovery process combined with a clear communication that no medication refills would be provided may explain our ability to prescribe low amounts of

opioids at discharge while maintaining a low rate of pain-related phone calls. This is despite the fact that the rates of anxiety, depression, substance abuse, and chronic opioid use in our study population were higher than those in other TJA cohorts [6,18]. In addition, previous studies have reported opioid refill rates as high as 67% for primary TJA [14] and a 23% pain-related phone call rate in the first 7 postoperative days [19]. In contrast, we were able to adhere to the target of zero refills and our pain-related phone call rate was only 10% despite including revision cases and extending follow-up to 90 days.

Another feature of the opioid protocol used in this study was it allowed for shared decision-making with patients on chronic opioids to either wean off or submit a letter from their original prescribers attesting that they will continue to assume opioid stewardship postoperatively. It was positive to find that a majority of patients were able to completely wean off and not require any refills postoperatively. The findings of this study are encouraging as we seek to tackle the opioid epidemic. Encouraging patients to wean off preoperatively may lead to decreased postoperative utilization and is an important intervention where surgeons can make a tremendous impact, not just clinically but also at a societal level. Brock et al [18] found that cessation of opioids 3 months before surgery in chronic opiate users resulted in a similar postoperative opioid use as opioid-naïve patients. Nguyen et al [20] reported significantly improved postoperative disease-specific and generic measures of health outcomes in patients who halved their opioid consumption before TJA compared with those with no intervention for chronic preoperative opioid use.

An interesting finding of this was that 16% of patients, including non-opioid-naïve and revision patients, never filled their opioid prescription. Although our small sample size limited our ability to draw statistical inferences, the finding gives hope that we may eventually be able to develop opioid-free analgesic protocols for select patients. This would also effectively reduce the number of dispensed opioids that go unused and pose a potential for misuse. Sabatino et al [14] reported a mean of 31.9 unconsumed pills after THA and 28.9 unconsumed pills after TKA. They estimated that approximately 43,216 opioid pills prescribed at their institution went unused during a 6-month period.

Orthopaedic surgeons are major prescribers of opioid medications, and therefore, we must be cognizant of our role in combating the opioid epidemic. Preoperative counseling and engagement of patients and their families have been shown to increase patient satisfaction and decrease opioid consumption [17,21–23]. Limiting the size of opioid prescriptions is not only an effective strategy to protect our patients [19] but also increasingly mandated by state laws and regulatory authorities. Several states have enacted legislative measures limiting the amount of opioid that can be prescribed, and all 50 states have implemented prescription drug monitoring programs [21,24].

There are some limitations for this study. First, it is a retrospective review from a single surgeon operating at a single, tertiary-care academic center. Second, the patient sample is too small to infer statistical associations. Third, the 60-pill opioid prescription amount was chosen to accommodate the use of 1 pill every 3 hours as needed. A lower amount could have been sufficient for most patients as only one patient made a refill request and some pharmacies did not fill the full 60-pill prescription. A refill request is not a huge burden for the prescriber but may be more difficult for the patient to fill. The more important goal should be limiting opioid consumption and not the refill number. Fourth, it is unknown how many pills were actually consumed by patients in the cohort. Nonetheless, we reported on a consecutive series of all incomers (primary, complex, and revision cases) with no exclusions, and all patients underwent a standardized care pathway.

**Table 3**  
Reasons for emergency room visits and complications.

ER visits	N
Decompensated congestive heart failure	1
Pneumonia	1
Lower extremity swelling/ultrasound examination	2
Complications	
Failure of arthroscopy repair secondary to forced flexion	1
Instability after revision total hip arthroplasty	1
Wound dehiscence requiring medial gastrocnemius rotation flap	1
Mortality secondary to metastatic lung cancer	1

In conclusion, a single prescription of low-dose opioids appears to be sufficient for patients undergoing TJA when using preoperative counseling and opioid-sparing multimodal analgesia even in complex primary and revision procedures. As we combat the opioid epidemic, standardized guidelines are needed to guide best practices for pain management in patients on chronic opioid therapy. This information can help accelerate the decline of opioid use and hopefully pave the way for opioid-free TJA.

### Conflict of interest

M.A. Harrington is a paid consultant for Zimmer Inc and is a board member for AOA, AAHKS, AAOS, and J Robert Gladden Orthopaedic Society. M.J. Halawi is a board member for AAHKS Publications Committee.

### References

- [1] Centers for Disease Control and Prevention. National Ambulatory Medical Care Survey: 2016. [https://www.cdc.gov/nchs/data/ahcd/namcs\\_summary/2016\\_namcs\\_web\\_tables.pdf](https://www.cdc.gov/nchs/data/ahcd/namcs_summary/2016_namcs_web_tables.pdf). [Accessed 15 December 2019].
- [2] Food and Drug Administration. Joint Meeting of the Drug Safety and Risk Management Advisory Committee (DSaRM) and the Anesthetic and Analgesic Drug Products Advisory Committee (AADPAC): FDA Introductory Remarks. <https://www.fda.gov/media/97531/download>. [Accessed 15 December 2019].
- [3] Rose ME. Are prescription opioids driving the opioid crisis? Assumptions vs facts. *Pain Med* 2018;19(4):793.
- [4] Hannon CP, Keating TC, Lange JK, Ricciardi BF, Waddell BS, Della Valle CJ. Anesthesia and analgesia practices in total joint arthroplasty: a Survey of the American Association of Hip and Knee Surgeons Membership. *J Arthroplasty* 2019;34(12):2872.
- [5] Kim KY, Anoushiravani AA, Chen KK, Roof M, Long WJ, Schwarzkopf R. Preoperative chronic opioid users in total knee arthroplasty—which patients persistently abuse opiates following surgery? *J Arthroplasty* 2018;33(1):107.
- [6] Prentice HA, Inacio MCS, Singh A, Namba RS, Paxton EW. Preoperative risk factors for opioid utilization after total hip arthroplasty. *J Bone Joint Surg Am* 2019;101(18):1670.
- [7] Pivec R, Issa K, Naziri Q, Kapadia BH, Bonutti PM, Mont MA. Opioid use prior to total hip arthroplasty leads to worse clinical outcomes. *Int Orthop* 2014;38(6):1159.
- [8] Vakharia RM, Sabeh KG, Vakharia AM, Damodar DM, Law TY, Roche MW. Comparison of implant related complications amongst patients with opioid use disorder and non-users following total knee arthroplasty. *World J Orthop* 2019;10(3):137.
- [9] Zywiell MG, Stroh DA, Lee SY, Bonutti PM, Mont MA. Chronic opioid use prior to total knee arthroplasty. *J Bone Joint Surg Am* 2011;93(21):1988.
- [10] Kim SC, Jin Y, Lee YC, et al. Association of preoperative opioid use with mortality and short-term safety outcomes after total knee replacement. *JAMA Netw Open* 2019;2(7):e198061.
- [11] Buvanendran A, Kroin JS, Tuman KJ, et al. Effects of perioperative administration of a selective cyclooxygenase 2 inhibitor on pain management and recovery of function after knee replacement: a randomized controlled trial. *JAMA* 2003;290(18):2411.
- [12] Golladay GJ, Balch KR, Dalury DF, Satpathy J, Jiranek WA. Oral multimodal Analgesia for total joint arthroplasty. *J Arthroplasty* 2017;32(9S):S69.
- [13] Sawhney M, Mehdian H, Kashin B, et al. Pain after unilateral total knee arthroplasty: a prospective randomized controlled trial examining the analgesic effectiveness of a combined adductor Canal peripheral nerve block with periarticular infiltration versus adductor canal nerve block alone versus periarticular infiltration alone. *Anesth Analg* 2016;122(6):2040.
- [14] Sabatino MJ, Kunkel ST, Ramkumar DB, Keeney BJ, Jevsevar DS. Excess opioid medication and variation in prescribing patterns following common orthopaedic procedures. *J Bone Joint Surg Am* 2018;100(3):180.
- [15] Hernandez NM, Parry JA, Taunton MJ. Patients at risk: large opioid prescriptions after total knee arthroplasty. *J Arthroplasty* 2017;32(8):2395.
- [16] Premkumar A, Lovecchio FC, Stepan JG, et al. Characterization of opioid consumption and disposal patterns after total knee arthroplasty. *Bone Joint J* 2019;101-B(7\_Supple\_C):98.
- [17] Louw A, Puenteadura EJ, Reed J, Zimney K, Grimm D, Landers MR. A controlled clinical trial of preoperative pain neuroscience education for patients about to undergo total knee arthroplasty. *Clin Rehabil* 2019;33(11):1722.
- [18] Brock JL, Jain N, Phillips FM, Malik AT, Khan SN. Postoperative opioid cessation rates based on preoperative opioid use: analysis of common orthopaedic procedures. *Bone Joint J* 2019;101-B(12):1570.
- [19] Shah RP, Karas V, Berger RA. Rapid discharge and outpatient total joint arthroplasty introduce a burden of care to the surgeon. *J Arthroplasty* 2019;34(7):1307.
- [20] Nguyen LC, Sing DC, Bozic KJ. Preoperative reduction of opioid use before total joint arthroplasty. *J Arthroplasty* 2016;31(9 Suppl):282.
- [21] American Academy of Orthopaedic Surgeons. Opioid use, misuse, and abuse in orthopaedic practice. [https://www.aaos.org/uploadedFiles/PreProduction/About/Opinion\\_Statements/advistmt/1045%20Opioid%20Use,%20Misuse,%20and%20Abuse%20in%20Practice.pdf](https://www.aaos.org/uploadedFiles/PreProduction/About/Opinion_Statements/advistmt/1045%20Opioid%20Use,%20Misuse,%20and%20Abuse%20in%20Practice.pdf). [Accessed 15 December 2019].
- [22] Levinson W, Hudak P, Tricco AC. A systematic review of surgeon-patient communication: strengths and opportunities for improvement. *Patient Educ Couns* 2013;93(1):3.
- [23] Yajnik M, Hill JN, Hunter OO, et al. Patient education and engagement in postoperative pain management decreases opioid use following knee replacement surgery. *Patient Educ Couns* 2019;102(2):383.
- [24] Lovecchio F, Derman P, Stepan J, et al. Support for safer opioid prescribing practices: a Catalog of published use after orthopaedic surgery. *J Bone Joint Surg Am* 2017;99(22):1945.