



Review

# Hydrogen Peroxide: Its Role in Plant Biology and Crosstalk with Signalling Networks

Martin Černý<sup>1,2,\*,†</sup> , Hana Habánová<sup>1,3,4,†</sup>, Miroslav Berka<sup>1</sup>, Markéta Luklová<sup>1,3</sup> and Břetislav Brzobohatý<sup>1,3,5</sup>

<sup>1</sup> Department of Molecular Biology and Radiobiology, Faculty of AgriSciences Mendel University in Brno, 613 00 Brno, Czech Republic; habanova.ha@gmail.com (H.H.); miroslavberka94@gmail.com (M.B.); luklovam@gmail.com (M.L.); brzoboha@ibp.cz (B.B.)

<sup>2</sup> Phytophthora Research Centre, Faculty of AgriSciences, Mendel University in Brno, 613 00 Brno, Czech Republic

<sup>3</sup> CEITEC—Central European Institute of Technology, Faculty of AgriSciences Mendel University in Brno, 613 00 Brno, Czech Republic

<sup>4</sup> Brno Ph.D. Talent, South Moravian Centre for International Mobility, 602 00 Brno, Czech Republic

<sup>5</sup> Institute of Biophysics AS CR, 613 00 Brno, Czech Republic

\* Correspondence: martincerny83@gmail.com; Tel.: +420-545-133-374

† These authors contributed equally to this work.

Received: 24 July 2018; Accepted: 15 September 2018; Published: 18 September 2018



**Abstract:** Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) is steadily gaining more attention in the field of molecular biology research. It is a major REDOX (reduction–oxidation reaction) metabolite and at high concentrations induces oxidative damage to biomolecules, which can culminate in cell death. However, at concentrations in the low nanomolar range, H<sub>2</sub>O<sub>2</sub> acts as a signalling molecule and in many aspects, resembles phytohormones. Though its signalling network in plants is much less well characterized than are those of its counterparts in yeast or mammals, accumulating evidence indicates that the role of H<sub>2</sub>O<sub>2</sub>-mediated signalling in plant cells is possibly even more indispensable. In this review, we summarize hydrogen peroxide metabolism in plants, the sources and sinks of this compound and its transport via peroxiporins. We outline H<sub>2</sub>O<sub>2</sub> perception, its direct and indirect effects and known targets in the transcriptional machinery. We focus on the role of H<sub>2</sub>O<sub>2</sub> in plant growth and development and discuss the crosstalk between it and phytohormones. In addition to a literature review, we performed a meta-analysis of available transcriptomics data which provided further evidence for crosstalk between H<sub>2</sub>O<sub>2</sub> and light, nutrient signalling, temperature stress, drought stress and hormonal pathways.

**Keywords:** H<sub>2</sub>O<sub>2</sub>; plant hormone; signalling; growth and development; stress

## 1. Introduction

Hydrogen peroxide, a chemical compound discovered by Louis Jacques Thenard a hundred years ago, has properties that could justify classifying it as a phytohormone. In nature, it can be of inorganic origin, for example, via reactions in the atmosphere [1] but H<sub>2</sub>O<sub>2</sub> from this source has only an indirect effect on living organisms. Thenard was the first to discover not only that H<sub>2</sub>O<sub>2</sub> decomposes into water but also that it can cause skin blistering at a high concentration. However, oxidative stress is not the sole effect of this molecule. It is an evolutionarily conserved signalling molecule and in plants, it has gained attention also for its role in the regulation of growth and development. Indeed, the number of H<sub>2</sub>O<sub>2</sub>-related research articles published each year has doubled since 2008, with Web of Science listing over 3000 plant science publications on this topic in the last five years. In this review, we summarize different aspects of H<sub>2</sub>O<sub>2</sub>-mediated responses in plants, starting with the sources,


catabolism and transport of H<sub>2</sub>O<sub>2</sub>. We then describe mechanisms for its perception and discuss its role in plant signalling networks and its effects on plant growth and development.

## 2. Metabolism

Hydrogen peroxide H<sub>2</sub>O<sub>2</sub> is a non-radical reactive oxygen species (ROS) and it, like singlet oxygen <sup>1</sup>O<sub>2</sub> and free radicals such as superoxide anion O<sub>2</sub><sup>-</sup> and hydroxyl radical •OH, is one of the major members of the ROS family [2]. In contrast to other ROS, H<sub>2</sub>O<sub>2</sub> is relatively stable, with a half-life of ms and its level in a plant leaf oscillates around 1 μmol per gram of fresh weight under natural conditions [3]. There are numerous routes, both enzymatic and non-enzymatic, for H<sub>2</sub>O<sub>2</sub> production in plant cells. The key sources include photorespiration, electron transport chains and redox reactions in the apoplast [4,5]. The KEGG (Kyoto Encyclopedia of Genes and Genomes) database lists 150 classes of enzyme that produce or utilize hydrogen peroxide. Of these, only 29 enzymes encoded by 227 genes are annotated in *Arabidopsis* and the largest enzyme family formed by peroxidases has 75 entries (Figure 1, Supplementary tables). However, not all of these enzymes necessarily participate in peroxide metabolism in plants. For instance, a flavin-containing monooxygenase like YUC6 may produce hydrogen peroxide in the absence of its substrate but in vitro experiments indicate that in this case the uncoupled reaction represents less than 4% of the enzyme's activity [6]. In contrast, mammalian flavin-containing monooxygenases are clearly a source of hydrogen peroxide [7]. The key enzymes that are involved in *Arabidopsis* H<sub>2</sub>O<sub>2</sub> metabolism reside in the apoplast, peroxisome, chloroplast and mitochondria and they will be described in detail.

### 2.1. Electron Transport Chains and Superoxide Dismutase

Under favourable conditions, the majority of intracellular H<sub>2</sub>O<sub>2</sub> is produced from molecular oxygen by a stepwise reaction via a superoxide anion intermediate which undergoes enzymatic reduction to H<sub>2</sub>O<sub>2</sub>. Excessive energy and/or malfunctioning of chloroplast and mitochondrial energetic metabolism are key causes of superoxide anion generation in plant cells. In chloroplasts, superoxide anions are produced when the electron-transport chain of photosystem I is oversaturated by excessive irradiation and electrons are transmitted by the Mehler reaction to oxygen molecules [8]. The resulting superoxide anions are then converted to H<sub>2</sub>O<sub>2</sub>. This dismutation step is a pH-dependent non-enzymatic event (for details see for example, [9]) but cells also catalyse the process by means of superoxide dismutase (SOD) in order to rapidly remove the toxic superoxide radical. Besides photosystem I, H<sub>2</sub>O<sub>2</sub> may also originate at the manganese-containing, oxygen-evolving complex which is the donor site of photosystem II and by the reduction of singlet oxygen or superoxide anions by photosynthetic electron transport chain components such as plastoquinol [10]. In seeds and non-photosynthetic parts of plants, the main sources of superoxide anion are coupled with the processes of cell respiration in mitochondria. Electron leakage occurs especially in complexes I, II and III and it is estimated that 1–5% of the oxygen entering the plant respiratory chain is converted into H<sub>2</sub>O<sub>2</sub> [11–13]. The *Arabidopsis* genome encodes eight SOD isozymes which can be divided into three classes according to their metal cofactor (Fe<sup>2+</sup>, Mn<sup>2+</sup>, Cu<sup>2+</sup>). There are three chloroplastic Fe-SODs and two Mn-SODs localized in mitochondria. The Fe-SODs are considered to be the oldest in evolutionary terms but the two classes share structural similarities and can also be found in prokaryotes. In contrast, the Cu/Zn-SOD class, which has three isozymes in *Arabidopsis*, most likely emerged after oxygen saturated the atmosphere. It is specific to eukaryotes and can be present in different cell compartments [14,15].



**Figure 1.** Key enzymes of hydrogen peroxide metabolism in plants. The list shows enzymes that directly catalyse hydrogen peroxide production or degradation in *Arabidopsis*, including the numbers of different isozymes, a comparison of relative gene expression profiles in seedlings, roots and shoots and the figure indicates subcellular localization. Colour coding: anabolic processes (green), catabolic processes (red), based on UniProt [16], SUBA 3.0 [17] and average gene expression profiles in 45, 24 and 7 NGS experiments for seedlings, leaf and root respectively (ThaleMine [18]).

### 2.2. NADPH Oxidase

The second largest group of H<sub>2</sub>O<sub>2</sub>-producing enzymes consists of the respiratory burst oxidases (Figure 1), which are also known as respiratory burst oxidase homologs (RBOHs) based on their homology to mammalian phagocyte NADPH oxidase (nicotinamide adenine dinucleotide phosphate

oxidase). RBOHs, together with the type III cell wall peroxidases, are associated with the so-called “oxidative burst,” which is considered to be one of the main responses of plant cells to biotic or abiotic stress [19,20] but is also a crucial part of normal plant growth and development [21]. RBOHs are plasma membrane-localized proteins which oxidize cytosolic NADPH, transferring the released electron to  $O_2$  and producing superoxide which is then dismutated. In *Arabidopsis*, there are ten RBOH genes which are divided into three classes according to their tissue-specificity [22,23]. RBOHs are probably the best studied enzymatic ROS-generating system in plants and different regulatory mechanisms have been described. RBOHs undergo multiple post-translational modifications (PTMs), including S-nitrosylation and phosphorylation, that are required for enzyme activity and are regulated by calcium ions and phosphatidic acid [24,25].

### 2.3. Polyamine Oxidase

Hydrogen peroxide is an end product of oxidative degradation of amines and polyamine degradation is considered to be an especially important source of hydrogen peroxide in plants (e.g., [26]). Plant polyamines are catabolized by two distinct classes of amine oxidases, the flavin adenine dinucleotide (FAD)-dependent polyamine oxidases and the copper amine oxidases, of which there are, respectively, five and eight putative functional isozymes encoded by the *Arabidopsis* genome [27]. The copper amine oxidases oxidize primary amino groups, producing ammonia,  $H_2O_2$  and an aminoaldehyde, whereas the polyamine oxidases oxidize the secondary amino groups and the reaction products depend on the catalytic mechanism and substrate specificity of a given isozyme. All five *Arabidopsis* polyamine oxidases are reportedly intracellular and oxidize the carbon on the exo-side of the  $N^4$  atom of spermine and spermidine to produce 1,3-diaminopropane,  $H_2O_2$  and an aminoaldehyde [28]. Polyamines play an important role in plant tolerance of abiotic stress and at least part of this tolerance is associated with hydrogen peroxide production (see for example, review [29]). Furthermore, polyamines represent a direct link between  $H_2O_2$  and hormonal pathways, as it has been shown that cytokinin increases the polyamine content of plants [30].

### 2.4. Peroxisomal Production of $H_2O_2$

Peroxisomal enzymes represent a major site of  $H_2O_2$  production in a plant cell. In *Arabidopsis*, in addition to SOD and amine oxidases that are present in multiple compartments, peroxisomes contain acyl-CoA oxidases, glycolate oxidases, uricase, sulphite oxidase, aldehyde oxidase and sarcosine oxidase (Figure 1). Xanthine oxidase, which converts xanthine to urate and  $H_2O_2$ , can be also localized in peroxisomes [2] but a putative *Arabidopsis* homolog that preferentially accepts  $NAD^+$  as the electron acceptor [31] reportedly resides in the cytosol. A significant proportion of peroxisomal  $H_2O_2$  originates during the beta-oxidation of long-chain fatty acids via acyl-CoA oxidase [32], which is an especially important process in germinating seeds that contain glyoxysomes, specialized peroxisome-like organelles. However, in photosynthetic tissues, the role of peroxisomes in  $H_2O_2$  metabolism is predominantly via photorespiration reactions that may contribute up to 70% of the total production of  $H_2O_2$  in a plant cell [33,34]. In this reaction, glycolate produced in chloroplasts is converted to glyoxylate by glycolate oxidase in peroxisomes. The *Arabidopsis* genome contains five genes encoding glycolate oxidase and their combined relative expression in photosynthetic tissues is the highest of all  $H_2O_2$ -producing enzymes (Figure 1). However, the actual levels of  $H_2O_2$  in peroxisomes are kept in check by catalase and it is estimated that the peroxisomal  $H_2O_2$  concentration is under 10  $\mu M$  [35].

### 2.5. The $H_2O_2$ Scavenging System

Plant cells survive with  $H_2O_2$  levels that would kill animal cells and the estimated endogenous  $H_2O_2$  content of plant cells is usually much higher than that found in animals and bacteria [36].  $H_2O_2$  accumulation increases the probability of hydroxyl radical production via the Fenton reaction and this would cause significant oxidative damage to cellular structures if it were not for the presence of a highly

efficient antioxidant system. Higher plants contain several types of peroxidases, including catalases, ascorbate peroxidases (APX), thiol-specific peroxidases and classical secretory plant peroxidase. Furthermore, non-enzymatic compounds like tocopherols, ascorbic acid and flavonoids and glutathione play significant roles in  $H_2O_2$  scavenging [37,38]. The plastoquinone and ubiquinone pool also contribute to the ROS scavenging process as illustrated in recent reports [39,40]. In accordance, inhibition of enzymes that maintain the oxidized plastoquinone and ubiquinone pool, plastid terminal oxidases and mitochondrial alternative oxidases, respectively, stimulates  $H_2O_2$  production [41,42].

### 2.6. Catalases

Though catalase belongs to the peroxidase family, it is usually considered separately due to its unique ability to convert two molecules of  $H_2O_2$  into water and molecular oxygen without the need for any reductant. This heme-containing enzyme is first oxidized to a high-valence iron intermediate, which is then reduced by a further reaction with  $H_2O_2$  [43]. Under specific circumstances, the intermediate may also react with a different substrate and catalase may oxidize donors such as alcohols or phenols. Catalase has a high turnover rate but a low substrate affinity, with a  $K_m$  value in the millimolar range, a far greater concentration of  $H_2O_2$  than that expected to be present in the cell [35]. As an illustration, the activity of a single molecule of rice catalase ( $k_{cat}$  80,000;  $K_m$  100 mM) [44] would be equivalent to more than 2200% of tobacco APX ( $k_{cat}$  1800;  $K_m$  0.022 mM) [45] at 100 mM  $H_2O_2$  but to only 1% at concentrations below 1  $\mu$ M  $H_2O_2$ , which would render catalase redundant. Of course, the constants determined in vitro may be misleading; the active form of catalase is a tetramer and it has been shown that, for example, PTMs may significantly affect the kinetics of a multimeric enzyme (e.g., [46]). Nevertheless, even though catalase activity has also been reported in the cytosol and mitochondria, its predominant localization is in peroxisomes, compartments with a high  $H_2O_2$  concentration where its efficiency should be greatest (e.g., [47]). There are three functionally conserved classes of catalase with different spatial and developmental localizations in plants. For example, in tobacco catalase class I detoxifies  $H_2O_2$  produced in photorespiration reactions, class II is localized in the vascular system and class III is present predominantly in flowers and fruits [48].

### 2.7. Ascorbate and Thiol-Specific Peroxidases

APX and glutathione peroxidases belong to the most important group of intracellular peroxidases [49]. Several types of APX have been described in plants; they include soluble enzymes in the cytosol, chloroplast and mitochondria and membrane-bound peroxidases in peroxisomes, glyoxysomes and thylakoids [50]. APX is the first enzyme in the so-called ascorbate-glutathione cycle, which includes monodehydroascorbate reductase, dehydroascorbate reductase and glutathione reductase and reduces  $H_2O_2$  and regenerates ascorbate via NAD(P)H [49]. The *Arabidopsis* genome encodes seven different APX isozymes and as indicated above, APX may be more important than catalase for  $H_2O_2$  metabolism. Indeed, it has been shown that in the absence of cytosolic APX1, the entire chloroplastic  $H_2O_2$ -scavenging system in *Arabidopsis* collapses,  $H_2O_2$  levels increase and protein oxidation occurs [51]. The thiol-specific peroxidases peroxiredoxins and glutathione peroxidases detoxify a broad spectrum of peroxide substrates [8]. However, recent evidence from *S. cerevisiae* indicates that this could be a secondary role and that thiol peroxidases perceive and transfer oxidative signals to signalling proteins and regulate transcription [52]. In plants and bacteria, six groups of peroxiredoxins are recognized on the basis of differences in sequence, structure and positions of conserved cysteinyl residues [53].

### 2.8. Peroxidases (Class III)

Peroxidases are by far the most abundant family of enzymes in  $H_2O_2$  metabolism (Figure 1). These so-called class III peroxidases probably have a correspondingly diverse range of functions, of which only a few, in certain plant species, have been revealed (see for example [54,55] for details). From the point of view of this review, it is important to note that the class III peroxidases participate not only



in  $H_2O_2$  catabolism via oxidation of phenolic compounds but also in producing it via an oxidative cycle using apoplastic reductants. For instance, it has been shown that in *Arabidopsis* cell culture they contribute to ca. 50% of the  $H_2O_2$  produced during the oxidative burst in pathogen defence [56]. Class III peroxidases can be found in vacuoles but the majority are apoplastic or associated with cell walls in the apoplast as they play a key role in maintaining cell wall integrity by catalysing its cross-linking and loosening, lignification and suberization [57].

### 3. Transport

Normal levels of  $H_2O_2$  leaf extracts are reported to be in the  $\mu\text{mol}$  per gram of fresh weight range but they may significantly vary within the same plant [3]. For instance, localization of hydrogen peroxide in different regions of the leaf reveals a pattern of increasing accumulation from the base to the leaf tip [58]. There is no clear evidence for long distance transport of  $H_2O_2$  but it is the least reactive ROS and this allows it to travel at least among neighbouring cells or cellular compartments and to serve as an important signalling molecule [59]. Thus, if it is able to escape the  $H_2O_2$ -scavenging mechanisms described above and is not reduced to the highly reactive hydroxyl radical, it may freely diffuse from the site of its generation and reach its putative target. Questions of how it overcomes the competing  $H_2O_2$ -scavengers that prevent the targeted oxidation of redox-regulated proteins are still not fully answered [60] but it is now clear that transport mediated by simple diffusion would not explain, for example, rapid stress-induced transfer of  $H_2O_2$  generated in apoplast by NADPH oxidases into cytosol and that a  $H_2O_2$ -specific transporter or channel must therefore exist.

#### *Peroxioporins*

Henzer and Steudle found that treatment with  $HgCl_2$  (an aquaporin activity inhibitor) caused a rapid decrease in  $H_2O_2$  and water influx and they postulated the existence of an aquaporin subclass, peroxiporins [61]. The similarity of  $H_2O_2$  to the water molecule indicates that aquaporins could have such a function. Plant aquaporins are recognized as multifunctional proteins transporting not only water but also many other small uncharged molecules (e.g.,  $CO_2$  and nutrients) and they thus play a role in the regulation of plant growth and development and in responses to a wide range of stresses. Aquaporins belong to the ancient superfamily of major intrinsic proteins (MIPs) and are present throughout living organisms with the exception of some Archea and bacteria [62]. Plant aquaporins are divided into five subfamilies: plasma membrane intrinsic proteins (PIPs), tonoplast intrinsic proteins (TIPs), nodulin26-like intrinsic proteins (NIPs), small basic intrinsic proteins (SIPs) and uncategorized intrinsic proteins (XIPs). The latter two groups, which were discovered more recently, are not present in some plant species [63].  $H_2O_2$  has a higher polarity than water and thus not all aquaporins are peroxiporins. For instance, Hooijmaijers et al. employed heterologous expression of all 13 *Arabidopsis* PIPs in yeast and found that only five of them inhibited yeast growth in the presence of  $H_2O_2$  [64]. Since the first report of  $H_2O_2$  transport by an aquaporin appeared, this phenomenon has been studied in diverse plant species, including maize [65], rice and barley [66], *Arabidopsis* [64,67–69], tulip [70], tobacco, potato and tomato [71]. Kim and Steudle (2009) suggested the occurrence of feedback regulation in aquaporin-facilitated  $H_2O_2$  transport, based on the observed inhibition of aquaporin transport capacity after  $H_2O_2$  treatment [72]. Further studies showed that this inhibition may occur indirectly by the internalization of aquaporin into vesicles that is caused by the change in the phosphorylation status of aquaporins [73,74]. Hooijmaijers et al. (2012) also found that  $H_2O_2$  treatment can alter aquaporin expression, indicating a feedback loop between  $H_2O_2$  concentration and peroxiporin expression [64].

### 4. Signalling

It has been widely reported that  $H_2O_2$  effects are dose-specific and that at low concentrations it serves as a signalling molecule. Despite  $H_2O_2$  being rapidly removed by protective enzymes, the scavenging mechanisms are less effective at concentrations of around 10 nM, enabling  $H_2O_2$  to be a second

messenger [59,75]. In general, proteins are primary targets of all oxidative species and there are two modes of action by which  $H_2O_2$  is perceived: direct oxidation of amino acid residues or reaction with reactive intermediates (e.g., [76]). The latter represents an indirect effect mediated via peroxide decomposition products (hydroxyl radical and singlet oxygen) and is usually considered to be a non-specific oxidative stress response. However, it has been shown that the transcription factor PerR, a major regulator of the peroxide inducible stress response in bacteria, senses  $H_2O_2$  via this pathway, employing metal-catalysed histidine oxidation [77]. The complexity of ROS-mediated processes in plants somewhat limits our understanding of  $H_2O_2$  signalling circuits and the present state of this understanding lags far behind that for bacteria, yeasts or mammalian cells. For instance, the ratio of superoxide radical to hydrogen peroxide may regulate the respiratory chain in mitochondria [78] and it is believed that the ratio of singlet oxygen plus superoxide radical to hydrogen peroxide determines the activation of cell death programs [79]. Some mechanisms have been conserved during evolution, whereas others seem to be plant-specific. Here, we summarize the main circuits that have been found to operate in plants.

#### 4.1. Oxidation of Cysteine Residues

Targets of direct oxidation are predominantly cysteinyl residues and reactive thiol side chains can act as sensors or switches in both signal transduction and regulation of enzyme activity [76]. Depending on  $H_2O_2$  concentration a cysteinyl residue can react to undergo several reversible or irreversible modifications, starting with sulfenic acid, which is highly reactive and reacts with other proximal thiolates resulting in the formation of inter/intramolecular disulphide bonds or S-glutathionylation. The reduction of disulphide bonds and the removal of glutathione are regulated by members of the thioredoxin and glutaredoxin enzyme families. Sulfenic acid can be also further oxidized by  $H_2O_2$  to sulfinic or even sulfonic acid [76,80]. Some signalling models predict that a hypothetical receptor may undergo successive oxidation steps and that each step would correspond to a physiological response but it remains to be seen whether such a receptor exists. Experiments carried out in vitro have shown that the rate of reaction of hydrogen peroxide with cysteine is relatively low but this does not apply to  $H_2O_2$ -scavenging enzymes. The reaction of the cysteinyl residue in peroxiredoxin has an apparent second order rate constant seven orders of magnitude higher than that for cysteinyl in BSA [81] and Marinho et al. calculated that the  $H_2O_2$  concentration needed for a peroxiredoxin-mediated response time of 5 min is as low as 0.2 nM [82]. The thiol-specific peroxidases thus act as receptors and, upon oxidation, interact with and oxidize effector proteins, forming a redox relay. For example, *Arabidopsis* glutathione peroxidase functions as both a redox transducer and a scavenger in stomatal closure [83]. Key enzymes in photosynthesis and carbohydrate metabolism are oxidized in response to  $H_2O_2$ , including RuBisCO, phosphoribulokinase, glyceraldehyde-3-phosphate dehydrogenase, transketolase and sedoheptulose-1,7-bisphosphatase [84]. It is very likely that this is also a redox relay mediated by peroxiredoxins present in the chloroplast but evidence for this is lacking.

#### 4.2. Oxidation of Methionine Residues

Methionine, the second proteinaceous sulphur-containing amino acid, is usually not considered to be a regulatory target in  $H_2O_2$  signalling but its first oxidized form (methionine sulfoxide) is the product of a PTM that can be reversed via the action of a specific reductase [76]. The fact that this enzyme increases  $H_2O_2$  tolerance indicates that methionine residues have a role at least in the  $H_2O_2$ -induced stress response [85]. Jacques et al. studied protein methionine sulfoxide dynamics in catalase knock-out *Arabidopsis* and found that 51 proteins were significantly more oxidized compared to wild-type. They also demonstrated that the activity of glutathione S-transferase is reduced upon methionine oxidation [86].

#### 4.3. Other Protein PTMs

It should be noted that the direct effect of  $H_2O_2$  on protein PTMs is not limited to cysteine or methionine residues. In fact, the presence of oxidative PTMs has been shown to interfere with other

PTMs close to the oxidized site [87]. An alteration in the PTM pattern can play a crucial role in signalling. The well-known regulator TP53, which participates in mammalian H<sub>2</sub>O<sub>2</sub> signalling, has to integrate a complex network of PTMs [82]. Its *Arabidopsis* orthologue SOG1 (suppressor of gamma response 1) is hyperphosphorylated in response to ROS and it has been proposed that H<sub>2</sub>O<sub>2</sub> regulates its hyperphosphorylation, ultimately leading to cell cycle regulation [88]. Examples from mammalian systems also indicate that PTM by ubiquitination and targeted protein degradation is key to the H<sub>2</sub>O<sub>2</sub> response [82]. However, our knowledge about its role in plant H<sub>2</sub>O<sub>2</sub> circuits is limited. It has been found that UPL5 ubiquitin ligase mediates degradation of the transcription factor WRKY53 [89] but there are more than 1500 E3 enzymes in *Arabidopsis* and this, together with extensive crosstalk with phytohormonal networks (which all to some extent converge on the proteasome) [90], represents a substantial obstacle to the elucidation of H<sub>2</sub>O<sub>2</sub> signal transduction.

#### 4.4. Transcription Factors

##### 4.4.1. HsfA

Heat-shock transcription factors are transcriptional activators that, once trimerized, specifically bind *cis*-elements called heat shock elements, palindromic DNA sequences that are found in the promoters of heat stress-inducible genes of all eukaryotes, including that encoding APX, the major catabolic enzyme in *Arabidopsis* H<sub>2</sub>O<sub>2</sub> metabolism [91,92]. The trimerization mechanism requires intramolecular disulphide bonds and it can be directly induced by H<sub>2</sub>O<sub>2</sub> (reviewed in, for example, [93]). In *Arabidopsis*, HsfA2 was found to be involved in H<sub>2</sub>O<sub>2</sub> signalling and it was shown that both its transcript and the transcript levels of its target genes were induced by treating with exogenous H<sub>2</sub>O<sub>2</sub> [94].

##### 4.4.2. NAC Domain-Containing Protein

NAC (No apical meristem *Arabidopsis* transcription activation factor Cup-shaped cotyledon) domain-containing proteins constitute one of the largest transcription factor families in plants and they are involved in multiple developmental and physiological processes, including senescence and abiotic stress responses. Multiple genes of this family have been found to be upregulated in response to H<sub>2</sub>O<sub>2</sub> [95] treatment and it has been suggested that NAC042 (JUB1) functions as a modulator of cellular H<sub>2</sub>O<sub>2</sub> levels [96]. NAC059-dependent gene expression was triggered after H<sub>2</sub>O<sub>2</sub> treatment [97] indicating that NAC could be a primary target of H<sub>2</sub>O<sub>2</sub>. Furthermore, two transcription factors (NAC013 and NAC017) that apparently shuttle between the nucleus and endoplasmic reticulum membrane mediate redox-related retrograde signalling [98,99].

##### 4.4.3. Mediators of RNA Polymerase

In yeast and mammals, an RNA polymerase inhibitor localized in cytosol is activated by H<sub>2</sub>O<sub>2</sub> through the thioredoxin system and translocated into the nucleus [82]. Its putative orthologue in *Arabidopsis* is not known to be a H<sub>2</sub>O<sub>2</sub>-responsive protein but mediators of RNA polymerase II have been found to be upregulated in response to H<sub>2</sub>O<sub>2</sub>, including MED37C [94]. Shaikhali et al. showed that members of this family readily form oligomers *in vitro* via intramolecular disulphide bonds [100] and showed that root growth in the knock-out mutant *med32* was significantly less affected by H<sub>2</sub>O<sub>2</sub> than that in wild-type plants [101].

##### 4.4.4. WRKY and ZAT (Zinc finger of *Arabidopsis thaliana*) Transcription Factors

There are 74 WRKY amino acid signature sequence-containing transcription factors in *Arabidopsis* that contain four-stranded  $\beta$ -sheet WRKY DNA binding domain/s ca 60 amino acids in length and zinc-finger motifs [102]. Like Nascent polypeptide-Associated Complex NAC domain-containing proteins, these transcription factors participate in stress-related responses and some have been found to be upregulated in response to H<sub>2</sub>O<sub>2</sub>. WRKY30 and WRKY53 were found to be upregulated in response



to ozone and H<sub>2</sub>O<sub>2</sub> exposure, with WRKY53 being much more responsive to H<sub>2</sub>O<sub>2</sub> than WRKY30 and vice versa for ozone [103]. WRKY46 was upregulated by H<sub>2</sub>O<sub>2</sub> [104] and WRKY70 is a putative interactor of the H<sub>2</sub>O<sub>2</sub>-responsive zinc finger protein ZAT7 [105]. ZAT12, another H<sub>2</sub>O<sub>2</sub>-responsive transcription factor, was proposed to mediate iron uptake control via its interaction with the FIT protein and with H<sub>2</sub>O<sub>2</sub> as a signal in iron deficiency responses [106]. The present evidence indicates that WRKY transcription factors and ZAT zinc finger proteins participate in responses to H<sub>2</sub>O<sub>2</sub> but a more detailed analysis of WRKY/ZAT-mediated transcription is needed in order to test the hypothesis that they play a role as the primary target. The fact that ZAT12 and ZAT5 respond positively to both ascorbate and H<sub>2</sub>O<sub>2</sub> [107] indicates that this is probably not the case, at least for the ZATs.

#### 4.5. Calcium Ions


Calcium ions play a key role in a vast array of signalling pathways in plants (e.g., [108]). Ca<sup>2+</sup> is a second messenger like H<sub>2</sub>O<sub>2</sub> and multiple characterized cascades require their combined effect, for example, via the opening of H<sub>2</sub>O<sub>2</sub>-dependent Ca<sup>2+</sup> channels [109–111]. The Ca<sup>2+</sup>-binding protein calmodulin is an activator of catalase [112] and calmodulin-binding transcription activators have been found to be upregulated by H<sub>2</sub>O<sub>2</sub>. BT2, another calmodulin-binding protein which is upregulated by H<sub>2</sub>O<sub>2</sub>, is also part of an E3 ligase complex [113,114]. Moreover, Ca<sup>2+</sup>-dependent phosphorylation activates NADPH oxidases (e.g., [115]) and plays a role in the so-called ROS-Ca<sup>2+</sup> hubs described in Section 5.11.

### 5. H<sub>2</sub>O<sub>2</sub> in Growth and Development

The role of H<sub>2</sub>O<sub>2</sub> in the life of plants is illustrated in Figure 2 and outlined in the following text, which presents examples from different developmental stages.

#### 5.1. The Crosstalk between H<sub>2</sub>O<sub>2</sub> and Phytohormones

The first genome-wide analyses of plant H<sub>2</sub>O<sub>2</sub> signalling revealed a connection between ethylene and H<sub>2</sub>O<sub>2</sub>. Ethylene signalling is induced in response to H<sub>2</sub>O<sub>2</sub> accumulation [116] but the ethylene receptor ETR1 itself perceives H<sub>2</sub>O<sub>2</sub> directly in an ethylene-independent manner that does not require its kinase domain [117]. ROS is a key component of phytohormonal signalling networks and does not only mediate stress-related pathways. From the proteome-wide point of view, catalases, peroxiredoxins, disulphide isomerases and thioredoxins have been detected at high frequencies in phytohormone-responsive proteomics studies and APX, glutathione S-transferase and class III peroxidase were found at least once in all reported hormone-responsive proteomes; see Table 1 [90]. H<sub>2</sub>O<sub>2</sub> mediates hormonal homeostasis (e.g., auxin conjugation [118] and degradation [119]) but enzymes involved in hormone metabolism may produce H<sub>2</sub>O<sub>2</sub>. These include abscisic acid aldehyde oxidases, enzymes that catalyse the final step in abscisic acid biosynthesis producing H<sub>2</sub>O<sub>2</sub> in the process. Auxin aldehyde oxidases are also present in *Arabidopsis* but it is not clear to what extent these enzymes contribute to auxin metabolism [120]. Furthermore, monooxygenases may catalyse a H<sub>2</sub>O<sub>2</sub>-producing side reaction, as illustrated above for the auxin biosynthetic enzyme YUC6, which is encoded by a member of a multigene family in *Arabidopsis*. Similar enzymes are present in the pathways of cytokinin metabolism (hydroxylases cytochrome P450 735A1 and 735A2), gibberellin and brassinosteroid biosynthesis (ent-kaurene oxidase, ent-kaurenoic acid oxidase) and abscisic acid metabolism (hydroxylases cytochrome P450 707A1-707A4). Our comparison of expression profiles revealed that multiple hormonal metabolism genes share patterns of expression with those of H<sub>2</sub>O<sub>2</sub> metabolism; the former include *ABA4* (31 similar patterns), tryptophan aminotransferases *TAR3* (29) and *TAR4* (29), methyl esterase *MES1* (28), cytokinin biosynthetic genes *IPT2* (26), *IPT6* (22) and *LOG2* (24), ethylene metabolism genes *ACO2* (23) and *ACS4* (20) and Ent-copalyl diphosphate synthase *GA1* (27) (see Table 2 and Supplementary Materials for details).



**Figure 2.** Hydrogen peroxide-mediated processes in plant growth and development. This figure summarizes our present-day knowledge about the role of H<sub>2</sub>O<sub>2</sub> in the life of plants as described, with references, in Section 5. ABA—abscisic acid, AUX—auxin, BR—brassinosteroids, ET—ethylene, GA—gibberellins, SA—salicylic acid, JA—jasmonic acid, AOX—amine oxidases, PRX—peroxidases, RBOH—NADPH oxidases, PCD—programmed cell death. The water droplet shape indicates flooding and absence of water, for hypoxia and drought, respectively.

**Table 1.** Proteins of hydrogen peroxide metabolism in *Arabidopsis* identified in phytohormone-responsive proteomics analyses. Based on a previously published overview of hormone-responsive proteins [90].

AGI	Protein Name (UniProt)	Relative Protein Abundance						
		Auxin	Abscisic Acid	Brassinosteroid	Cytokinin	Salicylic Acid	Jasmonate/Oxylipins	Strigolactone
AT1G05260	Peroxidase 3		down [121]					
AT1G06290	Acyl-coenzyme A oxidase 3						up [122]	
AT1G07890	L-Ascorbate peroxidase 1		down [121]		down [30,123,124]		up [125,126]	
AT1G08830	Superoxide dismutase [Cu-Zn] 1					up [127]		
AT1G20620	Catalase-3		up [128]		down [129]	up [130]		up [130]
AT1G20630	Catalase-1		up [131]					
AT1G31710	Amine oxidase		down [121]					
AT1G44446	Chlorophyllide a oxygenase			down [132]				
AT1G65980	Peroxiredoxin-2B		down [121]				up [126]	
AT1G71695	Peroxidase 12		down [121]	down [132]				
AT1G77490	L-Ascorbate peroxidase T	up [133]						
AT2G18150	Peroxidase 15					up [127]		
AT2G22420	Peroxidase 17					up [127]		
AT2G26230	Uricase		down [121]					
AT2G28190	Superoxide dismutase [Cu-Zn] 2						up [134]	
AT2G30490	Trans-cinnamate 4-monooxygenase		up [131]					
AT2G43350	Probable glutathione peroxidase 3		down [121]					
AT3G06050	Peroxiredoxin-2F						up [134]	
AT3G10920	Superoxide dismutase [Mn] 1				down [135]	up [136]	down [126]	
AT3G11630	2-Cys peroxiredoxin BAS1				up [30,129]		up [125]	
AT3G14415	(S)-2-hydroxy-acid oxidase		down [125]			up [130]	up [125,130]	
AT3G14420	(S)-2-hydroxy-acid oxidase GLO1				up [30]	up [130]	up [126,130]	
AT3G26060	Peroxiredoxin Q, chloroplastic						up [134]	
AT3G32980	Peroxidase 32				down [30]	up [127]		
AT3G49120	Peroxidase 34		up [128,131]		down [30]	up [127]		
AT3G56350	Superoxide dismutase [Mn] 2					up [137]		
AT4G08390	L-Ascorbate peroxidase S		up [126]		down [30]			
AT4G08770	Peroxidase 37					up [127]		
AT4G08780	Peroxidase 38					up [127]		
AT4G15760	Monooxygenase 1							up [137]
AT4G16760	Acyl-coenzyme A oxidase 1	up [133]					up [122]	
AT4G25100	Superoxide dismutase [Fe] 1					up [127]	up [125]	
AT4G35000	L-Ascorbate peroxidase 3				down [30]			

Table 1. Cont.

AGI	Protein Name (UniProt)	Relative Protein Abundance						
		Auxin	Abscisic Acid	Brassinosteroid	Cytokinin	Salicylic Acid	Jasmonate/Oxylipins	Strigolactone
AT4G35090	Catalase-2		up [131]		up/down [30,123]	up [130]	up [130]	
AT4G36430	Peroxidase 49					up [127]		
AT5G06290	2-Cys peroxiredoxin BAS1-like						up [126]	
AT5G14220	Protoporphyrinogen oxidase 2			up [132]	up [132]			
AT5G17820	Peroxidase 57		up [128]					
AT5G18100	Superoxide dismutase [Cu-Zn] 3					up [127]		
AT5G23310	Superoxide dismutase [Fe] 3			down [138]			down [122]	
AT5G49970	PYRIDOXINE/PYRIDOXAMINE 5'-PHOSPHATE OXIDASE 1						up [122]	
AT5G51100	Superoxide dismutase [Fe] 2		up [139]					
AT5G64120	Peroxidase 71		down [131]				up [122]	down [137]
AT5G65110	Acyl-coenzyme A oxidase 2			down [132]				

### 5.2. Light Signalling

Light signal transduction is involved in H<sub>2</sub>O<sub>2</sub> metabolism and/or signalling. It has been demonstrated that blue-light perception by cryptochrome is directly coupled with H<sub>2</sub>O<sub>2</sub> generation [140–142]. It has also been proposed that phytochrome B modulates homeostasis of reactive oxygen species in roots via synthesis and transport of abscisic acid [143]. Our comparison of expression profiles revealed that genes participating in light signalling share patterns of expression with H<sub>2</sub>O<sub>2</sub> metabolism genes; the former include *MED25* which acts in the repression of phytochrome B-mediated light signalling (26 similarities), *COP1* (32), *phytochrome A* (29), *PIF1* (28), *phytochrome B* (26), *phytochrome C* (25) and *cryptochrome 1* (18) (see Table 2 and Supplementary Materials for details).

### 5.3. Dry Seed

The majority of plants from temperate climate zones produce so-called orthodox seeds which pass through a phase of intensive desiccation and in this state, they are able to survive for periods ranging from months to tens of years (or even hundreds of years in some cases) [144]. The quiescent state limits enzymatic activity to a minimal level but H<sub>2</sub>O<sub>2</sub> and other ROS are still produced and can be accumulated during seed ageing. H<sub>2</sub>O<sub>2</sub> accumulation in seeds may cause significant damage to storage molecules and loss of viability but the degree to which it accumulates and the sensitivity to oxidative damage is species specific. For example, H<sub>2</sub>O<sub>2</sub> does not accumulate in *Brassica napus* seeds [145].

### 5.4. Germination

Seed germination is defined as a three-phase process, starting with rapid water intake and ending with seed coat rupture, usually by radicle protuberance. In imbibed and germinating seeds, high levels of H<sub>2</sub>O<sub>2</sub> are produced mainly as a product of intensive metabolism in mitochondria, peroxisomes and glyoxysomes but also by NADPH oxidases and through lipid peroxidation [146]. Though seeds contain both enzymatic and nonenzymatic ROS scavenging machinery to prevent oxidative damage [147], H<sub>2</sub>O<sub>2</sub> is also needed to remove mechanical and hormonal barriers that inhibit germination. H<sub>2</sub>O<sub>2</sub> promotes endosperm weakening [148,149] and triggers an increase in gibberellin biosynthesis and a decrease in abscisic acid levels [150–154]. It also mediates selective oxidation of mRNA and proteins [155,156], for example irreversible carbonylation of storage proteins that enables their rapid mobilization via proteasomes [157]. Another key aspect of seed germination, especially in cereals, is the activation of  $\alpha$ -amylase and the promotion of programmed cell death (PCD) in the aleurone layer. Here, H<sub>2</sub>O<sub>2</sub> is produced by NADPH oxidase and it functions via interplay with DELLA proteins (proteins with the highly conserved amino acid sequence motif DELLA), key components of the gibberellin signalling pathway [158–160]. In many respects, the role of H<sub>2</sub>O<sub>2</sub> in germination is similar to that of a growth regulator and studies of exogenous H<sub>2</sub>O<sub>2</sub> application have demonstrated that its effect is dose dependent [161–163].

### 5.5. Root Development

Ample evidence showed that ROS regulates root development via NADPH oxidases [111,164,165]. The phytohormone that is key to the regulation of root growth is auxin, which is well known to mediate changes in H<sub>2</sub>O<sub>2</sub> levels and thus promote cell growth and lateral root formation [166–168]. However, a recent study indicated that in mediating the induction of lateral roots, H<sub>2</sub>O<sub>2</sub> acts downstream of melatonin, an auxin-like indoleamine compound [169]. Root tip growth is also known to be affected by H<sub>2</sub>O<sub>2</sub> [170,171]. Polar auxin transport seems to regulate H<sub>2</sub>O<sub>2</sub>-induced root gravitropism [163] and exogenous H<sub>2</sub>O<sub>2</sub> treatment can disrupt this sensing, probably due to a change in auxin receptor distribution [172]. Abscisic acid inhibits primary root growth by activating NADPH oxidases and thus reducing auxin sensitivity [173] and a RBOH was proposed to interact with abscisic acid in the



regulation of lateral root growth in *Arabidopsis* under drought stress [174]. H<sub>2</sub>O<sub>2</sub>-mediated root growth in response to stress was also found in cucumber [175], cotton [176] and rice [177].

### 5.6. Shoot Development

Shoot growth and development of shoot architecture are driven by phytohormones, especially auxin and cytokinin, levels of which are highly correlated with environmental conditions. Auxin participates in cell growth by inducing cell wall peroxidases (peroxidases class III) and NADPH oxidases to produce ROS and promote cell wall loosening and further cell elongation [178,179]. H<sub>2</sub>O<sub>2</sub> has been reported to mediate apical dominance [180], photosynthesis [181] and leaf epinasty [182].

### 5.7. Stomatal Movement

Stomata are formed as gaps between pairs of guard cells and changes in guard cell turgor mediate the opening and closure of the stomatal pore. Stomatal closure is an example of rapid leaf-to-leaf communication mediated by ROS (e.g., [183]). Under conditions of excessive irradiation, an autopropagating wave of ROS rapidly transfers a signal to leaves that are not directly exposed to light and initiates stomatal closure. This closure can be induced by multiple stimuli, including brassinosteroids [184,185], strigolactones [186], jasmonic acid and salicylic acid [187], CO<sub>2</sub> [188], ethylene [189,190], glucose [191] and interactions among them [192]. The best-described mechanism is that mediated by abscisic acid that recruits calcium ions, nitric oxide (NO), H<sub>2</sub>O<sub>2</sub> and regulatory phosphorylation [193,194]. Guard cells generate H<sub>2</sub>O<sub>2</sub> by means of amine oxidases [195], peroxidases and RBOHs [196,197]. The activity of RBOHs is regulated by Ca<sup>2+</sup> binding [198] and phosphorylation by protein kinase OST1 (OPEN STOMATA 1) [199], which in turn is regulated by abscisic acid and interacts with a peroxiporin [200,201]. The overall H<sub>2</sub>O<sub>2</sub>-mediated machinery is much more complex. For example, hydrogen sulphide promotes H<sub>2</sub>O<sub>2</sub> production by stimulating RBOH activity [202] but the presence of flavonols in guard cells and H<sub>2</sub>O<sub>2</sub> scavenging inhibits stomatal closure [203] and it has been shown that this flavanol accumulation is induced by 5-aminolevulinic acid [204,205].

### 5.8. Pollination

In generatively propagating plants, H<sub>2</sub>O<sub>2</sub> and other ROS play a key role in pollen navigation and gametophyte fusion. Angiosperms have developed different reproductive barriers to avoid self-fertilization, one of the most widespread being self-incompatibility [206]. H<sub>2</sub>O<sub>2</sub> level is elevated during the incompatible reaction, triggering PCD. In contrast, the compatible reaction decreases the level of H<sub>2</sub>O<sub>2</sub> in the stigma and the development of the pollen tube is promoted. The further growth of and the penetration of the ovule by, the pollen tube is guided by quite complicated signalling machinery, including the FERONIA protein which modulates NADPH oxidase activity [207,208]. ROS accumulation, especially that of the hydroxyl radical which is largely generated from H<sub>2</sub>O<sub>2</sub>, is then crucial for pollen tube rupture and the release of sperm cells [209].

### 5.9. Fruit Ripening

The involvement of H<sub>2</sub>O<sub>2</sub> in the ripening process is known but not fully understood. Huan et al. proposed that H<sub>2</sub>O<sub>2</sub> acts as a signalling molecule in the middle stage of peach fruit development but that it serves as an important toxic molecule, stimulating lipid peroxidation and oxidative stress, during the late stage of fruit ripening [210]. Kumar et al. analysed ripening in tomato and found changes in the redox state during different stages of ripening with a significant increase of H<sub>2</sub>O<sub>2</sub> at the so-called breaker point (defined by the initial change in fruit colour) [211]. The increase in H<sub>2</sub>O<sub>2</sub> is most likely regulated by ethylene, the key regulator of fruit ripening that enhances respiration rate and ROS production [212].

### 5.10. Senescence and Cell Death

Senescence ultimately leads to the death of plant organs or whole plants. It is a multistep process by which the plant recovers and recycles valuable nutrient components that would otherwise be lost [213]. The role of H<sub>2</sub>O<sub>2</sub> in plant senescence was investigated by Bieker et al., who showed time-dependent levels of H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub>-scavenging enzymes in senescent leaves [214]. In such tissues H<sub>2</sub>O<sub>2</sub> mediates PCD together with stress phytohormones like ethylene [215] or salicylic acid [216]. H<sub>2</sub>O<sub>2</sub> levels are transiently elevated at the initial point of leaf senescence and peak again during the terminal stage [217,218] and this accumulation is reportedly more pronounced inside interveinal tissue [219]. Furthermore, transgenic lines with lower H<sub>2</sub>O<sub>2</sub> levels display delayed senescence [96,214].

### 5.11. Stress

The key phytohormones orchestrating plant stress responses are abscisic acid, salicylic acid, jasmonates and ethylene and all of these phytohormones employ H<sub>2</sub>O<sub>2</sub> in their signalling cascades in an either upstream or downstream manner [220]. Putative markers of nutrient status, temperature stress and drought stress share patterns of expression with those of H<sub>2</sub>O<sub>2</sub> metabolism (Table 2) and H<sub>2</sub>O<sub>2</sub> has been implicated in cold acclimation [221], salt stress responses and salt stress tolerance [222–224] and hypoxia stress [225]. Important targets in these responses are RBOHs [177,226,227]. Recently, maintenance of acquired thermotolerance was found to be interlinked with generation of H<sub>2</sub>O<sub>2</sub> by RBOHs [228] and these NADH oxidases also participate in H<sub>2</sub>O<sub>2</sub> production in biotic interactions. Under pathogen attack, ROS accumulation is involved in PCD of infected and surrounding cells [229]. This hypersensitive response is orchestrated by the phytohormones ethylene, JA and SA (e.g., [214]) but high cytokinin levels also induce H<sub>2</sub>O<sub>2</sub> accumulation [230]. H<sub>2</sub>O<sub>2</sub> has been implicated in the susceptibility of *Brassica napus* to *Leptosphaeria maculans* [231], resistance to root-knot nematode in tomato [232], systemic virus resistance in *Nicotiana benthamiana* [233] and reduction of rot in postharvest citrus fruits [234]. In accordance, plants primed with H<sub>2</sub>O<sub>2</sub> or with a higher basal level of H<sub>2</sub>O<sub>2</sub> formation display enhanced resistance to stressors [42,235].

It is well established that a significant proportion of H<sub>2</sub>O<sub>2</sub>-mediated stress response originates from its decomposition products. This decomposition is enhanced by the presence of transient metal catalysts through the so-called Haber-Weiss reaction. It is widely postulated that this reaction accounts for the in vivo generation of the highly reactive hydroxyl radical, which is a prime cause of oxidative damage to biomolecules (e.g., [9,236]). The hydroxyl radical is one of the strongest oxidants known and reacts at nearly diffusion-limited rates near the site of its formation [237]. Besides its ability to damage anything in its close vicinity and generate further radicals, the hydroxyl radical seems to be a potent effector in calcium and potassium homeostasis, regulation of cell elongation and stress-induced cell death [111,238–241]. Furthermore, hydroxyl radical-mediated activation of calcium channels is also proposed to be a part of the so-called ROS-Ca<sup>2+</sup> hub, the mechanism that is utilized to perceive and amplify signal. This self-amplifying system employs Ca<sup>2+</sup>-dependent phosphorylation of NADPH oxidases and promotes hydroxyl radical production that, in turn, stimulates Ca<sup>2+</sup>-influx and NADPH oxidases' activity (see for example [242]). The ROS-Ca<sup>2+</sup> hub is believed to be central to hypersensitive response, phytohormonal signalling or abiotic stress responses [115,158,243,244]

Organelles like chloroplasts or mitochondria are key cellular sensors of environmental fluctuations and integral parts of plant stress responses. They communicate information by signalling to nuclei via stress-triggered retrograde signals, including ROS (reviewed in Reference [245]). Recent reports show not only that H<sub>2</sub>O<sub>2</sub> participates indirectly via ROS triggered signals but also that it can transfer from chloroplasts to nuclei and facilitate photosynthetic control over gene expression [246].

**Table 2.** Hydrogen peroxide metabolism genes have expression patterns similar to those of genes related to light signalling, nutrient status, temperature stress, drought stress and hormonal metabolism. Based on average gene expression profiles in stress-related experiments (ThaleMine [11]) and reference stress-related genes [247]. Numbers indicate the number of analysed genes (numbers in brackets) and the number of detected co-expressed genes (hydrogen peroxide metabolism/candidate signalling and metabolism genes). See Supplementary Materials for the full list of co-expressed genes.

	Nutrient Stress (142)	Temperature Stress (43)	Drought Stress (13)	Light Signalling (27)	Abscisic Acid Metabolism (16)	Auxin Metabolism (31)	Brassinosteroid Metabolism (13)	Cytokinin Metabolism (37)	Ethylene Metabolism (12)	Gibberellin Metabolism (23)	Jasmonate Metabolism (17)	Salicylic Acid Metabolism (9)	Strigolactone Metabolism (3)
Amine/polyamine oxidase (15)	11/40	4/6	1/1	7/16	6/5	5/6	5/5	10/15	5/4	4/7	3/5	2/1	0/0
Respiratory burst oxidase (10)	10/58	6/6	4/1	9/16	5/3	7/14	8/8	9/21	4/4	8/6	3/2	4/4	4/3
Superoxide dismutase (8)	7/46	5/5	0/0	4/13	5/2	5/9	4/6	5/15	2/3	5/10	1/3	2/2	0/0
L-Gulonolactone oxidase (7)	7/31	2/2	2/1	4/5	4/5	4/7	2/1	7/10	3/3	3/5	3/4	3/2	2/2
Acyl-coenzyme A oxidase (7)	5/19	4/6	4/2	4/14	4/3	6/9	4/2	5/11	0/0	5/7	6/6	1/1	2/2
Glycolate oxidase (5)	5/10	2/1	0/0	5/8	0/0	3/3	4/2	3/4	0/0	2/1	3/2	0/0	3/1
Aldehyde/acetaldehyde oxidase (5)	4/36	2/1	2/1	4/9	3/3	5/8	4/4	4/13	1/1	3/7	1/2	3/3	1/2
Long-chain-alcohol oxidase (4)	3/18	3/4	0/0	2/7	2/2	3/5	1/1	3/6	1/1	1/1	1/2	2/2	0/0
Sulfhydryl oxidase (3)	3/25	3/5	0/0	3/13	3/2	3/8	3/3	3/5	2/2	3/6	3/3	2/1	1/1
Protoporphyrinogen oxidase (2)	2/21	2/1	0/0	2/5	2/2	2/6	2/3	2/6	2/1	2/3	0/0	1/2	0/0
Pyridoxal 5'-phosphate synthase (2)	2/28	2/3	0/0	2/9	2/2	2/7	2/3	2/8	2/2	2/5	1/1	2/2	1/1
L-Aspartate oxidase (1)	1/5	1/1	1/1	1/1	0/0	0/0	0/0	0/0	0/0	1/1	0/0	0/0	1/1
Sarcosine oxidase (1)	1/1	0/0	0/0	1/1	0/0	1/1	0/0	0/0	0/0	0/0	0/0	0/0	0/0
Uricase (1)	1/6	1/1	1/1	1/3	1/1	1/1	1/1	1/1	0/0	0/0	1/1	0/0	1/1
Sulphite oxidase (1)	1/12	1/3	0/0	1/9	1/1	1/5	1/1	1/4	1/1	1/3	1/1	1/1	0/0
Peroxidase (73)	53/106	21/9	9/2	33/25	21/7	35/21	29/13	43/30	21/6	27/15	17/8	18/7	10/3
Peroxiredoxin (10)	8/43	4/7	1/1	5/14	5/3	3/7	3/5	6/15	4/4	7/12	2/4	4/3	0/0
L-Ascorbate peroxidase (7)	6/44	6/7	2/2	6/16	3/2	4/10	5/7	6/16	3/4	6/9	4/3	3/2	2/2
Glutathione peroxidase (6)	5/26	3/6	0/0	3/10	1/1	5/8	3/3	3/6	3/2	3/3	1/4	1/1	2/2
Catalase (3)	3/13	2/2	1/1	2/13	1/1	3/5	2/3	3/8	0/0	3/3	2/4	0/0	2/1

## 6. Conclusions

H<sub>2</sub>O<sub>2</sub> represents a key signalling molecule, connecting the signalling pathways of multiple phytohormones and acting as a second messenger in response to diverse conditions modulating plant growth and development. Its dose-dependent effect on growth clearly indicates that H<sub>2</sub>O<sub>2</sub> is a growth regulator but can we also refer to H<sub>2</sub>O<sub>2</sub> as a putative phytohormone? It is produced and degraded by the plant in response to stimuli and it is perceived by specialized proteins and elicits a response at low nanomolar concentrations. However, the limiting factor is its transport. Though it can be readily transported within a single cell and exported to extracellular space, it is not believed to serve as a long-distance signal due to its low stability and the presence of H<sub>2</sub>O<sub>2</sub> scavengers. Exogenous treatment with H<sub>2</sub>O<sub>2</sub> elicits a response and H<sub>2</sub>O<sub>2</sub> gradients are established in plant organs but it is believed that signal propagation is sequential and that H<sub>2</sub>O<sub>2</sub> reaches only neighbouring cells [248]. In conclusion, the recent literature offers multiple examples that reveal H<sub>2</sub>O<sub>2</sub> as a versatile mediator of molecular communication in plants and whether we classify it as a phytohormone or not, this does not change its importance in the life of plants. There are new perspectives emerging in the field of H<sub>2</sub>O<sub>2</sub> research with tools being developed for the detection of low micromolar and even picomolar H<sub>2</sub>O<sub>2</sub> concentrations [249,250] and it is likely that their eventual application in plant sciences will provide answers to some of our questions about H<sub>2</sub>O<sub>2</sub> transport and concentration dynamics. Similarly, we may expect that increasing sensitivity in proteomics approaches combined with imaging or laser microdissection techniques (e.g., [251]) will reveal more H<sub>2</sub>O<sub>2</sub> targets and their spatio-temporal distribution.

**Supplementary Materials:** Supplementary materials can be found at <http://www.mdpi.com/1422-0067/19/9/2812/s1>.

**Author Contributions:** M.Č., H.H. performed the analytical and systematic search of the literature, analysed and interpreted the data and wrote the manuscript. M.L. performed and interpreted meta analyses, M.Č., H.H. and M.B. prepared figures, B.B. reviewed the final text and provided critical comments.

**Funding:** This research was funded partially by grant TE02000177 (TACR), AF-IGA-IP-2018/014 (Internal Grant Agency of Faculty of AgriSciences, Mendel University in Brno) and the LQ1601 (CEITEC 2020) project with financial contribution made by the Ministry of Education, Youths and Sports of the Czech Republic from within special support paid from the National Programme for Sustainability II funds.

**Conflicts of Interest:** The authors declare that they have no conflicts of interest.

## References

1. Möller, D. Atmospheric hydrogen peroxide: Evidence for aqueous-phase formation from a historic perspective and a one-year measurement campaign. *Atmos. Environ.* **2009**, *43*, 5923–5936. [CrossRef]
2. Das, K.; Roychoudhury, A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Front. Environ. Sci.* **2014**, *2*, 53. [CrossRef]
3. Cheeseman, J.M. Hydrogen peroxide concentrations in leaves under natural conditions. *J. Exp. Bot.* **2006**, *57*, 2435–2444. [CrossRef] [PubMed]
4. Wrzaczek, M.; Brosché, M.; Kangasjärvi, J. ROS signalling loops—Production, perception, regulation. *Curr. Opin. Plant Biol.* **2013**, *16*, 575–582. [CrossRef] [PubMed]
5. Foyer, C.H.; Bloom, A.J.; Queval, G.; Noctor, G. Photorespiratory Metabolism: Genes, Mutants, Energetics and Redox Signalling. *Annu. Rev. Plant Biol.* **2009**, *60*, 455–484. [CrossRef] [PubMed]
6. Dai, X.; Mashiguchi, K.; Chen, Q.; Kasahara, H.; Kamiya, Y.; Ojha, S.; DuBois, J.; Ballou, D.; Zhao, Y. The biochemical mechanism of auxin biosynthesis by an arabidopsis YUCCA flavin-containing monooxygenase. *J. Biol. Chem.* **2013**, *288*, 1448–1457. [CrossRef] [PubMed]
7. Siddens, L.K.; Krueger, S.K.; Henderson, M.C.; Williams, D.E. Mammalian flavin-containing monooxygenase (FMO) as a source of hydrogen peroxide. *Biochem. Pharmacol.* **2014**, *89*, 141–147. [CrossRef] [PubMed]
8. Dietz, K.-J.; Turkan, I.; Krieger-Liszka, A. Redox- and Reactive Oxygen Species-Dependent Signaling into and out of the Photosynthesizing Chloroplast. *Plant Physiol.* **2016**, *171*, 1541–1550. [CrossRef] [PubMed]

9. Demidchik, V. Mechanisms of oxidative stress in plants: From classical chemistry to cell biology. *Environ. Exp. Bot.* **2015**, *109*, 212–228. [[CrossRef](#)]
10. Khorobrykh, S.A.; Karonen, M.; Tyystjärvi, E. Experimental evidence suggesting that H<sub>2</sub>O<sub>2</sub> is produced within the thylakoid membrane in a reaction between plastoquinol and singlet oxygen. *FEBS Lett.* **2015**, *589*, 779–786. [[CrossRef](#)] [[PubMed](#)]
11. Huang, S.; Van Aken, O.; Schwarzländer, M.; Belt, K.; Millar, A.H. The Roles of Mitochondrial Reactive Oxygen Species in Cellular Signaling and Stress Response in Plants. *Plant Physiol.* **2016**, *171*, 1551–1559. [[CrossRef](#)] [[PubMed](#)]
12. Mignolet-Spruyt, L.; Xu, E.; Idänheimo, N.; Hoerberichts, F.A.; Mühlenbock, P.; Brosché, M.; Van Breusegem, F.; Kangasjärvi, J. Spreading the news: Subcellular and organellar reactive oxygen species production and signalling. *J. Exp. Bot.* **2016**, *67*, 3831–3844. [[CrossRef](#)] [[PubMed](#)]
13. Møller, I.M. Plant Mitochondria and Oxidative Stress: Electron Transport, NADPH Turnover and Metabolism of Reactive Oxygen Species. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **2001**, *52*, 561–591. [[CrossRef](#)] [[PubMed](#)]
14. Gill, S.S.; Anjum, N.A.; Gill, R.; Yadav, S.; Hasanuzzaman, M.; Fujita, M.; Mishra, P.; Sabat, S.C.; Tuteja, N. Superoxide dismutase—Mentor of abiotic stress tolerance in crop plants. *Environ. Sci. Pollut. Res.* **2015**, *22*, 10375–10394. [[CrossRef](#)] [[PubMed](#)]
15. Alscher, R.G.; Erturk, N.; Heath, L.S. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J. Exp. Bot.* **2002**, *53*, 1331–1341. [[CrossRef](#)] [[PubMed](#)]
16. The UniProt Consortium. UniProt: The universal protein knowledgebase. *Nucleic Acids Res.* **2017**, *45*, D158–D169. [[CrossRef](#)]
17. Tanz, S.K.; Castleden, I.; Hooper, C.M.; Vacher, M.; Small, I.; Millar, H.A. SUBA3: A database for integrating experimentation and prediction to define the SUBcellular location of proteins in Arabidopsis. *Nucleic Acids Res.* **2012**, *41*, D1185–D1191. [[CrossRef](#)] [[PubMed](#)]
18. Krishnakumar, V.; Hanlon, M.R.; Contrino, S.; Ferlanti, E.S.; Karamycheva, S.; Kim, M.; Rosen, B.D.; Cheng, C.-Y.; Moreira, W.; Mock, S.A.; et al. Araport: The Arabidopsis Information Portal. *Nucleic Acids Res.* **2015**, *43*, D1003–D1009. [[CrossRef](#)] [[PubMed](#)]
19. Marino, D.; Dunand, C.; Puppo, A.; Pauly, N. A burst of plant NADPH oxidases. *Trends Plant Sci.* **2012**, *17*, 9–15. [[CrossRef](#)] [[PubMed](#)]
20. Suzuki, N.; Miller, G.; Morales, J.; Shulaev, V.; Torres, M.A.; Mittler, R. Respiratory burst oxidases: The engines of ROS signaling. *Curr. Opin. Plant Biol.* **2011**, *14*, 691–699. [[CrossRef](#)] [[PubMed](#)]
21. Swanson, S.; Gilroy, S. ROS in plant development. *Physiol. Plant.* **2010**, *138*, 384–392. [[CrossRef](#)] [[PubMed](#)]
22. Tripathy, B.C.; Oelmüller, R. Reactive oxygen species generation and signaling in plants. *Plant Signal. Behav.* **2012**, *7*, 1621–1633. [[CrossRef](#)] [[PubMed](#)]
23. Sagi, M.; Fluhr, R. Production of reactive oxygen species by plant NADPH oxidases. *Plant Physiol.* **2006**, *141*, 336–340. [[CrossRef](#)] [[PubMed](#)]
24. Yun, B.-W.; Feechan, A.; Yin, M.; Saidi, N.B.B.; Le Bihan, T.; Yu, M.; Moore, J.W.; Kang, J.-G.; Kwon, E.; Spoel, S.H.; et al. S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* **2011**, *478*, 264–268. [[CrossRef](#)] [[PubMed](#)]
25. Qu, Y.; Yan, M.; Zhang, Q. Functional regulation of plant NADPH oxidase and its role in signaling. *Plant Signal. Behav.* **2017**, *12*, e1356970. [[CrossRef](#)] [[PubMed](#)]
26. Yoda, H.; Yamaguchi, Y.; Sano, H. Induction of hypersensitive cell death by hydrogen peroxide produced through polyamine degradation in tobacco plants. *Plant Physiol.* **2003**, *132*, 1973–1981. [[CrossRef](#)] [[PubMed](#)]
27. Tavladoraki, P.; Cona, A.; Angelini, R. Copper-Containing Amine Oxidases and FAD-Dependent Polyamine Oxidases Are Key Players in Plant Tissue Differentiation and Organ Development. *Front. Plant Sci.* **2016**, *7*, 824. [[CrossRef](#)] [[PubMed](#)]
28. Fincato, P.; Moschou, P.N.; Spedaletti, V.; Tavazza, R.; Angelini, R.; Federico, R.; Roubelakis-Angelakis, K.A.; Tavladoraki, P. Functional diversity inside the Arabidopsis polyamine oxidase gene family. *J. Exp. Bot.* **2011**, *62*, 1155–1168. [[CrossRef](#)] [[PubMed](#)]
29. Gupta, K.; Sengupta, A.; Chakraborty, M.; Gupta, B. Hydrogen Peroxide and Polyamines Act as Double Edged Swords in Plant Abiotic Stress Responses. *Front. Plant Sci.* **2016**, *7*, 1343. [[CrossRef](#)] [[PubMed](#)]
30. Černý, M.; Kuklová, A.; Hoehenwarter, W.; Fagner, L.; Novák, O.; Rotková, G.; Jedelský, P.L.; Žáková, K.; Šmehilová, M.; Strnad, M.; et al. Proteome and metabolome profiling of cytokinin action in Arabidopsis



- identifying both distinct and similar responses to cytokinin down- and up-regulation. *J. Exp. Bot.* **2013**, *64*, 4193–4206. [[CrossRef](#)] [[PubMed](#)]
31. Hesberg, C.; Hänsch, R.; Mendel, R.R.; Bittner, F. Tandem Orientation of Duplicated Xanthine Dehydrogenase Genes from *Arabidopsis thaliana*: Differential gene expression and enzyme activities. *J. Biol. Chem.* **2004**, *279*, 13547–13554. [[CrossRef](#)] [[PubMed](#)]
  32. Hu, J.; Baker, A.; Bartel, B.; Linka, N.; Mullen, R.T.; Reumann, S.; Zolman, B.K. Plant Peroxisomes: Biogenesis and Function. *Plant Cell* **2012**, *24*, 2279–2303. [[CrossRef](#)] [[PubMed](#)]
  33. Bauwe, H.; Hagemann, M.; Fernie, A.R. Photorespiration: Players, partners and origin. *Trends Plant Sci.* **2010**, *15*, 330–336. [[CrossRef](#)] [[PubMed](#)]
  34. Maurino, V.G.; Peterhansel, C. Photorespiration: Current status and approaches for metabolic engineering. *Curr. Opin. Plant Biol.* **2010**, *13*, 248–255. [[CrossRef](#)] [[PubMed](#)]
  35. Foyer, C.H.; Noctor, G. Stress-triggered redox signalling: What's in pROspect? *Plant Cell Environ.* **2016**, *39*, 951–964. [[CrossRef](#)] [[PubMed](#)]
  36. Costa, A.; Drago, I.; Behera, S.; Zottini, M.; Pizzo, P.; Schroeder, J.I.; Pozzan, T.; Schiavo, F.L. H<sub>2</sub>O<sub>2</sub> in plant peroxisomes: An in vivo analysis uncovers a Ca<sup>2+</sup>-dependent scavenging system. *Plant J.* **2010**, *62*, 760–772. [[CrossRef](#)] [[PubMed](#)]
  37. del Río, L.A. ROS and RNS in plant physiology: An overview. *J. Exp. Bot.* **2015**, *66*, 2827–2837. [[CrossRef](#)] [[PubMed](#)]
  38. Petrov, V.D.; Van Breusegem, F. Hydrogen peroxide—A central hub for information flow in plant cells. *AoB Plants* **2012**, *2012*, pls014. [[CrossRef](#)] [[PubMed](#)]
  39. Ksas, B.; Légeret, B.; Ferretti, U.; Chevalier, A.; Pospíšil, P.; Alric, J.; Havaux, M. The plastoquinone pool outside the thylakoid membrane serves in plant photoprotection as a reservoir of singlet oxygen scavengers. *Plant Cell Environ.* **2018**. [[CrossRef](#)] [[PubMed](#)]
  40. Khorobrykh, S.; Tyystjärvi, E. Plastoquinol generates and scavenges reactive oxygen species in organic solvent: Potential relevance for thylakoids. *Biochim. Biophys. Acta-Bioenerg.* **2018**, *1859*, 1119–1131. [[CrossRef](#)] [[PubMed](#)]
  41. Popov, V.; Simonian, R.; Skulachev, V.; Starkov, A. Inhibition of the alternative oxidase stimulates H<sub>2</sub>O<sub>2</sub> production in plant mitochondria. *FEBS Lett.* **1997**, *415*, 87–90. [[CrossRef](#)]
  42. Wiczarz, M.; Gubernator, B.; Kruk, J.; Niewiadomska, E. Enhanced chloroplastic generation of H<sub>2</sub>O<sub>2</sub> in stress-resistant *Thellungiella salsuginea* in comparison to *Arabidopsis thaliana*. *Physiol. Plant.* **2015**, *153*, 467–476. [[CrossRef](#)] [[PubMed](#)]
  43. Alfonso-Prieto, M.; Biarnés, X.; Vidossich, P.; Rovira, C. The molecular mechanism of the catalase reaction. *J. Am. Chem. Soc.* **2009**, *131*, 11751–11761. [[CrossRef](#)] [[PubMed](#)]
  44. Ray, M.; Mishra, P.; Das, P.; Sabat, S.C. Expression and purification of soluble bio-active rice plant catalase-A from recombinant *Escherichia coli*. *J. Biotechnol.* **2012**, *157*, 12–19. [[CrossRef](#)] [[PubMed](#)]
  45. Kitajima, S.; Kitamura, M.; Kojima, N. Triple mutation of Cys26, Trp35 and Cys126 in stromal ascorbate peroxidase confers H<sub>2</sub>O<sub>2</sub> tolerance comparable to that of the cytosolic isoform. *Biochem. Biophys. Res. Commun.* **2008**, *372*, 918–923. [[CrossRef](#)] [[PubMed](#)]
  46. Černý, M.; Doubnerová, V.; Müller, K.; Ryšlavá, H. Characterization of phosphoenolpyruvate carboxylase from mature maize seeds: Properties of phosphorylated and dephosphorylated forms. *Biochimie* **2010**, *92*, 1362–1370. [[CrossRef](#)] [[PubMed](#)]
  47. Mhamdi, A.; Queval, G.; Chaouch, S.; Vanderauwera, S.; Van Breusegem, F.; Noctor, G. Catalase function in plants: A focus on *Arabidopsis* mutants as stress-mimic models. *J. Exp. Bot.* **2010**, *61*, 4197–4220. [[CrossRef](#)] [[PubMed](#)]
  48. Mhamdi, A.; Noctor, G.; Baker, A. Plant catalases: Peroxisomal redox guardians. *Arch. Biochem. Biophys.* **2012**, *525*, 181–194. [[CrossRef](#)] [[PubMed](#)]
  49. Foyer, C.H.; Noctor, G. Ascorbate and glutathione: The heart of the redox hub. *Plant Physiol.* **2011**, *155*, 2–18. [[CrossRef](#)] [[PubMed](#)]
  50. Caverzan, A.; Passaia, G.; Rosa, S.B.; Ribeiro, C.W.; Lazzarotto, F.; Margis-Pinheiro, M. Plant responses to stresses: Role of ascorbate peroxidase in the antioxidant protection. *Genet. Mol. Biol.* **2012**, *35*, 1011–1019. [[CrossRef](#)] [[PubMed](#)]

51. Davletova, S.; Rizhsky, L.; Liang, H.; Shengqiang, Z.; Oliver, D.J.; Coutu, J.; Shulaev, V.; Schlauch, K.; Mittler, R. Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of Arabidopsis. *Plant Cell* **2005**, *17*, 268–281. [[CrossRef](#)] [[PubMed](#)]
52. Fomenko, D.E.; Koc, A.; Agisheva, N.; Jacobsen, M.; Kaya, A.; Malinouski, M.; Rutherford, J.C.; Siu, K.-L.; Jin, D.-Y.; Winge, D.R.; et al. Thiol peroxidases mediate specific genome-wide regulation of gene expression in response to hydrogen peroxide. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 2729–2734. [[CrossRef](#)] [[PubMed](#)]
53. Liebthal, M.; Maynard, D.; Dietz, K.-J. Peroxiredoxins and Redox Signaling in Plants. *Antioxid. Redox Signal.* **2018**, *28*, 609–624. [[CrossRef](#)] [[PubMed](#)]
54. Shigeto, J.; Tsutsumi, Y. Diverse functions and reactions of class III peroxidases. *New Phytol.* **2016**, *209*, 1395–1402. [[CrossRef](#)] [[PubMed](#)]
55. Podgórska, A.; Burian, M.; Szal, B. Extra-Cellular But Extra-Ordinarily Important for Cells: Apoplastic Reactive Oxygen Species Metabolism. *Front. Plant Sci.* **2017**, *8*, 1353. [[CrossRef](#)] [[PubMed](#)]
56. O'Brien, J.A.; Daudi, A.; Finch, P.; Butt, V.S.; Whitelegge, J.P.; Souda, P.; Ausubel, F.M.; Bolwell, G.P. A peroxidase-dependent apoplastic oxidative burst in cultured Arabidopsis cells functions in MAMP-elicited defense. *Plant Physiol.* **2012**, *158*, 2013–2027. [[CrossRef](#)] [[PubMed](#)]
57. Jovanović, S.V.; Kukavica, B.; Vidović, M.; Morina, F.; Menckhoff, L. Class III Peroxidases: Functions, Localization and Redox Regulation of Isoenzymes. In *Antioxidants and Antioxidant Enzymes in Higher Plants*; Springer International Publishing: Cham, Switzerland, 2018; pp. 269–300.
58. Tewari, R.K.; Singh, P.K.; Watanabe, M. The spatial patterns of oxidative stress indicators co-locate with early signs of natural senescence in maize leaves. *Acta Physiol. Plant.* **2013**, *35*, 949–957. [[CrossRef](#)]
59. Winterbourn, C.C. Biological Production, Detection and Fate of Hydrogen Peroxide. *Antioxid. Redox Signal.* **2017**, *29*, 541–551. [[CrossRef](#)] [[PubMed](#)]
60. Stöcker, S.; Van Laer, K.; Mijuskovic, A.; Dick, T.P. The Conundrum of Hydrogen Peroxide Signaling and the Emerging Role of Peroxiredoxins as Redox Relay Hubs. *Antioxid. Redox Signal.* **2018**, *28*, 558–573. [[CrossRef](#)] [[PubMed](#)]
61. Henzler, T.; Steudle, E. Transport and metabolic degradation of hydrogen peroxide in Chara corallina: Model calculations and measurements with the pressure probe suggest transport of H<sub>2</sub>O<sub>2</sub> across water channels. *J. Exp. Bot.* **2000**, *51*, 2053–2066. [[CrossRef](#)] [[PubMed](#)]
62. Abascal, F.; Irisarri, I.; Zardoya, R. Diversity and evolution of membrane intrinsic proteins. *Biochim. Biophys. Acta-Gen. Subj.* **2014**, *1840*, 1468–1481. [[CrossRef](#)] [[PubMed](#)]
63. Maurel, C.; Boursiac, Y.; Luu, D.-T.; Santoni, V.; Shahzad, Z.; Verdoucq, L. Aquaporins in Plants. *Physiol. Rev.* **2015**, *95*, 1321–1358. [[CrossRef](#)] [[PubMed](#)]
64. Hooijmaijers, C.; Rhee, J.Y.; Kwak, K.J.; Chung, G.C.; Horie, T.; Katsuhara, M.; Kang, H. Hydrogen peroxide permeability of plasma membrane aquaporins of *Arabidopsis thaliana*. *J. Plant Res.* **2012**, *125*, 147–153. [[CrossRef](#)] [[PubMed](#)]
65. Bienert, G.P.; Heinen, R.B.; Berny, M.C.; Chaumont, F. Maize plasma membrane aquaporin ZmPIP2;5 but not ZmPIP1;2, facilitates transmembrane diffusion of hydrogen peroxide. *Biochim. Biophys. Acta-Biomembr.* **2014**, *1838*, 216–222. [[CrossRef](#)] [[PubMed](#)]
66. Katsuhara, M.; Sasano, S.; Horie, T.; Matsumoto, T.; Rhee, J.; Shibasaki, M. Functional and molecular characteristics of rice and barley NIP aquaporins transporting water, hydrogen peroxide and arsenite. *Plant Biotechnol.* **2014**, *31*, 213–219. [[CrossRef](#)]
67. Bienert, G.P.; Møller, A.L.B.; Kristiansen, K.A.; Schulz, A.; Møller, I.M.; Schjoerring, J.K.; Jahn, T.P. Specific Aquaporins Facilitate the Diffusion of Hydrogen Peroxide across Membranes. *J. Biol. Chem.* **2007**, *282*, 1183–1192. [[CrossRef](#)] [[PubMed](#)]
68. Dynowski, M.; Schaaf, G.; Loque, D.; Moran, O.; Ludewig, U. Plant plasma membrane water channels conduct the signalling molecule H<sub>2</sub>O<sub>2</sub>. *Biochem. J.* **2008**, *414*, 53–61. [[CrossRef](#)] [[PubMed](#)]
69. Tian, S.; Wang, X.; Li, P.; Wang, H.; Ji, H.; Xie, J.; Qiu, Q.; Shen, D.; Dong, H. Plant Aquaporin AtPIP1;4 Links Apoplastic H<sub>2</sub>O<sub>2</sub> Induction to Disease Immunity Pathways. *Plant Physiol.* **2016**, *171*, 1635–1650. [[CrossRef](#)] [[PubMed](#)]
70. Azad, A.K.; Yoshikawa, N.; Ishikawa, T.; Sawa, Y.; Shibata, H. Substitution of a single amino acid residue in the aromatic/arginine selectivity filter alters the transport profiles of tonoplast aquaporin homologs. *Biochim. Biophys. Acta* **2012**, *1818*, 1–11. [[CrossRef](#)] [[PubMed](#)]

71. Bienert, G.P.; Bienert, M.D.; Jahn, T.P.; Boutry, M.; Chaumont, F. Solanaceae XIPs are plasma membrane aquaporins that facilitate the transport of many uncharged substrates. *Plant J.* **2011**, *66*, 306–317. [[CrossRef](#)] [[PubMed](#)]
72. Kim, Y.X.; Steudle, E. Gating of aquaporins by light and reactive oxygen species in leaf parenchyma cells of the midrib of *Zea mays*. *J. Exp. Bot.* **2009**, *60*, 547–556. [[CrossRef](#)] [[PubMed](#)]
73. Bienert, G.P.; Chaumont, F. Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochim. Biophys. Acta* **2014**, *1840*, 1596–1604. [[CrossRef](#)] [[PubMed](#)]
74. Verdoucq, L.; Rodrigues, O.; Martinière, A.; Luu, D.T.; Maurel, C. Plant aquaporins on the move: Reversible phosphorylation, lateral motion and cycling. *Curr. Opin. Plant Biol.* **2014**, *22*, 101–107. [[CrossRef](#)] [[PubMed](#)]
75. Sies, H. Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. *Redox Biol.* **2017**, *11*, 613–619. [[CrossRef](#)] [[PubMed](#)]
76. Černý, M.; Skalák, J.; Cerna, H.; Brzobohatý, B. Advances in purification and separation of posttranslationally modified proteins. *J. Proteomics* **2013**, *92*, 2–27. [[CrossRef](#)] [[PubMed](#)]
77. Lee, J.-W.; Helmann, J.D. The PerR transcription factor senses H<sub>2</sub>O<sub>2</sub> by metal-catalysed histidine oxidation. *Nature* **2006**, *440*, 363–367. [[CrossRef](#)] [[PubMed](#)]
78. Andronis, E.A.; Moschou, P.N.; Toumi, I.; Roubelakis-Angelakis, K.A. Peroxisomal polyamine oxidase and NADPH-oxidase cross-talk for ROS homeostasis which affects respiration rate in *Arabidopsis thaliana*. *Front. Plant Sci.* **2014**, *5*, 132. [[CrossRef](#)] [[PubMed](#)]
79. Sabater, B.; Martín, M. Hypothesis: Increase of the ratio singlet oxygen plus superoxide radical to hydrogen peroxide changes stress defense response to programmed leaf death. *Front. Plant Sci.* **2013**, *4*, 479. [[CrossRef](#)] [[PubMed](#)]
80. Couturier, J.; Chibani, K.; Jacquot, J.-P.; Rouhier, N. Cysteine-based redox regulation and signaling in plants. *Front. Plant Sci.* **2013**, *4*, 105. [[CrossRef](#)] [[PubMed](#)]
81. Davies, M.J. Protein oxidation and peroxidation. *Biochem. J.* **2016**, *473*, 805–825. [[CrossRef](#)] [[PubMed](#)]
82. Marinho, H.S.; Real, C.; Cyrne, L.; Soares, H.; Antunes, F. Hydrogen peroxide sensing, signaling and regulation of transcription factors. *Redox Biol.* **2014**, *2*, 535–562. [[CrossRef](#)] [[PubMed](#)]
83. Miao, Y.; Lv, D.; Wang, P.; Wang, X.-C.; Chen, J.; Miao, C.; Song, C.-P. An Arabidopsis Glutathione Peroxidase Functions as Both a Redox Transducer and a Scavenger in Abscisic Acid and Drought Stress Responses. *Plant Cell Online* **2006**, *18*, 2749–2766. [[CrossRef](#)] [[PubMed](#)]
84. Muthuramalingam, M.; Matros, A.; Scheibe, R.; Mock, H.-P.; Dietz, K.-J. The hydrogen peroxide-sensitive proteome of the chloroplast in vitro and in vivo. *Front. Plant Sci.* **2013**, *4*, 54. [[CrossRef](#)] [[PubMed](#)]
85. Li, C.-W.; Lee, S.-H.; Chieh, P.-S.; Lin, C.-S.; Wang, Y.-C.; Chan, M.-T. Arabidopsis Root-Abundant Cytosolic Methionine Sulfoxide Reductase B Genes MsrB7 and MsrB8 are Involved in Tolerance to Oxidative Stress. *Plant Cell Physiol.* **2012**, *53*, 1707–1719. [[CrossRef](#)] [[PubMed](#)]
86. Jacques, S.; Ghesquière, B.; De Bock, P.-J.; Demol, H.; Wahni, K.; Willems, P.; Messens, J.; Van Breusegem, F.; Gevaert, K. Protein Methionine Sulfoxide Dynamics in *Arabidopsis thaliana* under Oxidative Stress. *Mol. Cell. Proteomics* **2015**, *14*, 1217–1229. [[CrossRef](#)] [[PubMed](#)]
87. Hardin, S.C.; Larue, C.T.; Oh, M.-H.; Jain, V.; Huber, S.C. Coupling oxidative signals to protein phosphorylation via methionine oxidation in Arabidopsis. *Biochem. J.* **2009**, *422*, 305–312. [[CrossRef](#)] [[PubMed](#)]
88. Yi, D.; Alvim Kamei, C.L.; Cools, T.; Vanderauwera, S.; Takahashi, N.; Okushima, Y.; Eekhout, T.; Yoshiyama, K.O.; Larkin, J.; Van den Daele, H.; et al. The Arabidopsis SIAMESE-RELATED cyclin-dependent kinase inhibitors SMR5 and SMR7 regulate the DNA damage checkpoint in response to reactive oxygen species. *Plant Cell* **2014**, *26*, 296–309. [[CrossRef](#)] [[PubMed](#)]
89. Miao, Y.; Zentgraf, U. A HECT E3 ubiquitin ligase negatively regulates Arabidopsis leaf senescence through degradation of the transcription factor WRKY53. *Plant J.* **2010**, *63*, 179–188. [[CrossRef](#)] [[PubMed](#)]
90. Černý, M.; Novák, J.; Habánová, H.; Cerna, H.; Brzobohatý, B. Role of the proteome in phytohormonal signaling. *Biochim. Biophys. Acta-Proteins Proteom.* **2016**, *1864*, 1003–1015. [[CrossRef](#)] [[PubMed](#)]
91. Schöffl, F.; Rieping, M.; Baumann, G.; Bevan, M.; Angermüller, S. The function of plant heat shock promoter elements in the regulated expression of chimaeric genes in transgenic tobacco. *Mol. Gen. Genet.* **1989**, *217*, 246–253. [[CrossRef](#)] [[PubMed](#)]
92. Panchuk, I.I.; Volkov, R.A.; Schöffl, F. Heat stress- and heat shock transcription factor-dependent expression and activity of ascorbate peroxidase in Arabidopsis. *Plant Physiol.* **2002**, *129*, 838–853. [[CrossRef](#)] [[PubMed](#)]

93. Miller, G.; Mittler, R. Could Heat Shock Transcription Factors Function as Hydrogen Peroxide Sensors in Plants? *Ann. Bot.* **2006**, *98*, 279–288. [[CrossRef](#)] [[PubMed](#)]
94. Nishizawa, A.; Yabuta, Y.; Yoshida, E.; Maruta, T.; Yoshimura, K.; Shigeoka, S. Arabidopsis heat shock transcription factor A2 as a key regulator in response to several types of environmental stress. *Plant J.* **2006**, *48*, 535–547. [[CrossRef](#)] [[PubMed](#)]
95. Balazadeh, S.; Wu, A.; Mueller-Roeber, B. Salt-triggered expression of the ANAC092-dependent senescence regulon in *Arabidopsis thaliana*. *Plant Signal. Behav.* **2010**, *5*, 733–735. [[CrossRef](#)] [[PubMed](#)]
96. Wu, A.; Allu, A.D.; Garapati, P.; Siddiqui, H.; Dortay, H.; Zanon, M.-I.; Asensi-Fabado, M.A.; Munné-Bosch, S.; Antonio, C.; Tohge, T.; et al. JUNGBRUNNEN1, a reactive oxygen species-responsive NAC transcription factor, regulates longevity in Arabidopsis. *Plant Cell* **2012**, *24*, 482–506. [[CrossRef](#)] [[PubMed](#)]
97. Balazadeh, S.; Kwasniewski, M.; Caldana, C.; Mehrnia, M.; Zanon, M.I.; Xue, G.-P.; Mueller-Roeber, B. ORS1, an H<sub>2</sub>O<sub>2</sub>-responsive NAC transcription factor, controls senescence in *Arabidopsis thaliana*. *Mol. Plant* **2011**, *4*, 346–360. [[CrossRef](#)] [[PubMed](#)]
98. Ng, S.; Ivanova, A.; Duncan, O.; Law, S.R.; Van Aken, O.; De Clercq, I.; Wang, Y.; Carrie, C.; Xu, L.; Kmiec, B.; et al. A Membrane-Bound NAC Transcription Factor, ANAC017, Mediates Mitochondrial Retrograde Signaling in Arabidopsis. *Plant Cell* **2013**, *25*, 3450–3471. [[CrossRef](#)] [[PubMed](#)]
99. De Clercq, I.; Vermeirssen, V.; Van Aken, O.; Vandepoele, K.; Murcha, M.W.; Law, S.R.; Inzé, A.; Ng, S.; Ivanova, A.; Rombaut, D.; et al. The membrane-bound NAC transcription factor ANAC013 functions in mitochondrial retrograde regulation of the oxidative stress response in Arabidopsis. *Plant Cell* **2013**, *25*, 3472–3490. [[CrossRef](#)] [[PubMed](#)]
100. Shaikhali, J.; Davoine, C.; Brännström, K.; Rouhier, N.; Bygdell, J.; Björklund, S.; Wingsle, G. Biochemical and redox characterization of the mediator complex and its associated transcription factor GeBPL, a GLABROUS1 enhancer binding protein. *Biochem. J.* **2015**, *468*, 385–400. [[CrossRef](#)] [[PubMed](#)]
101. Shaikhali, J.; Davoine, C.; Björklund, S.; Wingsle, G. Redox regulation of the MED28 and MED32 mediator subunits is important for development and senescence. *Protoplasma* **2016**, *253*, 957–963. [[CrossRef](#)] [[PubMed](#)]
102. Phukan, U.J.; Jeena, G.S.; Shukla, R.K. WRKY Transcription Factors: Molecular Regulation and Stress Responses in Plants. *Front. Plant Sci.* **2016**, *7*, 760. [[CrossRef](#)] [[PubMed](#)]
103. Besseau, S.; Li, J.; Palva, E.T. WRKY54 and WRKY70 co-operate as negative regulators of leaf senescence in *Arabidopsis thaliana*. *J. Exp. Bot.* **2012**, *63*, 2667–2679. [[CrossRef](#)] [[PubMed](#)]
104. Ding, Z.J.; Yan, J.Y.; Xu, X.Y.; Yu, D.Q.; Li, G.X.; Zhang, S.Q.; Zheng, S.J. Transcription factor WRKY46 regulates osmotic stress responses and stomatal movement independently in Arabidopsis. *Plant J.* **2014**, *79*, 13–27. [[CrossRef](#)] [[PubMed](#)]
105. Ciftci-Yilmaz, S.; Morsy, M.R.; Song, L.; Coutu, A.; Krizek, B.A.; Lewis, M.W.; Warren, D.; Cushman, J.; Connolly, E.L.; Mittler, R. The EAR-motif of the Cys2/His2-type zinc finger protein Zat7 plays a key role in the defense response of Arabidopsis to salinity stress. *J. Biol. Chem.* **2007**, *282*, 9260–9268. [[CrossRef](#)] [[PubMed](#)]
106. Le, C.T.T.; Brumbarova, T.; Ivanov, R.; Stoof, C.; Weber, E.; Mohrbacher, J.; Fink-Straube, C.; Bauer, P. ZINC FINGER OF ARABIDOPSIS THALIANA12 (ZAT12) Interacts with FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR (FIT) Linking Iron Deficiency and Oxidative Stress Responses. *Plant Physiol.* **2016**, *170*, 540–557. [[CrossRef](#)] [[PubMed](#)]
107. Pavet, V.; Olmos, E.; Kiddle, G.; Mowla, S.; Kumar, S.; Antoniw, J.; Alvarez, M.E.; Foyer, C.H. Ascorbic Acid Deficiency Activates Cell Death and Disease Resistance Responses in Arabidopsis. *Plant Physiol.* **2005**, *139*, 1291–1303. [[CrossRef](#)] [[PubMed](#)]
108. Scrase-Field, S.A.M.G.; Knight, M.R. Calcium: Just a chemical switch? *Curr. Opin. Plant Biol.* **2003**, *6*, 500–506. [[CrossRef](#)]
109. Peiter, E. The Ever-Closer Union of Signals: Propagating Waves of Calcium and ROS Are Inextricably Linked. *Plant Physiol.* **2016**, *172*, 3–4. [[CrossRef](#)] [[PubMed](#)]
110. Pei, Z.-M.; Murata, Y.; Benning, G.; Thomine, S.; Klüsener, B.; Allen, G.J.; Grill, E.; Schroeder, J.I. Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* **2000**, *406*, 731–734. [[CrossRef](#)] [[PubMed](#)]
111. Foreman, J.; Demidchik, V.; Bothwell, J.H.F.; Mylona, P.; Miedema, H.; Torres, M.A.; Linstead, P.; Costa, S.; Brownlee, C.; Jones, J.D.G.; et al. Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* **2003**, *422*, 442–446. [[CrossRef](#)] [[PubMed](#)]



112. Yang, T.; Poovaiah, B.W. Hydrogen peroxide homeostasis: Activation of plant catalase by calcium/calmodulin. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 4097–4102. [[CrossRef](#)] [[PubMed](#)]
113. Mandadi, K.K.; Misra, A.; Ren, S.; McKnight, T.D. BT2, a BTB protein, mediates multiple responses to nutrients, stresses and hormones in Arabidopsis. *Plant Physiol.* **2009**, *150*, 1930–1939. [[CrossRef](#)] [[PubMed](#)]
114. Hua, Z.; Vierstra, R.D. The Cullin-RING Ubiquitin-Protein Ligases. *Annu. Rev. Plant Biol.* **2011**, *62*, 299–334. [[CrossRef](#)] [[PubMed](#)]
115. Grant, M.; Brown, I.; Adams, S.; Knight, M.; Ainslie, A.; Mansfield, J. The RPM1 plant disease resistance gene facilitates a rapid and sustained increase in cytosolic calcium that is necessary for the oxidative burst and hypersensitive cell death. *Plant J.* **2000**, *23*, 441–450. [[CrossRef](#)] [[PubMed](#)]
116. Vanderauwera, S.; Zimmermann, P.; Rombauts, S.; Vandenabeele, S.; Langebartels, C.; Grissem, W.; Inzé, D.; Van Breusegem, F. Genome-wide analysis of hydrogen peroxide-regulated gene expression in Arabidopsis reveals a high light-induced transcriptional cluster involved in anthocyanin biosynthesis. *Plant Physiol.* **2005**, *139*, 806–821. [[CrossRef](#)] [[PubMed](#)]
117. Desikan, R.; Hancock, J.T.; Bright, J.; Harrison, J.; Weir, I.; Hooley, R.; Neill, S.J. A role for ETR1 in hydrogen peroxide signaling in stomatal guard cells. *Plant Physiol.* **2005**, *137*, 831–834. [[CrossRef](#)] [[PubMed](#)]
118. Tognetti, V.B.; Van Aken, O.; Morreel, K.; Vandenbroucke, K.; van de Cotte, B.; De Clercq, I.; Chiwocha, S.; Fenske, R.; Prinsen, E.; Boerjan, W.; et al. Perturbation of Indole-3-Butyric Acid Homeostasis by the UDP-Glucosyltransferase UGT74E2 Modulates Arabidopsis Architecture and Water Stress Tolerance. *Plant Cell* **2010**, *22*, 2660–2679. [[CrossRef](#)] [[PubMed](#)]
119. Zhang, S.; Wu, J.; Yuan, D.; Zhang, D.; Huang, Z.; Xiao, L.; Yang, C. Perturbation of auxin homeostasis caused by mitochondrial FtSH4 gene-mediated peroxidase accumulation regulates arabidopsis architecture. *Mol. Plant.* **2014**, *7*, 856–873. [[CrossRef](#)] [[PubMed](#)]
120. Mashiguchi, K.; Tanaka, K.; Sakai, T.; Sugawara, S.; Kawaide, H.; Natsume, M.; Hanada, A.; Yaeno, T.; Shirasu, K.; Yao, H.; et al. The main auxin biosynthesis pathway in Arabidopsis. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 18512–18517. [[CrossRef](#)] [[PubMed](#)]
121. Komatsu, S.; Han, C.; Nanjo, Y.; Altaf-Un-Nahar, M.; Wang, K.; He, D.; Yang, P. Label-free quantitative proteomic analysis of abscisic acid effect in early-stage soybean under flooding. *J. Proteome Res.* **2013**, *12*, 4769–4784. [[CrossRef](#)] [[PubMed](#)]
122. Gfeller, A.; Baerenfaller, K.; Loscos, J.; Chételat, A.; Baginsky, S.; Farmer, E.E. Jasmonate controls polypeptide patterning in undamaged tissue in wounded Arabidopsis leaves. *Plant Physiol.* **2011**, *156*, 1797–1807. [[CrossRef](#)] [[PubMed](#)]
123. Černý, M.; Jedelský, P.L.; Novák, J.; Schlosser, A.; Brzobohatý, B. Cytokinin modulates proteomic, transcriptomic and growth responses to temperature shocks in Arabidopsis. *Plant Cell Environ.* **2014**, *37*, 1641–1655. [[CrossRef](#)] [[PubMed](#)]
124. Ždárská, M.; Zatloukalová, P.; Benítez, M.; Šedo, O.; Potěšil, D.; Novák, O.; Svačinová, J.; Pešek, B.; Malbeck, J.; Vašíčková, J.; et al. Proteome analysis in Arabidopsis reveals shoot- and root-specific targets of cytokinin action and differential regulation of hormonal homeostasis. *Plant Physiol.* **2013**, *161*, 918–930. [[CrossRef](#)] [[PubMed](#)]
125. Zhu, M.; Dai, S.; Zhu, N.; Booy, A.; Simons, B.; Yi, S.; Chen, S. Methyl jasmonate responsive proteins in Brassica napus guard cells revealed by iTRAQ-based quantitative proteomics. *J. Proteome Res.* **2012**, *11*, 3728–3742. [[CrossRef](#)] [[PubMed](#)]
126. Zhu, M.; Zhu, N.; Song, W.; Harmon, A.C.; Assmann, S.M.; Chen, S. Thiol-based redox proteins in abscisic acid and methyl jasmonate signaling in Brassica napus guard cells. *Plant J.* **2014**, *78*, 491–515. [[CrossRef](#)] [[PubMed](#)]
127. Cheng, F.; Blackburn, K.; Lin, Y.; Goshe, M.B.; Williamson, J.D. Absolute protein quantification by LC/MS(E) for global analysis of salicylic acid-induced plant protein secretion responses. *J. Proteome Res.* **2009**, *8*, 82–93. [[CrossRef](#)] [[PubMed](#)]
128. Li, B.; Takahashi, D.; Kawamura, Y.; Uemura, M. Comparison of plasma membrane proteomic changes of Arabidopsis suspension-cultured cells (T87 Line) after cold and ABA treatment in association with freezing tolerance development. *Plant Cell Physiol.* **2012**, *53*, 543–554. [[CrossRef](#)] [[PubMed](#)]
129. Lochmanová, G.; Zdráhal, Z.; Konečná, H.; Koukalová, Š.; Malbeck, J.; Souček, P.; Válková, M.; Kiran, N.S.; Brzobohatý, B. Cytokinin-induced photomorphogenesis in dark-grown Arabidopsis: A proteomic analysis. *J. Exp. Bot.* **2008**, *59*, 3705–3719. [[CrossRef](#)] [[PubMed](#)]



130. Proietti, S.; Bertini, L.; Timperio, A.M.; Zolla, L.; Caporale, C.; Caruso, C. Crosstalk between salicylic acid and jasmonate in *Arabidopsis* investigated by an integrated proteomic and transcriptomic approach. *Mol. Biosyst.* **2013**, *9*, 1169–1187. [[CrossRef](#)] [[PubMed](#)]
131. Böhmer, M.; Schroeder, J.I. Quantitative transcriptomic analysis of abscisic acid-induced and reactive oxygen species-dependent expression changes and proteomic profiling in *Arabidopsis* suspension cells. *Plant J.* **2011**, *67*, 105–118. [[CrossRef](#)] [[PubMed](#)]
132. Zhang, Y.; Liu, S.; Dai, S.Y.; Yuan, J.S. Integration of shot-gun proteomics and bioinformatics analysis to explore plant hormone responses. *BMC Bioinform.* **2012**, *13*, S8. [[CrossRef](#)] [[PubMed](#)]
133. Xing, M.; Xue, H. A proteomics study of auxin effects in *Arabidopsis thaliana*. *Acta Biochim. Biophys. Sin. (Shanghai)* **2012**, *44*, 783–796. [[CrossRef](#)] [[PubMed](#)]
134. Dueckershoff, K.; Mueller, S.; Mueller, M.J.; Reinders, J. Impact of cyclopentenone-oxylipins on the proteome of *Arabidopsis thaliana*. *Biochim. Biophys. Acta* **2008**, *1784*, 1975–1985. [[CrossRef](#)] [[PubMed](#)]
135. Černý, M.; Dyčka, F.; Bobál'ová, J.; Brzobohatý, B. Early cytokinin response proteins and phosphoproteins of *Arabidopsis thaliana* identified by proteome and phosphoproteome profiling. *J. Exp. Bot.* **2011**, *62*, 921–937. [[CrossRef](#)] [[PubMed](#)]
136. Rajjou, L.; Belghazi, M.; Huguet, R.; Robin, C.; Moreau, A.; Job, C.; Job, D. Proteomic investigation of the effect of salicylic acid on *Arabidopsis* seed germination and establishment of early defense mechanisms. *Plant Physiol.* **2006**, *141*, 910–923. [[CrossRef](#)] [[PubMed](#)]
137. Li, Z.; Czarnecki, O.; Chourey, K.; Yang, J.; Tuskan, G.A.; Hurst, G.B.; Pan, C.; Chen, J.-G. Strigolactone-regulated proteins revealed by iTRAQ-based quantitative proteomics in *Arabidopsis*. *J. Proteome Res.* **2014**, *13*, 1359–1372. [[CrossRef](#)] [[PubMed](#)]
138. Shigeta, T.; Yasuda, D.; Mori, T.; Yoshimitsu, Y.; Nakamura, Y.; Yoshida, S.; Asami, T.; Okamoto, S.; Matsuo, T. Characterization of brassinosteroid-regulated proteins in a nuclear-enriched fraction of *Arabidopsis* suspension-cultured cells. *Plant Physiol. Biochem.* **2011**, *49*, 985–995. [[CrossRef](#)] [[PubMed](#)]
139. Wang, P.; Xue, L.; Batelli, G.; Lee, S.; Hou, Y.-J.; Van Oosten, M.J.; Zhang, H.; Tao, W.A.; Zhu, J.-K. Quantitative phosphoproteomics identifies SnRK2 protein kinase substrates and reveals the effectors of abscisic acid action. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 11205–11210. [[CrossRef](#)] [[PubMed](#)]
140. Jourdan, N.; Martino, C.F.; El-Esawi, M.; Witczak, J.; Bouchet, P.-E.; d'Harlingue, A.; Ahmad, M. Blue-light dependent ROS formation by *Arabidopsis* cryptochrome-2 may contribute toward its signaling role. *Plant Signal. Behav.* **2015**, *10*, e1042647. [[CrossRef](#)] [[PubMed](#)]
141. El-Esawi, M.; Arthaut, L.-D.; Jourdan, N.; d'Harlingue, A.; Link, J.; Martino, C.F.; Ahmad, M. Blue-light induced biosynthesis of ROS contributes to the signaling mechanism of *Arabidopsis* cryptochrome. *Sci. Rep.* **2017**, *7*, 13875. [[CrossRef](#)] [[PubMed](#)]
142. Consentino, L.; Lambert, S.; Martino, C.; Jourdan, N.; Bouchet, P.-E.; Witczak, J.; Castello, P.; El-Esawi, M.; Corbineau, F.; d'Harlingue, A.; et al. Blue-light dependent reactive oxygen species formation by *Arabidopsis* cryptochrome may define a novel evolutionarily conserved signaling mechanism. *New Phytol.* **2015**, *206*, 1450–1462. [[CrossRef](#)] [[PubMed](#)]
143. Ha, J.-H.; Kim, J.-H.; Kim, S.-G.; Sim, H.-J.; Lee, G.; Halitschke, R.; Baldwin, I.T.; Kim, J.-I.; Park, C.-M. Shoot phytochrome B modulates reactive oxygen species homeostasis in roots via abscisic acid signaling in *Arabidopsis*. *Plant J.* **2018**, *94*, 790–798. [[CrossRef](#)] [[PubMed](#)]
144. Rajjou, L.; Debeaujon, I. Seed longevity: Survival and maintenance of high germination ability of dry seeds. *C. R. Biol.* **2008**, *331*, 796–805. [[CrossRef](#)] [[PubMed](#)]
145. Yin, X.; He, D.; Gupta, R.; Yang, P. Physiological and proteomic analyses on artificially aged *Brassica napus* seed. *Front. Plant Sci.* **2015**, *6*, 112. [[CrossRef](#)] [[PubMed](#)]
146. KOCSY, G. Die or survive? Redox changes as seed viability markers. *Plant Cell Environ.* **2015**, *38*, 1008–1010. [[CrossRef](#)] [[PubMed](#)]
147. Nguyen, T.-P.; Cuff, G.; Hegedus, D.D.; Rajjou, L.; Bentsink, L. A role for seed storage proteins in *Arabidopsis* seed longevity. *J. Exp. Bot.* **2015**, *66*, 6399–6413. [[CrossRef](#)] [[PubMed](#)]
148. Gomes, M.; Garcia, Q. Reactive oxygen species and seed germination. *Biologia* **2013**, *68*, 351–357. [[CrossRef](#)]
149. Zhang, Y.; Chen, B.; Xu, Z.; Shi, Z.; Chen, S.; Huang, X.; Chen, J.; Wang, X. Involvement of reactive oxygen species in endosperm cap weakening and embryo elongation growth during lettuce seed germination. *J. Exp. Bot.* **2014**, *65*, 3189–3200. [[CrossRef](#)] [[PubMed](#)]

150. Oracz, K.; Karpiński, S. Phytohormones Signaling Pathways and ROS Involvement in Seed Germination. *Front. Plant Sci.* **2016**, *7*, 864. [[CrossRef](#)] [[PubMed](#)]
151. Kai, K.; Kasa, S.; Sakamoto, M.; Aoki, N.; Watabe, G.; Yuasa, T.; Iwaya-Inoue, M.; Ishibashi, Y. Role of reactive oxygen species produced by NADPH oxidase in gibberellin biosynthesis during barley seed germination. *Plant Signal. Behav.* **2016**, *11*, e1180492. [[CrossRef](#)] [[PubMed](#)]
152. Ishibashi, Y.; Aoki, N.; Kasa, S.; Sakamoto, M.; Kai, K.; Tomokiyo, R.; Watabe, G.; Yuasa, T.; Iwaya-Inoue, M. The Interrelationship between Abscisic Acid and Reactive Oxygen Species Plays a Key Role in Barley Seed Dormancy and Germination. *Front. Plant Sci.* **2017**, *8*, 275. [[CrossRef](#)] [[PubMed](#)]
153. Lariguet, P.; Ranocha, P.; De Meyer, M.; Barbier, O.; Penel, C.; Dunand, C. Identification of a hydrogen peroxide signalling pathway in the control of light-dependent germination in Arabidopsis. *Planta* **2013**, *238*, 381–395. [[CrossRef](#)] [[PubMed](#)]
154. Wojtyła, Ł.; Lechowska, K.; Kubala, S.; Garnczarska, M. Different Modes of Hydrogen Peroxide Action During Seed Germination. *Front. Plant Sci.* **2016**, *7*, 66. [[CrossRef](#)] [[PubMed](#)]
155. Barba-Espín, G.; Hernández, J.A.; Diaz-Vivancos, P. Role of H<sub>2</sub>O<sub>2</sub> in pea seed germination. *Plant Signal. Behav.* **2012**, *7*, 193–195. [[CrossRef](#)] [[PubMed](#)]
156. Bazin, J.; Langlade, N.; Vincourt, P.; Arribat, S.; Balzergue, S.; El-Maarouf-Bouteau, H.; Bailly, C. Targeted mRNA Oxidation Regulates Sunflower Seed Dormancy Alleviation during Dry After-Ripening. *Plant Cell* **2011**, *23*, 2196–2208. [[CrossRef](#)] [[PubMed](#)]
157. El-Maarouf-Bouteau, H.; Meimoun, P.; Job, C.; Job, D.; Bailly, C. Role of protein and mRNA oxidation in seed dormancy and germination. *Front. Plant Sci.* **2013**, *4*, 77. [[CrossRef](#)] [[PubMed](#)]
158. Ishibashi, Y.; Kasa, S.; Sakamoto, M.; Aoki, N.; Kai, K.; Yuasa, T.; Hanada, A.; Yamaguchi, S.; Iwaya-Inoue, M. A Role for Reactive Oxygen Species Produced by NADPH Oxidases in the Embryo and Aleurone Cells in Barley Seed Germination. *PLoS ONE* **2015**, *10*, e0143173. [[CrossRef](#)] [[PubMed](#)]
159. Aoki, N.; Ishibashi, Y.; Kai, K.; Tomokiyo, R.; Yuasa, T.; Iwaya-Inoue, M. Programmed cell death in barley aleurone cells is not directly stimulated by reactive oxygen species produced in response to gibberellin. *J. Plant Physiol.* **2014**, *171*, 615–618. [[CrossRef](#)] [[PubMed](#)]
160. Ishibashi, Y.; Tawaratsumida, T.; Kondo, K.; Kasa, S.; Sakamoto, M.; Aoki, N.; Zheng, S.-H.; Yuasa, T.; Iwaya-Inoue, M. Reactive Oxygen Species Are Involved in Gibberellin/Abscisic Acid Signaling in Barley Aleurone Cells. *Plant Physiol.* **2012**, *158*, 1705–1714. [[CrossRef](#)] [[PubMed](#)]
161. Farouk, S.; Abdul Qados, A.M.S. Enhancing seed quality and productivity as well as physio-anatomical responses of pea plants by folic acid and/or hydrogen peroxide application. *Sci. Hortic. (Amsterdam)* **2018**, *240*, 29–37. [[CrossRef](#)]
162. Bouallègue, A.; Souissi, F.; Nouairi, I.; Souibgui, M.; Abbes, Z.; Mhadhbi, H. Salicylic acid and hydrogen peroxide pretreatments alleviate salt stress in faba bean (*Vicia faba*) seeds during germination. *Seed Sci. Technol.* **2017**, *45*, 675–690. [[CrossRef](#)]
163. Li, B.; Cai, Q.; Ma, S.; Li, S.; Zhang, X.; Yu, Y. Regulation of NPA and ACC on H<sub>2</sub>O<sub>2</sub>-Induced Pea Primary Horizontal Bending Root. *J. Plant Growth Regul.* **2018**, *37*, 246–254. [[CrossRef](#)]
164. Carol, R.J.; Takeda, S.; Linstead, P.; Durrant, M.C.; Kakesova, H.; Derbyshire, P.; Drea, S.; Zarsky, V.; Dolan, L. A RhoGDP dissociation inhibitor spatially regulates growth in root hair cells. *Nature* **2005**, *438*, 1013–1016. [[CrossRef](#)] [[PubMed](#)]
165. Takeda, S.; Gapper, C.; Kaya, H.; Bell, E.; Kuchitsu, K.; Dolan, L. Local positive feedback regulation determines cell shape in root hair cells. *Science* **2008**, *319*, 1241–1244. [[CrossRef](#)] [[PubMed](#)]
166. Qu, Y.; Wang, Q.; Guo, J.; Wang, P.; Song, P.; Jia, Q.; Zhang, X.; Kudla, J.; Zhang, W.; Zhang, Q. Peroxisomal CuAOZ and its product H<sub>2</sub>O<sub>2</sub> regulate the distribution of auxin and IBA-dependent lateral root development in Arabidopsis. *J. Exp. Bot.* **2017**, *68*, 4851–4867. [[CrossRef](#)] [[PubMed](#)]
167. Su, C.; Liu, L.; Liu, H.; Ferguson, B.J.; Zou, Y.; Zhao, Y.; Wang, T.; Wang, Y.; Li, X. H<sub>2</sub>O<sub>2</sub> regulates root system architecture by modulating the polar transport and redistribution of auxin. *J. Plant Biol.* **2016**, *59*, 260–270. [[CrossRef](#)]
168. Takáč, T.; Obert, B.; Rolčík, J.; Šamaj, J. Improvement of adventitious root formation in flax using hydrogen peroxide. *New Biotechnol.* **2016**, *33*, 728–734. [[CrossRef](#)] [[PubMed](#)]
169. Chen, Z.; Gu, Q.; Yu, X.; Huang, L.; Xu, S.; Wang, R.; Shen, W.; Shen, W. Hydrogen peroxide acts downstream of melatonin to induce lateral root formation. *Ann. Bot.* **2018**, *121*, 1127–1136. [[CrossRef](#)] [[PubMed](#)]

170. Xiong, J.; Yang, Y.; Fu, G.; Tao, L. Novel roles of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) in regulating pectin synthesis and demethylesterification in the cell wall of rice (*Oryza sativa*) root tips. *New Phytol.* **2015**, *206*, 118–126. [[CrossRef](#)] [[PubMed](#)]
171. Ivanchenko, M.G.; den Os, D.; Monshausen, G.B.; Dubrovsky, J.G.; Bednárová, A.; Krishnan, N. Auxin increases the hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) concentration in tomato (*Solanum lycopersicum*) root tips while inhibiting root growth. *Ann. Bot.* **2013**, *112*, 1107–1116. [[CrossRef](#)] [[PubMed](#)]
172. Zhou, L.; Hou, H.; Yang, T.; Lian, Y.; Sun, Y.; Bian, Z.; Wang, C. Exogenous hydrogen peroxide inhibits primary root gravitropism by regulating auxin distribution during Arabidopsis seed germination. *Plant Physiol. Biochem. PPB* **2018**, *128*, 126–133. [[CrossRef](#)] [[PubMed](#)]
173. Jiao, Y.; Sun, L.; Song, Y.; Wang, L.; Liu, L.; Zhang, L.; Liu, B.; Li, N.; Miao, C.; Hao, F. AtrbohD and AtrbohF positively regulate abscisic acid-inhibited primary root growth by affecting Ca<sup>2+</sup> signalling and auxin response of roots in Arabidopsis. *J. Exp. Bot.* **2013**, *64*, 4183–4192. [[CrossRef](#)] [[PubMed](#)]
174. He, H.; Yan, J.; Yu, X.; Liang, Y.; Fang, L.; Scheller, H.V.; Zhang, A. The NADPH-oxidase AtRbohI plays a positive role in drought-stress response in Arabidopsis thaliana. *Biochem. Biophys. Res. Commun.* **2017**, *491*, 834–839. [[CrossRef](#)] [[PubMed](#)]
175. Li, H.; He, J.; Yang, X.; Li, X.; Luo, D.; Wei, C.; Ma, J.; Zhang, Y.; Yang, J.; Zhang, X. Glutathione-dependent induction of local and systemic defense against oxidative stress by exogenous melatonin in cucumber (*Cucumis sativus* L.). *J. Pineal Res.* **2016**, *60*, 206–216. [[CrossRef](#)] [[PubMed](#)]
176. Kong, X.; Luo, Z.; Dong, H.; Eneji, A.E.; Li, W. H<sub>2</sub>O<sub>2</sub> and ABA signaling are responsible for the increased Na<sup>+</sup> efflux and water uptake in *Gossypium hirsutum* L. roots in the non-saline side under non-uniform root zone salinity. *J. Exp. Bot.* **2016**, *67*, 2247–2261. [[CrossRef](#)] [[PubMed](#)]
177. Yamauchi, T.; Yoshioka, M.; Fukazawa, A.; Mori, H.; Nishizawa, N.K.; Tsutsumi, N.; Yoshioka, H.; Nakazono, M. An NADPH Oxidase RBOH Functions in Rice Roots during Lysigenous Aerenchyma Formation under Oxygen-Deficient Conditions. *Plant Cell* **2017**, *29*, 775–790. [[CrossRef](#)] [[PubMed](#)]
178. Mangano, S.; Denita-Juarez, S.P.; Choi, H.-S.; Marzol, E.; Hwang, Y.; Ranocha, P.; Velasquez, S.M.; Borassi, C.; Barberini, M.L.; Aptekmann, A.A.; et al. Molecular link between auxin and ROS-mediated polar growth. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 5289–5294. [[CrossRef](#)] [[PubMed](#)]
179. Voxeur, A.; Höfte, H. Cell wall integrity signaling in plants: “To grow or not to grow that’s the question”. *Glycobiology* **2016**, *26*, 950–960. [[CrossRef](#)] [[PubMed](#)]
180. Chen, X.-J.; Xia, X.-J.; Guo, X.; Zhou, Y.-H.; Shi, K.; Zhou, J.; Yu, J.-Q. Apoplastic H<sub>2</sub>O<sub>2</sub> plays a critical role in axillary bud outgrowth by altering auxin and cytokinin homeostasis in tomato plants. *New Phytol.* **2016**, *211*, 1266–1278. [[CrossRef](#)] [[PubMed](#)]
181. Guo, Z.; Wang, F.; Xiang, X.; Ahammed, G.J.; Wang, M.; Onac, E.; Zhou, J.; Xia, X.; Shi, K.; Yin, X.; et al. Systemic Induction of Photosynthesis via Illumination of the Shoot Apex Is Mediated Sequentially by Phytochrome, B<sub>1</sub>, Auxin and Hydrogen Peroxide in Tomato. *Plant Physiol.* **2016**, *172*, 1259–1272. [[CrossRef](#)] [[PubMed](#)]
182. Sandalio, L.M.; Rodríguez-Serrano, M.; Romero-Puertas, M.C. Leaf epinasty and auxin: A biochemical and molecular overview. *Plant Sci.* **2016**, *253*, 187–193. [[CrossRef](#)] [[PubMed](#)]
183. Devireddy, A.R.; Zandalinas, S.I.; Gómez-Cadenas, A.; Blumwald, E.; Mittler, R. Coordinating the overall stomatal response of plants: Rapid leaf-to-leaf communication during light stress. *Sci. Signal.* **2018**, *11*, eaam9514. [[CrossRef](#)] [[PubMed](#)]
184. Ha, Y.; Shang, Y.; Nam, K.H. Brassinosteroids modulate ABA-induced stomatal closure in Arabidopsis. *J. Exp. Bot.* **2016**, *67*, 6297–6308. [[CrossRef](#)] [[PubMed](#)]
185. Xia, X.-J.; Gao, C.-J.; Song, L.-X.; Zhou, Y.-H.; Shi, K.; Yu, J.-Q. Role of H<sub>2</sub>O<sub>2</sub> dynamics in brassinosteroid-induced stomatal closure and opening in *Solanum lycopersicum*. *Plant Cell Environ.* **2014**, *37*, 2036–2050. [[CrossRef](#)] [[PubMed](#)]
186. Lv, S.; Zhang, Y.; Li, C.; Liu, Z.; Yang, N.; Pan, L.; Wu, J.; Wang, J.; Yang, J.; Lv, Y.; et al. Strigolactone-triggered stomatal closure requires hydrogen peroxide synthesis and nitric oxide production in an abscisic acid-independent manner. *New Phytol.* **2018**, *217*, 290–304. [[CrossRef](#)] [[PubMed](#)]
187. Khokon, M.A.R.; Salam, M.A.; Jammes, F.; Ye, W.; Hossain, M.A.; Okuma, E.; Nakamura, Y.; Mori, I.C.; Kwak, J.M.; Murata, Y. MPK9 and MPK12 function in SA-induced stomatal closure in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* **2017**, *81*, 1394–1400. [[CrossRef](#)] [[PubMed](#)]

188. Ehonen, S.; Yarmolinsky, D.; Kollist, H.; Kangasjärvi, J. Reactive Oxygen Species, Photosynthesis and Environment in the Regulation of Stomata. *Antioxid. Redox Signal.* **2017**, ars.2017.7455. [[CrossRef](#)]
189. Ge, X.-M.; Cai, H.-L.; Lei, X.; Zhou, X.; Yue, M.; He, J.-M. Heterotrimeric G protein mediates ethylene-induced stomatal closure via hydrogen peroxide synthesis in Arabidopsis. *Plant J.* **2015**, *82*, 138–150. [[CrossRef](#)] [[PubMed](#)]
190. Shi, C.; Qi, C.; Ren, H.; Huang, A.; Hei, S.; She, X. Ethylene mediates brassinosteroid-induced stomatal closure via G $\alpha$  protein-activated hydrogen peroxide and nitric oxide production in Arabidopsis. *Plant J.* **2015**, *82*, 280–301. [[CrossRef](#)] [[PubMed](#)]
191. Li, Y.; Xu, S.; Wang, Z.; He, L.; Xu, K.; Wang, G. Glucose triggers stomatal closure mediated by basal signaling through HXK1 and PYR/RCAR receptors in Arabidopsis. *J. Exp. Bot.* **2018**, *69*, 1471–1484. [[CrossRef](#)] [[PubMed](#)]
192. Nazareno, A.L.; Hernandez, B.S. A mathematical model of the interaction of abscisic acid, ethylene and methyl jasmonate on stomatal closure in plants. *PLoS ONE* **2017**, *12*, e0171065. [[CrossRef](#)] [[PubMed](#)]
193. Dubovskaya, L.V.; Bakakina, Y.S.; Kolesneva, E.V.; Sodel, D.L.; Mcainsh, M.R.; Hetherington, A.M.; Volotovskii, I.D. cGMP-dependent ABA-induced stomatal closure in the ABA-insensitive Arabidopsis mutant *abi1-1*. *New Phytol.* **2011**, *191*, 57–69. [[CrossRef](#)] [[PubMed](#)]
194. Sun, L.; Li, Y.; Miao, W.; Piao, T.; Hao, Y.; Hao, F.-S. NADK2 positively modulates abscisic acid-induced stomatal closure by affecting accumulation of H<sub>2</sub>O<sub>2</sub>, Ca<sup>2+</sup> and nitric oxide in Arabidopsis guard cells. *Plant Sci.* **2017**, *262*, 81–90. [[CrossRef](#)] [[PubMed](#)]
195. Agurla, S.; Gayatri, G.; Raghavendra, A.S. Polyamines increase nitric oxide and reactive oxygen species in guard cells of Arabidopsis thaliana during stomatal closure. *Protoplasma* **2018**, *255*, 153–162. [[CrossRef](#)] [[PubMed](#)]
196. Jardim-Messeder, D.; Caverzan, A.; Rauber, R.; Cunha, J.R.; Carvalho, F.E.L.; Gaeta, M.L.; da Fonseca, G.C.; Costa, J.M.; Frei, M.; Silveira, J.A.G.; et al. Thylakoidal APX modulates hydrogen peroxide content and stomatal closure in rice (*Oryza sativa* L.). *Environ. Exp. Bot.* **2018**, *150*, 46–56. [[CrossRef](#)]
197. Mao, X.; Zheng, Y.; Xiao, K.; Wei, Y.; Zhu, Y.; Cai, Q.; Chen, L.; Xie, H.; Zhang, J. OsPRX2 contributes to stomatal closure and improves potassium deficiency tolerance in rice. *Biochem. Biophys. Res. Commun.* **2018**, *495*, 461–467. [[CrossRef](#)] [[PubMed](#)]
198. Kimura, S.; Kaya, H.; Kawarazaki, T.; Hiraoka, G.; Senzaki, E.; Michikawa, M.; Kuchitsu, K. Protein phosphorylation is a prerequisite for the Ca<sup>2+</sup>-dependent activation of Arabidopsis NADPH oxidases and may function as a trigger for the positive feedback regulation of Ca<sup>2+</sup> and reactive oxygen species. *Biochim. Biophys. Acta* **2012**, *1823*, 398–405. [[CrossRef](#)] [[PubMed](#)]
199. Sirichandra, C.; Gu, D.; Hu, H.-C.; Davanture, M.; Lee, S.; Djaoui, M.; Valot, B.; Zivy, M.; Leung, J.; Merlot, S.; et al. Phosphorylation of the Arabidopsis AtrbohF NADPH oxidase by OST1 protein kinase. *FEBS Lett.* **2009**, *583*, 2982–2986. [[CrossRef](#)] [[PubMed](#)]
200. Rodrigues, O.; Reshetnyak, G.; Grondin, A.; Saijo, Y.; Leonhardt, N.; Maurel, C.; Verdoucq, L. Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 9200–9205. [[CrossRef](#)] [[PubMed](#)]
201. Assmann, S.M.; Jegla, T. Guard cell sensory systems: Recent insights on stomatal responses to light, abscisic acid and CO<sub>2</sub>. *Curr. Opin. Plant Biol.* **2016**, *33*, 157–167. [[CrossRef](#)] [[PubMed](#)]
202. Scuffi, D.; Nietzel, T.; Di Fino, L.M.; Meyer, A.J.; Lamattina, L.; Schwarzländer, M.; Laxalt, A.M.; García-Mata, C. Hydrogen Sulfide Increases Production of NADPH Oxidase-Dependent Hydrogen Peroxide and Phospholipase D-Derived Phosphatidic Acid in Guard Cell Signaling. *Plant Physiol.* **2018**, *176*, 2532–2542. [[CrossRef](#)] [[PubMed](#)]
203. Watkins, J.M.; Chapman, J.M.; Muday, G.K. Abscisic Acid-Induced Reactive Oxygen Species Are Modulated by Flavonols to Control Stomata Aperture. *Plant Physiol.* **2017**, *175*, 1807–1825. [[CrossRef](#)] [[PubMed](#)]
204. An, Y.; Feng, X.; Liu, L.; Xiong, L.; Wang, L. ALA-Induced Flavonols Accumulation in Guard Cells Is Involved in Scavenging H<sub>2</sub>O<sub>2</sub> and Inhibiting Stomatal Closure in Arabidopsis Cotyledons. *Front. Plant Sci.* **2016**, *7*, 1713. [[CrossRef](#)] [[PubMed](#)]
205. An, Y.; Liu, L.; Chen, L.; Wang, L. ALA Inhibits ABA-induced Stomatal Closure via Reducing H<sub>2</sub>O<sub>2</sub> and Ca(2+) Levels in Guard Cells. *Front. Plant Sci.* **2016**, *7*, 482. [[CrossRef](#)] [[PubMed](#)]



206. Serrano, I.; Romero-Puertas, M.C.; Sandalio, L.M.; Olmedilla, A. The role of reactive oxygen species and nitric oxide in programmed cell death associated with self-incompatibility. *J. Exp. Bot.* **2015**, *66*, 2869–2876. [[CrossRef](#)] [[PubMed](#)]
207. Lassig, R.; Guterath, T.; Bey, T.D.; Konrad, K.R.; Romeis, T. Pollen tube NAD(P)H oxidases act as a speed control to dampen growth rate oscillations during polarized cell growth. *Plant J.* **2014**, *78*, 94–106. [[CrossRef](#)] [[PubMed](#)]
208. Kaya, H.; Nakajima, R.; Iwano, M.; Kanaoka, M.M.; Kimura, S.; Takeda, S.; Kawarazaki, T.; Senzaki, E.; Hamamura, Y.; Higashiyama, T.; et al. Ca<sup>2+</sup>-activated reactive oxygen species production by Arabidopsis RbohH and RbohJ is essential for proper pollen tube tip growth. *Plant Cell* **2014**, *26*, 1069–1080. [[CrossRef](#)] [[PubMed](#)]
209. Duan, Q.; Kita, D.; Johnson, E.A.; Aggarwal, M.; Gates, L.; Wu, H.-M.; Cheung, A.Y. Reactive oxygen species mediate pollen tube rupture to release sperm for fertilization in Arabidopsis. *Nat. Commun.* **2014**, *5*, 3129. [[CrossRef](#)] [[PubMed](#)]
210. Huan, C.; Jiang, L.; An, X.; Yu, M.; Xu, Y.; Ma, R.; Yu, Z. Potential role of reactive oxygen species and antioxidant genes in the regulation of peach fruit development and ripening. *Plant Physiol. Biochem. PPB* **2016**, *104*, 294–303. [[CrossRef](#)] [[PubMed](#)]
211. Kumar, V.; Irfan, M.; Ghosh, S.; Chakraborty, N.; Chakraborty, S.; Datta, A. Fruit ripening mutants reveal cell metabolism and redox state during ripening. *Protoplasma* **2016**, *253*, 581–594. [[CrossRef](#)] [[PubMed](#)]
212. Hurr, B.M.; Huber, D.J.; Vallejos, C.E.; Lee, E.; Sargent, S.A. Ethylene-induced overproduction of reactive oxygen species is responsible for the development of watersoaking in immature cucumber fruit. *J. Plant Physiol.* **2013**, *170*, 56–62. [[CrossRef](#)] [[PubMed](#)]
213. Avila-Ospina, L.; Moison, M.; Yoshimoto, K.; Masclaux-Daubresse, C. Autophagy, plant senescence and nutrient recycling. *J. Exp. Bot.* **2014**, *65*, 3799–3811. [[CrossRef](#)] [[PubMed](#)]
214. Bieker, S.; Riester, L.; Stahl, M.; Franzaring, J.; Zentgraf, U. Senescence-specific Alteration of Hydrogen Peroxide Levels in *Arabidopsis thaliana* and Oilseed Rape Spring Variety Brassica napus L. cv. MozartF. *J. Integr. Plant Biol.* **2012**, *54*, 540–554. [[CrossRef](#)] [[PubMed](#)]
215. Wang, H.; Lin, J.; Chang, Y.; Jiang, C.-Z. Comparative Transcriptomic Analysis Reveals That Ethylene/H<sub>2</sub>O<sub>2</sub>-Mediated Hypersensitive Response and Programmed Cell Death Determine the Compatible Interaction of Sand Pear and *Alternaria alternata*. *Front. Plant Sci.* **2017**, *8*, 195. [[CrossRef](#)] [[PubMed](#)]
216. Liu, J.; Xu, Y.; Zhang, Z.; Wei, J. Hydrogen peroxide promotes programmed cell death and salicylic acid accumulation during the induced production of sesquiterpenes in cultured cell suspensions of *Aquilaria sinensis*. *Funct. Plant Biol.* **2015**, *42*, 337–346. [[CrossRef](#)]
217. Zimmermann, P.; Heinlein, C.; Orendi, G.; Zentgraf, U. Senescence-specific regulation of catalases in *Arabidopsis thaliana* (L.) Heynh. *Plant Cell Environ.* **2006**, *29*, 1049–1060. [[CrossRef](#)] [[PubMed](#)]
218. Jajić, I.; Sarna, T.; Szewczyk, G.; Strzałka, K. Changes in production of reactive oxygen species in illuminated thylakoids isolated during development and senescence of barley. *J. Plant Physiol.* **2015**, *184*, 49–56. [[CrossRef](#)] [[PubMed](#)]
219. Niewiadomska, E.; Polzien, L.; Desel, C.; Rozpadek, P.; Miszalski, Z.; Krupinska, K. Spatial patterns of senescence and development-dependent distribution of reactive oxygen species in tobacco (*Nicotiana tabacum*) leaves. *J. Plant Physiol.* **2009**, *166*, 1057–1068. [[CrossRef](#)] [[PubMed](#)]
220. Saxena, I.; Srikanth, S.; Chen, Z. Cross Talk between H<sub>2</sub>O<sub>2</sub> and Interacting Signal Molecules under Plant Stress Response. *Front. Plant Sci.* **2016**, *7*, 570. [[CrossRef](#)] [[PubMed](#)]
221. Diao, Q.; Song, Y.; Shi, D.; Qi, H. Interaction of Polyamines, Abscisic Acid, Nitric Oxide and Hydrogen Peroxide under Chilling Stress in Tomato (*Lycopersicon esculentum* Mill.) Seedlings. *Front. Plant Sci.* **2017**, *8*, 203. [[CrossRef](#)] [[PubMed](#)]
222. Ren, C.-G.; Kong, C.-C.; Xie, Z.-H. Role of abscisic acid in strigolactone-induced salt stress tolerance in arbuscular mycorrhizal *Sesbania cannabina* seedlings. *BMC Plant Biol.* **2018**, *18*, 74. [[CrossRef](#)] [[PubMed](#)]
223. Freitas, V.S.; de Souza Miranda, R.; Costa, J.H.; de Oliveira, D.F.; de Oliveira Paula, S.; de Castro Migueld, E.; Freire, R.S.; Prisco, J.T.; Gomes-Filho, E. Ethylene triggers salt tolerance in maize genotypes by modulating polyamine catabolism enzymes associated with H<sub>2</sub>O<sub>2</sub> production. *Environ. Exp. Bot.* **2018**, *145*, 75–86. [[CrossRef](#)]



224. Kaur, N.; Kirat, K.; Saini, S.; Sharma, I.; Gantet, P.; Pati, P.K. Reactive oxygen species generating system and brassinosteroids are linked to salt stress adaptation mechanisms in rice. *Plant Signal. Behav.* **2016**, *11*, e1247136. [[CrossRef](#)] [[PubMed](#)]
225. Yang, C.-Y.; Huang, Y.-C.; Ou, S.-L. ERF73/HRE1 is involved in H<sub>2</sub>O<sub>2</sub> production via hypoxia-inducible Rboh gene expression in hypoxia signaling. *Protoplasma* **2017**, *254*, 1705–1714. [[CrossRef](#)] [[PubMed](#)]
226. Yao, Y.; He, R.J.; Xie, Q.L.; Zhao, X.H.; Deng, X.M.; He, J.B.; Song, L.; He, J.; Marchant, A.; Chen, X.-Y.; et al. ETHYLENE RESPONSE FACTOR 74 (ERF74) plays an essential role in controlling a respiratory burst oxidase homolog D (RbohD)-dependent mechanism in response to different stresses in Arabidopsis. *New Phytol.* **2017**, *213*, 1667–1681. [[CrossRef](#)] [[PubMed](#)]
227. Zhang, M.; Smith, J.A.C.; Harberd, N.P.; Jiang, C. The regulatory roles of ethylene and reactive oxygen species (ROS) in plant salt stress responses. *Plant Mol. Biol.* **2016**, *91*, 651–659. [[CrossRef](#)] [[PubMed](#)]
228. Sun, M.; Jiang, F.; Cen, B.; Wen, J.; Zhou, Y.; Wu, Z. Respiratory burst oxidase homologue-dependent H<sub>2</sub>O<sub>2</sub> and chloroplast H<sub>2</sub>O<sub>2</sub> are essential for the maintenance of acquired thermotolerance during recovery after acclimation. *Plant Cell Environ.* **2018**. [[CrossRef](#)] [[PubMed](#)]
229. Birch, P.R.J.; Avrova, A.O.; Dellagi, A.; Lacomme, C.; Cruz, S.S.; Lyon, G.D. Programmed Cell Death in Plants in Response to Pathogen Attack. In *Annual Plant. Reviews*; John Wiley & Sons, Ltd.: Chichester, UK, 2018; pp. 184–208.
230. Novák, J.; Pavlů, J.; Novák, O.; Nožková-Hlaváčková, V.; Špundová, M.; Hlavinka, J.; Koukalová, Š.; Skalák, J.; Černý, M.; Brzobohatý, B. High cytokinin levels induce a hypersensitive-like response in tobacco. *Ann. Bot.* **2013**, *112*, 41–55. [[CrossRef](#)] [[PubMed](#)]
231. Nováková, M.; Šašek, V.; Trdá, L.; Krutinová, H.; Mongin, T.; Valentová, O.; Balesdent, M.-H.; Rouxel, T.; Burketová, L. *L. leptosphaeria maculans* effector AvrLm4-7 affects salicylic acid (SA) and ethylene (ET) signalling and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) accumulation in *Brassica napus*. *Mol. Plant Pathol.* **2016**, *17*, 818–831. [[CrossRef](#)] [[PubMed](#)]
232. Song, L.-X.; Xu, X.-C.; Wang, F.-N.; Wang, Y.; Xia, X.-J.; Shi, K.; Zhou, Y.-H.; Zhou, J.; Yu, J.-Q. Brassinosteroids act as a positive regulator for resistance against root-knot nematode involving RESPIRATORY BURST OXIDASE HOMOLOG-dependent activation of MAPKs in tomato. *Plant Cell Environ.* **2018**, *41*, 1113–1125. [[CrossRef](#)] [[PubMed](#)]
233. Deng, X.-G.; Zhu, T.; Zou, L.-J.; Han, X.-Y.; Zhou, X.; Xi, D.-H.; Zhang, D.-W.; Lin, H.-H. Orchestration of hydrogen peroxide and nitric oxide in brassinosteroid-mediated systemic virus resistance in *Nicotiana benthamiana*. *Plant J.* **2016**, *85*, 478–493. [[CrossRef](#)] [[PubMed](#)]
234. Zhu, F.; Chen, J.; Xiao, X.; Zhang, M.; Yun, Z.; Zeng, Y.; Xu, J.; Cheng, Y.; Deng, X. Salicylic acid treatment reduces the rot of postharvest citrus fruit by inducing the accumulation of H<sub>2</sub>O<sub>2</sub>, primary metabolites and lipophilic polymethoxylated flavones. *Food Chem.* **2016**, *207*, 68–74. [[CrossRef](#)] [[PubMed](#)]
235. Ellouzi, H.; Sghayar, S.; Abdelly, C. H<sub>2</sub>O<sub>2</sub> seed priming improves tolerance to salinity; drought and their combined effect more than mannitol in *Cakile maritima* when compared to *Eutrema salsugineum*. *J. Plant Physiol.* **2017**, *210*, 38–50. [[CrossRef](#)] [[PubMed](#)]
236. Richards, S.L.; Wilkins, K.A.; Swarbreck, S.M.; Anderson, A.A.; Habib, N.; Smith, A.G.; McAinsh, M.; Davies, J.M. The hydroxyl radical in plants: From seed to seed. *J. Exp. Bot.* **2015**, *66*, 37–46. [[CrossRef](#)] [[PubMed](#)]
237. Imlay, J.A. Cellular Defenses against Superoxide and Hydrogen Peroxide. *Annu. Rev. Biochem.* **2008**, *77*, 755–776. [[CrossRef](#)] [[PubMed](#)]
238. Demidchik, V.; Shabala, S.N.; Coutts, K.B.; Tester, M.A.; Davies, J.M. Free oxygen radicals regulate plasma membrane Ca<sup>2+</sup>- and K<sup>+</sup>-permeable channels in plant root cells. *J. Cell Sci.* **2003**, *116*, 81–88. [[CrossRef](#)] [[PubMed](#)]
239. Laohavisit, A.; Shang, Z.; Rubio, L.; Cuin, T.A.; Véry, A.-A.; Wang, A.; Mortimer, J.C.; Macpherson, N.; Coxon, K.M.; Battey, N.H.; et al. Arabidopsis annexin1 mediates the radical-activated plasma membrane Ca<sup>2+</sup>- and K<sup>+</sup>-permeable conductance in root cells. *Plant Cell* **2012**, *24*, 1522–1533. [[CrossRef](#)] [[PubMed](#)]
240. Demidchik, V.; Cuin, T.A.; Svistunenko, D.; Smith, S.J.; Miller, A.J.; Shabala, S.; Sokolik, A.; Yurin, V. Arabidopsis root K<sup>+</sup>-efflux conductance activated by hydroxyl radicals: Single-channel properties, genetic basis and involvement in stress-induced cell death. *J. Cell Sci.* **2010**, *123*, 1468–1479. [[CrossRef](#)] [[PubMed](#)]

241. Demidchik, V.; Straltsova, D.; Medvedev, S.S.; Pozhvanov, G.A.; Sokolik, A.; Yurin, V. Stress-induced electrolyte leakage: The role of K<sup>+</sup>-permeable channels and involvement in programmed cell death and metabolic adjustment. *J. Exp. Bot.* **2014**, *65*, 1259–1270. [[CrossRef](#)] [[PubMed](#)]
242. Demidchik, V.; Shabala, S. Mechanisms of cytosolic calcium elevation in plants: The role of ion channels, calcium extrusion systems and NADPH oxidase-mediated “ROS-Ca<sup>2+</sup> Hub”. *Funct. Plant Biol.* **2018**, *45*, 9. [[CrossRef](#)]
243. Sun, J.; Wang, M.-J.; Ding, M.-Q.; Deng, S.-R.; Liu, M.-Q.; Lu, C.-F.; Zhou, X.-Y.; Shen, X.; Zheng, X.-J.; Zhang, Z.-K.; et al. H<sub>2</sub>O<sub>2</sub> and cytosolic Ca<sup>2+</sup> signals triggered by the PM H<sup>+</sup>-coupled transport system mediate K<sup>+</sup>/Na<sup>+</sup> homeostasis in NaCl-stressed *Populus euphratica* cells. *Plant Cell Environ.* **2010**, *33*, 943–958. [[CrossRef](#)] [[PubMed](#)]
244. Gilroy, S.; Białasek, M.; Suzuki, N.; Górecka, M.; Devireddy, A.R.; Karpiński, S.; Mittler, R. ROS, Calcium and Electric Signals: Key Mediators of Rapid Systemic Signaling in Plants. *Plant Physiol.* **2016**, *171*, 1606–1615. [[CrossRef](#)] [[PubMed](#)]
245. Crawford, T.; Lehotai, N.; Strand, Å. The role of retrograde signals during plant stress responses. *J. Exp. Bot.* **2018**, *69*, 2783–2795. [[CrossRef](#)] [[PubMed](#)]
246. Exposito-Rodriguez, M.; Laissue, P.P.; Yvon-Durocher, G.; Smirnov, N.; Mullineaux, P.M. Photosynthesis-dependent H<sub>2</sub>O<sub>2</sub> transfer from chloroplasts to nuclei provides a high-light signalling mechanism. *Nat. Commun.* **2017**, *8*, 49. [[CrossRef](#)] [[PubMed](#)]
247. Pavlí, J.; Novák, J.; Koukalová, V.; Luklová, M.; Brzobohatý, B.; Černý, M. Cytokinin at the crossroad of abiotic stress signalling pathways. *Int. J. Mol. Sci.* **2018**, *19*, 2450. [[CrossRef](#)] [[PubMed](#)]
248. Waszczak, C.; Carmody, M.; Kangasjärvi, J. Reactive Oxygen Species in Plant Signaling. *Annu. Rev. Plant Biol.* **2018**, *69*, 209–236. [[CrossRef](#)] [[PubMed](#)]
249. Pratsinis, A.; Kelesidis, G.A.; Zuercher, S.; Krumeich, F.; Bolisetty, S.; Mezzenga, R.; Leroux, J.-C.; Sotiriou, G.A. Enzyme-Mimetic Antioxidant Luminescent Nanoparticles for Highly Sensitive Hydrogen Peroxide Biosensing. *ACS Nano* **2017**, *11*, 12210–12218. [[CrossRef](#)] [[PubMed](#)]
250. Neal, C.J.; Gupta, A.; Barkam, S.; Saraf, S.; Das, S.; Cho, H.J.; Seal, S. Picomolar Detection of Hydrogen Peroxide using Enzyme-free Inorganic Nanoparticle-based Sensor. *Sci. Rep.* **2017**, *7*, 1324. [[CrossRef](#)] [[PubMed](#)]
251. Shabrangy, A.; Roustan, V.; Reipert, S.; Weidinger, M.; Roustan, P.-J.; Stoger, E.; Weckwerth, W.; Ibl, V. Using RT-qPCR, Proteomics and Microscopy to Unravel the Spatio-Temporal Expression and Subcellular Localization of Hordoinolines Across Development in Barley Endosperm. *Front. Plant Sci.* **2018**, *9*, 775. [[CrossRef](#)] [[PubMed](#)]

