

RESEARCH ARTICLE

Recommended next care following hospital-treated self-harm: Patterns and trends over time

Ella Arensman^{1,2*}, Eve Griffin¹, Caroline Daly¹, Paul Corcoran^{1,2}, Eugene Cassidy^{3,4}, Ivan J. Perry²

1 National Suicide Research Foundation, Cork, Ireland, **2** School of Public Health, University College Cork, Cork, Ireland, **3** Department of Psychiatry, University College Cork, Cork, Ireland, **4** Liaison Psychiatry Service, Cork University Hospital, Cork, Ireland

* earensman@ucc.ie



Abstract

Objective

The specific objectives of this study were to examine variation in the care of self-harm patients in hospital settings and to identify the factors that predict recommended next care following self-harm.

Methods

Data on consecutive presentations to Irish emergency departments (EDs) involving self-harm from the National Self-Harm Registry Ireland from 2004 to 2012 were utilised. Univariate and multivariate regression analyses were performed to assess the associations between patients' clinical and demographic characteristics, and recommended next care received.

Results

Across the study period a total 101,904 self-harm presentations were made to hospital EDs, involving 63,457 individuals. Over the course of the study there was a declining number of presentations resulting in patient admission following attendance with self-harm. Recommended next care varied according to hospital location, with general admission rates ranging from 11% to 61% across administrative health regions. Multinomial logistic regression identified that the factor which most strongly affected next care was the presenting hospital. Being male, older age, method, repeat self-harm, time of attendance and residence of the patient were all identified as influencing care received. Psychiatric admission was most common when highly lethal methods of self-harm were used (OR = 4.00, 95% CI, 3.63–4.41). A relatively large proportion of patients left the ED without being seen (15%) and the risk of doing so was highest for self-harm repeaters (1.64, 1.55–1.74 for those with 5+ presentations).

OPEN ACCESS

Citation: Arensman E, Griffin E, Daly C, Corcoran P, Cassidy E, Perry IJ (2018) Recommended next care following hospital-treated self-harm: Patterns and trends over time. PLoS ONE 13(3): e0193587. <https://doi.org/10.1371/journal.pone.0193587>

Editor: Keith M. Harris, University of Queensland, AUSTRALIA

Received: October 3, 2017

Accepted: February 14, 2018

Published: March 1, 2018

Copyright: © 2018 Arensman et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data used in this manuscript comes from the National Self-Harm Registry Ireland and contains sensitive patient information. As a result, access to this data set is restricted and facilitated by the National Suicide Research Foundation. All data requests may be made to the National Suicide Research Foundation (info@nsrf.ie).

Funding: The National Self-Harm Registry Ireland is funded by the Irish Health Service Executive's National Office for Suicide Prevention.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

The extensive hospital variation in recommended next care indicates that management of self-harm patients may be determined more by where they present than by the needs of the patient. The study outcomes underline the need to standardise the clinical management of self-harm patients in general hospital settings.

Introduction

Presentations of self-harm represent a substantial number of all hospital emergency department (ED) visits.[1–3] International guidelines for the short-term management of self-harm in emergency settings support and advocate the standardised assessment and management of self-harm.[4,5] However, previous studies have found regional variation in the admission rates and assessment procedures following hospital-treated self-harm, with rates of psychosocial assessment ranging from 36–82%.[6–8] Such variation suggests that care frequently appears to be below the standard recommended.[9–11] Clinical management and recommended next care of self-harm has also been shown to vary both by patient characteristics such as method of self-harm,[12] as well as by the presenting hospital.[13] Furthermore, there is limited evidence to suggest that clinical management of self-harm is associated with improved outcomes for self-harm patients.[14]

The focus of previous studies on this topic have tended to be narrow—for example, focusing on hospital admission as a sole outcome [15–17] or specific methods of self-harm.[12] Longitudinal data obtained by the Irish National Self-Harm Registry provides the opportunity to examine patterns of recommended next care for hospital-treated self-harm. The specific study objectives were to examine variation in the management of self-harm patients based on standard demographic and clinical characteristics; to map regional variation in recommended next following self-harm at national level and to examine trends over time; and to identify the factors that predict recommended next care.

Method

National Self-Harm Registry Ireland

Data for the period 2004 to 2012 were obtained from the National Self-Harm Registry Ireland. Between 2004 and 2005 the Registry had near-complete coverage of the hospital EDs in Ireland, with all hospitals contributing data to the Registry since 2006. A weighting was applied to adjust for the lack of data from two hospitals in 2004 and 2005.

Definition of self-harm

The Registry uses the following definition of self-harm: “an act with non-fatal outcome in which an individual deliberately initiates a non-habitual behaviour, that without intervention from others will cause self harm, or deliberately ingests a substance in excess of the prescribed or generally recognised dosage, and which is aimed at realising changes that a person desires via the actual or expected physical consequences”.[18] This definition was developed by the WHO/Euro Multicentre Study and is consistent with that used in the English multicentre study.[2] A repeat episode is defined as re-presentation to any ED due to self-harm after an index episode of self-harm.

Self-harm data is collected and collated by independent Data Registration Officers according to standardised operating procedures.³

Data items

The Registry has a core dataset including: gender, age, date and hour of attendance at hospital and method(s) of self-harm according to the tenth revision of the WHO's International Classification of Disease codes for intentional injury (X60-X84). In addition, patient initials (in an encrypted format) and area of residence, coded to administrative area, are recorded.

Recommended next care following treatment to the presenting ED is routinely recorded by the Registry. The outcomes that were captured included: a) admission to a general ward in the presenting hospital; b) admission to a psychiatric ward in the presenting hospital; c) patient refusing admission/ leaving without being seen; and d) discharge from ED following treatment.

Whether or not alcohol was consumed as part of the self-harm act was ascertained through hospital case notes—if it was recorded on registration by the attending clinician or if present on toxicology reports.

Data analysis

Univariate analyses were performed using chi-square tests with a significance level set at $p < 0.01$. A multinomial logistic regression model was developed to assess the associations between patients' clinical and demographic characteristics, and recommended next care received in order to identify which factors differentiated between general admission, psychiatric admission, leaving without being seen/refusing admission, and discharged from the ED. The dependent variable comparison group represented those patients discharged from the ED. Odds Ratios and their 95% Confidence Intervals were reported with the associated level of significance. All statistics were performed using SPSS Version 21.

Mapping of regional variations in recommended next care

The map (Fig 1) details the recommended next care following hospital-treated self-harm, according to the Health Service Executive (HSE) Hospital Groups in Ireland. The map was developed using Health Atlas (www.healthatlasireland.ie).

Ethical approval

Ethical approval has been granted by the National Research Ethics Committee of the Faculty of Public Health Medicine, Ireland and relevant hospital and regional ethics committees. The National Suicide Research Foundation is registered with the Irish Data Protection Agency and complies with the Irish Data Protection Act of 1988 and the Irish Data Protection (Amendment) Act of 2003. Only anonymised data are released in aggregate form in reports. The names and addresses of patients are not recorded.

Results

During the nine-year period 1st January 2004 to 31st December 2012, there were 101,904 self-harm presentations to hospital recorded by the Registry, involving 63,457 individuals. Intentional drug overdose was the most common method of self-harm, present in 72% of all episodes. Self-cutting was the only other common method, involved in 22% of all presentations.

Table 1 shows the patterns of recommended next care following self-harm, based on demographic, temporal and clinical characteristics. Most commonly, 41% of presentations were




Fig 1. Regional variation of recommended next care of hospital-treated self-harm by HSE Hospital Group, average 2004–2012.

<https://doi.org/10.1371/journal.pone.0193587.g001>

Table 1. Demographic and clinical characteristics according to recommended next care received.

	Discharge from ED (n = 42,047; 41.3%)		General admission (n = 33,348; 32.7%)		Psychiatric admission (n = 11,592; 11.4%)		Left without being seen/ refuse admission (n = 14,917; 14.6%)		X ² (df)
	n	(%)	n	(%)	n	(%)	n	(%)	
Gender									
Male	19010	(41.0%)	14103	(30.4%)	5616	(12.1%)	7637	(16.5%)	
Female	23037	(41.5%)	19245	(34.7%)	5976	(10.8%)	7280	(13.1%)	375.77(3), P<0.001
Age (years)									
<15	759	(37.5%)	1132	(55.9%)	54	(2.7%)	79	(3.9%)	
15–24	14845	(47.8%)	9283	(29.9%)	2696	(8.7%)	4247	(13.7%)	
25–34	10618	(41.4%)	7623	(29.7%)	3252	(12.7%)	4132	(16.1%)	
35–44	8263	(37.8%)	7133	(32.6%)	2785	(12.7%)	3666	(16.8%)	
45–54	4993	(36.0%)	5015	(36.2%)	1765	(12.7%)	2098	(15.1%)	
55+	2569	(34.4%)	3162	(42.4%)	1040	(13.9%)	695	(9.3%)	367.81(1), P<0.001
Presentation									
1 st	26929	(42.4%)	22114	(34.8%)	6103	(9.6%)	8311	(13.1%)	
2 nd	5780	(39.2%)	4942	(33.5%)	1874	(12.7%)	2159	(14.6%)	
3 rd	2549	(38.6%)	2037	(30.9%)	956	(14.5%)	1054	(16.0%)	
4 th	1459	(38.5%)	1121	(29.6%)	567	(15.0%)	640	(16.9%)	
5 th or subsequent	5330	(40.0%)	3134	(23.5%)	2092	(15.7%)	2753	(20.7%)	186.12(1), P<0.001
Method									
Drug overdose	25028	(37.4%)	26855	(40.1%)	5613	(8.4%)	9402	(14.1%)	
Self-cutting	9608	(56.8%)	200	(11.8%)	2321	(13.7%)	2978	(17.6%)	
Drug overdose and self-cutting	1854	(42.2%)	1287	(29.3%)	556	(12.6%)	700	(15.9%)	
Attempted hanging	1282	(39.0%)	639	(19.4%)	995	(30.3%)	370	(11.3%)	
Attempted drowning	833	(38.8%)	407	(18.9%)	550	(25.6%)	359	(16.7%)	
Other	3442	(41.6%)	2160	(26.1%)	1557	(18.8%)	1108	(13.4%)	7644.56(15), P<0.001
Alcohol involvement									
Yes	16121	(39.1%)	14426	(35.0%)	3578	(8.7%)	7131	(17.3%)	
No	25926	(42.7%)	18922	(31.2%)	8014	(13.2%)	7786	(12.8%)	963.57(3), P<0.001
Residence									
Household resident	38437	(41.2%)	31356	(33.6%)	9993	(10.7%)	9993	(14.4%)	
Hospital in-patient	205	(16.5%)	268	(21.6%)	732	(58.9%)	38	(3.1%)	
Prisoner	233	(68.7%)	76	(22.4%)	6	(1.8%)	6	(7.1%)	
Homeless	1360	(44.7%)	490	(16.1%)	396	(13.0%)	396	(26.2%)	
Other	847	(45.4%)	515	(27.6%)	217	(11.6%)	217	(15.3%)	3562.93(12), P<0.001
Year of presentation									
2004	3117	(30.4%)	4134	(40.3%)	1497	(14.6%)	1521	(14.8%)	
2008	5131	(44.0%)	3836	(32.9%)	1185	(10.2%)	1517	(13.0%)	
2012	5723	(47.7%)	3311	(27.6%)	1241	(10.3%)	1735	(14.4%)	46.14(1), P<0.001
Time of presentation									
12am-3am	9719	(42.1%)	7254	(31.5%)	2075	(9.0%)	4017	(17.4%)	
4am-7am	4652	(42.9%)	3588	(33.1%)	986	(9.1%)	1621	(14.9%)	
8am-11am	3757	(43.5%)	3028	(35.1%)	1086	(12.6%)	757	(8.8%)	
12pm-3pm	6561	(42.3%)	4944	(31.9%)	2191	(14.1%)	1800	(11.6%)	
4pm-7pm	7686	(39.2%)	6492	(33.1%)	2560	(13.1%)	2851	(14.6%)	
8pm-11pm	9453	(40.2%)	7658	(32.5%)	2599	(11.0%)	3819	(16.2%)	19.17(1), P<0.001
Weekend presentation									
Yes	12909	(41.4%)	10216	(32.7%)	3444	(11.0%)	4634	(14.9%)	

(Continued)

Table 1. (Continued)

	Discharge from ED (n = 42,047; 41.3%)		General admission (n = 33,348; 32.7%)		Psychiatric admission (n = 11,592; 11.4%)		Left without being seen/ refuse admission (n = 14,917; 14.6%)		X ² (df)
	n	(%)	n	(%)	n	(%)	n	(%)	
No	29138	(41.2%)	23132	(32.7%)	8148	(11.5%)	10283	(14.5%)	6.04(3), P = .110
City resident									
Yes	14404	(46.9%)	6202	(20.2%)	3822	(12.4%)	6316	(20.5%)	
No	27643	(38.8%)	27146	(38.1%)	7770	(10.9%)	8601	(12.1%)	3545.14(3), P<0.001

Note: Presentation variable indicates whether it was a patient’s first, second or subsequent attendance with self-harm. ‘Other’ residence would include non-residents, such as tourists and visitors. Data presented for year of presentation includes first, middle and last years of study period only.

<https://doi.org/10.1371/journal.pone.0193587.t001>

discharged from the ED following treatment. A similar proportion (44%) resulted in admission to either a general ward (33%) or a psychiatric unit (11%). In 15% of presentations, the patient either refused to be admitted or left the ED before a next care recommendation could be made. Patterns of recommended next care varied significantly across the study period [X^2 for trend (1) = 46.14; $p < 0.001$], with general ward admission decreasing by 30% (from 40% to 28%) (see also [S1 Fig](#)).

Recommended next care varied according to gender ($X^2(3) = 375.77$; $p < 0.001$). Overall, females were more likely to be admitted to a general ward than males (34.7% vs. 30.4%). Males more often left the ED before a next care recommendation could be made or they refused treatment (16.5% vs. 13.1%). Recommended next care also varied significantly by age [X^2 for trend (1) = 367.81; $p < 0.001$], with the rate of general admission to a hospital ward being highest for children and adolescents under 15 years (56%) and for those aged over 55 years (42%). Almost half of those aged between 15–24 years were discharged from the ED following treatment. In addition, psychiatric admission following a self-harm presentation was lowest among those aged under 15 years (2.7%).

Next care also varied according to method of self-harm [$X^2(15) = 7644.56$, $p < 0.001$]. General inpatient admission was most common following cases of poisoning, drug overdose and in cases where alcohol was involved (40%, 39% and 35%, respectively), and least common following self-cutting (16%). Psychiatric admission was most common following cases of attempted hanging and attempted drowning (30% and 25%, respectively). Patients were most likely to leave the ED without being seen if alcohol was involved (17.3% vs. 12.8%) or if the episode of self-harm involved self-cutting only (16.7%).

Admission was significantly associated with time of presentation, with variation in both general (32.5% to 39.8%) and psychiatric (9.0% to 14.1%) admission rates over the course of the day, and in the proportion of patients leaving without being seen/ refusing admission (8.8% to 17.4%).

Regional variation of recommended next care

Recommended next care varied by the Health Service Executive Hospital Groups where the presenting hospital was located ([Fig 1](#)). General admission ranged from 11.2% in the Dublin North East Hospital Group to 61.0% in the South Eastern Hospital Group. Admission to a psychiatric ward was lowest (3.7%) in the North Eastern Hospital Group and highest in the Dublin South Hospital Group (19.3%). The Dublin North East Hospital Group had the highest proportion of patients leaving the ED without being seen or without a clinical decision being made (18.9%).

Factors associated with next care following hospital-treated self-harm

Table 2 details the results of the multivariate multinomial regression which was conducted to identify the factors independently associated with self-harm patients and different types of recommended next care. The analysis controlled for the hospital where the presentation was

Table 2. Multinomial logistic regression analysis of the independent associations of factors and recommended next care following hospital-treated self-harm.

Variable	General admission		Psychiatric admission		Left without being seen/ refuse admission	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Gender (female)						
Male	1.10*	(1.06–1.13)	1.15*	(1.10–1.21)	1.30*	(1.25–1.35)
Age in years (55+)						
<15	1.11	(.97–1.27)	0.18*	(0.13–0.26)	0.48*	(0.36–0.64)
15–24	0.45*	(0.42–0.49)	0.43*	(0.39–0.47)	1.01	(0.92–1.11)
25–34	0.57*	(0.53–0.61)	0.71*	(0.65–0.78)	1.28*	(1.16–1.40)
35–44	0.66*	(0.61–0.70)	0.80*	(0.73–0.87)	1.39*	(1.26–1.52)
45–54	0.80*	(0.74–0.86)	0.88*	(0.80–0.97)	1.38*	(1.25–1.53)
Presentation (1st)						
2 nd	1.13*	(1.07–1.19)	1.44*	(1.35–1.54)	1.20*	(1.31–1.27)
3 rd	1.12*	(1.05–1.21)	1.63*	(1.50–1.78)	1.32*	(1.22–1.43)
4 th	1.11	(1.01–1.22)	1.67*	(1.50–1.87)	1.38*	(1.25–1.53)
5 th or subsequent	0.96	(0.91–1.02)	1.64*	(1.53–1.75)	1.64*	(1.55–1.74)
Method (drug overdose)						
Self-cutting	0.16*	(0.15–0.17)	0.99	(0.93–1.05)	0.77*	(0.73–0.81)
Drug overdose and self-cutting						
Attempted hanging	0.70*	(0.64–0.76)	1.45*	(1.30–1.61)	0.92	(0.83–1.01)
Attempted drowning	0.45*	(0.40–0.50)	4.00*	(3.63–4.41)	0.75*	(0.66–0.85)
Other	0.36*	(0.32–0.41)	2.95*	(2.61–3.33)	0.93	(0.81–1.06)
Alcohol involvement (No)						
Yes	0.95	(0.92–0.99)	0.68*	(0.65–0.71)	1.24*	(1.19–1.29)
Residence (Household resident)						
Hospital in-patient	1.75*	(1.42–2.15)	9.61*	(8.11–11.38)	0.47*	(0.33–0.66)
Prisoner	0.39*	(0.29–0.52)	0.05*	(0.22–0.12)	0.25*	(0.16–0.39)
Homeless	0.67*	(0.60–0.76)	0.80*	(0.71–0.90)	1.14	(1.04–1.26)
Other	0.69*	(0.60–0.78)	0.65*	(0.55–0.76)	0.89	(0.77–1.02)
Year of presentation (2004)						
2008	0.55*	(0.52–0.60)	0.41*	(0.37–0.45)	0.55*	(0.50–0.60)
2012	0.44*	(0.41–0.47)	0.38*	(0.34–0.41)	0.55*	(0.51–0.60)
Time of presentation (8pm–11pm)						
12am–3am	0.95	(0.90–0.99)	0.85*	(0.79–0.91)	1.00	(0.95–1.06)
4am–7am	0.97	(0.92–1.04)	0.87	(0.80–0.95)	0.85*	(0.79–0.91)
8am–11am	1.10	(1.03–1.17)	1.11	(1.02–1.21)	0.54*	(0.50–0.59)
12pm–3pm	0.94	(0.89–1.00)	1.20*	(1.11–1.28)	0.71*	(0.67–0.76)
4pm–7pm	1.03	(0.98–1.09)	1.18*	(1.11–1.27)	0.94	(0.89–1.00)
Weekend presentation (Yes)						
No	0.97	(0.94–1.01)	1.00	(0.95–1.05)	0.98	(0.93–1.02)
City resident (No)						
Yes	0.90	(0.85–0.94)	1.00	(0.94–1.07)	1.28*	(1.21–1.35)

* = p<0.001. Note: Reference category for dependent variable is 'discharge from ED'. Reference categories for independent variables shown in brackets. All factors listed were entered into the multivariate model.

made, and all 38 hospitals were included in the regression model. This variable was most strongly associated with recommended next care, with Odds Ratios (ORs) ranging from 0.04 to 2.38 (see S2–S4 Figs).

Gender was independently associated with recommended next care, with males more likely to be admitted to both general (OR = 1.10, 95% CI, 1.06–1.13) and psychiatric wards (1.15, 1.10 to 1.21) than females. In addition, males were more likely to leave the ED without being seen or refuse admission (1.30, 1.25–1.35). The likelihood of being admitted to a psychiatric ward increased with increasing age, and was lowest for those aged under 15 years (0.18, 0.13–0.26). A similar trend was observed for general admission (0.45, 0.42–0.49 for 15–24 year-olds), however those aged under 15 years were just as likely as older adults (55+) to be admitted to a general ward. Young and middle-aged adults (25–54 years) were most likely to leave the ED without being seen or to refuse admission, with those aged under 15 least likely to do so (0.48, 0.36–0.64). Patients residing in psychiatric hospitals were most likely to be both admitted to general (1.75, 1.42–2.15) and psychiatric (9.61, 8.11–11.38) wards. City residents were more likely to leave the ED without being seen (1.28, 1.21–1.35).

General and psychiatric admission following self-harm decreased over the study period, as did the proportion of those leaving without being seen/refusing admission. Presentations made between 12pm and 7pm (12pm–3pm: 1.20, 1.11–1.28; 4pm–7pm: 1.18, 1.11–1.27, respectively) were most likely to receive psychiatric admission, while those attending between 4am and 3pm were least likely to leave without being seen by a doctor. Whether the attendance occurred during the week or at the weekend did not significantly influence next care recommendations.

Characteristics of the self-harm presentation also influenced recommended next care. A person was most likely to be admitted to a general ward following intentional drug overdose, and least likely if self-cutting was involved (0.16, 0.15–0.17). Attempted hanging was most strongly associated with psychiatric admission (4.00, 3.63–4.41), while those engaging in self-cutting were least likely to leave without being seen (0.77, 0.73–0.81). Having used/misused alcohol was associated with leaving without being seen (1.24, 1.19–1.29). The number of previous presentations a person had made also influenced next care. The likelihood of being admitted to a psychiatric ward increased with each subsequent attendance for self-harm (5th or subsequent attendance: 1.64, 1.53–1.75). For admission to a general ward, only those patients on their second or third attendance were more likely to be admitted (1.13, 1.07–1.19; 1.12, 1.05–1.21, respectively). In contrast, patients on a repeat presentation were also more likely to leave the ED without being seen or to refuse admission (5th or subsequent attendance: 1.64, 1.55–1.74).

Discussion

This is the first study addressing recommended aftercare following self-harm presenting to general hospital at national level. The findings of this study indicate that over time there was a decreasing number of presentations resulting in inpatient admission to the presenting hospital following self-harm. In addition, significant variation in the patterns of recommended next care was observed across hospitals. The factor most strongly associated with recommended next care was the hospital to which the individual presented. The study also identified that being male, older age, method of self-harm, repeat attendance, time of attendance and residence of the patient all influenced the type of care patients received. Males were more likely to be admitted to hospital, and also more likely to leave without being seen, than females. Older patients were more likely to be admitted to hospital following self-harm, with those aged between 25–44 years most likely to leave without being seen. Psychiatric admission following self-harm was most common when the presentation was made during daytime hours.

This study identified a sub-group of patients who were at greatest risk of leaving the ED without being seen, or refusing admission. These patients were most likely to be male and aged between 25–44 years. These presentations were more likely to involve an intentional drug overdose or attempted drowning, often involving alcohol. City residents, those with no fixed abode, and those with a history of previous self-harm, were more likely to leave without treatment. Patients presenting due to self-harm in the evening or at night time were also more likely to leave without treatment.

Our study has a number of strengths and limitations. Firstly, Ireland's National Self-Harm Registry is the world's first national registry focusing specifically on non-fatal self-harm presenting to hospital EDs and provides a unique opportunity to establish, at a national level, the patterns of recommended next care in hospitals involving non-fatal suicidal behavior.[19] Secondly, over 100,000 presentations of self-harm made over a nine-year period were included in this analysis, providing robust data to examine longitudinal patterns of care for self-harm patients. A limitation of the study is the lack of information on psychosocial assessments offered to patients during their ED presentation, as well as on referrals provided to patients who were discharged from the presenting hospital. This information was not included in the data collection for the Registry during the study period. More detailed clinical data relating to psychiatric history, degree of suicidal intent, severity of the self-harm act and contributory factors were not recorded. To this end, we were not able to quantify their impact on recommended next care and acknowledge that these would be important to consider. Additionally, it may not always be recorded in the ED that a patient has been admitted directly to psychiatric inpatient care, or a patient may be admitted to an external psychiatric hospital for admission. Therefore, the figures presented for psychiatric admissions could be underestimated. Finally, establishing self-harm history was confined to presentations made that were recorded by the Registry database, thus assuming that a person's first episode in the Registry was their first presentation to hospital for self-harm.

To our knowledge only one other recent study has examined patterns of recommended next care for self-harm using a multinomial approach.[16] However, our research is based on national data and covers a longer time period. The variation in recommended next care across hospitals was similar to other studies.[6–8] However, general ward admissions were higher than those observed in a Spanish study [16] and lower than those reported in England.[6–8] and the improvement in admission rates in England since 2001 (42% to 54%)[8] was not reflected in our data. The independent risk factors for recommended next care following self-harm were consistent with previous literature.[17] In addition, the present study identified specific factors that were associated with increased likelihood of leaving the ED without being seen, which has not been examined previously. The observed variation in recommended next care across the study period and by hospital is undoubtedly complex and many contributing factors exist. The reduction in hospital admissions was most pronounced between 2008 and 2012, where numbers of general admission rates decreased by 16%. During this period of global economic recession, self-harm presentations increased by 31% for males and 22% for females.[20,21] The austerity measures introduced to Ireland during this time period involved substantial cuts in health care spending which may have impacted negatively on hospital resources, such as the availability of hospital beds and qualified staff to conduct psychosocial and psychiatric assessments. It is possible that the reduction in admissions during the study period was related to patients more often being treated in outpatient and community-based settings. We did not have access to data regarding referrals for discharged patients to examine this. However, given that the proportion of presentations resulting in the patient leaving the ED without being seen did not improve during the study period, it is unlikely that a reduction in admissions is fully attributable to service improvements.

The variation in recommended next care by hospital highlighted by the present study, poses a significant challenge for the assessment and management of self-harm in ED settings. The lowest admission rates following self-harm were observed in urbanised areas, coupled with a large proportion of patients leaving without being seen. In recent years, concerns have been expressed in the national media in relation to ED overcrowding in urban areas and to low per capita level of inpatient beds.[22] The recent Irish National Clinical Programme for the Management of Self Harm Presentations to Emergency Departments [23] aims to develop a standardised and effective process for the assessment and management of self-harm in an ED setting, and a specific aim of the programme to reduce the number of people who leave before assessment and reduce the number of repeat presentations. In terms of capacity building, new clinical posts have been allocated to each ED based on the data from the Registry to implement the programme and this resource will have a significant impact on availability of specialist staff on-site. A key to success of this programme will be effective implementation along with staff training as it is worth noting that in England, despite the comprehensive National Institute for Clinical Excellence (NICE) guidelines,[4] the proportion of patients who received a psychosocial assessment prior to leaving the ED did not significantly change.⁸

The present study highlights the variation in recommended next care for patients of self-harm according to previous self-harm history. Those who engaged in their second or third presentation were most likely to be admitted to a general ward, and those on their fourth or subsequent visit most likely to be admitted to a psychiatric ward. However, infrequent (less than five presentations) and frequent (five or more presentations) were all more likely to leave the ED without being seen, or to refuse admission, than those presenting on their index episode. This raises questions about the delivery and quality of initial assessments and current treatment for self-harm repeaters as well as the lack of awareness of the importance of self-harm and suicide risk assessment among these patients. Therefore, it is argued here that there is need for uniform assessment and referral procedures, in line with international best practice [4,5,24] to ensure the most appropriate treatment. For example, for patients with a pattern of frequent repetition of self-harm, Dialectical Behaviour Therapy is an indicated treatment.[25] However, if many of these patients do not receive an assessment at all, the likelihood of these patients receiving the required treatment is low.

In line with previous research, the hospital to which patients present appears to significantly influence the type of care and treatment a self-harm patient receives. In our study, data on hospital characteristics were lacking. Future research could use multi-level modelling approaches to explore specific hospital characteristics which affect the next care of self-harm. This research has identified that while repeaters of self-harm are more likely to be admitted to the presenting hospital, this occurs up to a point. Future research needs to explore the challenges associated with reaching this group of patients, and to ensure that the most appropriate treatment is being provided. There is varying evidence for the effectiveness of psychosocial assessments at the time of presentation in reducing repetition of self-harm.[14] Failure to routinely conduct a thorough and standardised psychosocial assessment of all self-harm patients may represent a missed opportunity to reduce the risk of both non-fatal and fatal repetition.[26] The Registry also has the potential to establish the effectiveness of out-patient referrals on repetition of self-harm, and future research should address this.

The results of this study show that non-clinical factors of self-harm presentations strongly influence patterns of hospital admission following self-harm. In particular, the hospital variation in recommended next care poses a challenge for the assessment and management of self-harm patients, and underlines the need for uniform guidelines.

Supporting information

S1 Fig. Changes in recommended next care of hospital-treated self-harm in Ireland, 2004–2012.

(TIF)

S2 Fig. Adjusted odds ratios by hospital for general admission.

(TIF)

S3 Fig. Adjusted odds ratios by hospital for psychiatric admission.

(TIF)

S4 Fig. Adjusted odds ratios for by hospital left without being seen/ refuse admission.

(TIF)

Acknowledgments

We would like to acknowledge the work of the Data Registration Officers in compiling data for the National Self-Harm Registry Ireland. We would also like to thank Eileen Williamson and Paul Williamson in their contributions to the design of this paper.

Author Contributions

Conceptualization: Ella Arensman, Eve Griffin, Paul Corcoran, Eugene Cassidy, Ivan J. Perry.

Data curation: Eve Griffin, Paul Corcoran.

Formal analysis: Eve Griffin, Paul Corcoran.

Investigation: Ella Arensman, Eve Griffin.

Methodology: Eve Griffin, Paul Corcoran, Ivan J. Perry.

Project administration: Eve Griffin.

Supervision: Paul Corcoran, Ivan J. Perry.

Visualization: Eve Griffin.

Writing – original draft: Ella Arensman, Eve Griffin, Caroline Daly.

Writing – review & editing: Ella Arensman, Eve Griffin, Caroline Daly, Paul Corcoran, Eugene Cassidy, Ivan J. Perry.

References

1. Claassen CA, Trivedi MH, Shimizu I, Stewart S, Larkin GL, Litovitz T. Epidemiology of nonfatal deliberate self-harm in the United States as described in three medical databases. *Suicide Life Threat Behav* 2006; 36: 192–212 PMID: [16704324](https://pubmed.ncbi.nlm.nih.gov/16704324/)
2. Hawton K, Bergen H, Casey D, Simkin S, Palmer B, Cooper J, et al. Self-harm in England: a tale of three cities. *Soc Psychiatry Psychiatr Epidemiol* 2007; 42: 513–521. <https://doi.org/10.1007/s00127-007-0199-7> PMID: [17516016](https://pubmed.ncbi.nlm.nih.gov/17516016/)
3. Perry IJ, Corcoran P, Fitzgerald AP, Keeley HS, Reulbach U, Arensman E. The incidence and repetition of hospital-treated deliberate self harm: findings from the world's first national registry. *PLoS One* 2012; 7: e31663. <https://doi.org/10.1371/journal.pone.0031663> PMID: [22363700](https://pubmed.ncbi.nlm.nih.gov/22363700/)
4. National Institute for Health and Care Excellence. Self-harm: The short-term physical and psychological management and secondary prevention of self-harm in primary and secondary care. London: National Institute for Health and Care Excellence, 2004.

5. Knesper DJ. Continuity of care for suicide prevention and research: Suicide attempts and suicide deaths subsequent to discharge from the emergency department or psychiatry inpatient unit. Newton, MA: Education Development Center, 2010.
6. Bennewith O, Gunnell D, Peters T, Hawton K, House A. Variations in the hospital management of self harm in adults in England: observational study. *BMJ* 2004; 328(7448): 1108–1109. <https://doi.org/10.1136/bmj.328.7448.1108> PMID: 15130979
7. Bergen H, Hawton K, Waters K, Cooper J, Kapur N. Psychosocial assessment and repetition of self-harm: The significance of single and multiple repeat episode analyses. *J Affect Disord* 2010; 127: 257–265. <https://doi.org/10.1016/j.jad.2010.05.001> PMID: 20684848
8. Cooper J, Steeg S, Bennewith O, Lowe M, Gunnell D, House A, et al. Are hospital services for self-harm getting better? An observational study examining management, service provision and temporal trends in England. *BMJ Open* 2013; 3: e003444. <https://doi.org/10.1136/bmjopen-2013-003444> PMID: 24253029
9. Cooper J, Murphy E, Jordan R, Mackway-Jones K. Communication between secondary and primary care following self-harm: are National Institute of Clinical Excellence (NICE) guidelines being met? *Ann Gen Psychiatry* 2008; 7: 21. <https://doi.org/10.1186/1744-859X-7-21> PMID: 18947385
10. Baraff LJ, Janowicz N, Asarnow JR. Survey of California emergency departments about practices for management of suicidal patients and resources available for their care. *Ann Emerg Med* 2006; 48: 452–458. <https://doi.org/10.1016/j.annemergmed.2006.06.026> PMID: 16997683
11. Kapur N, Steeg S, Webb R, Haigh M, Bergen H, Hawton K, et al. Does clinical management improve outcomes following self-harm? Results from the multicentre study of self-harm in England. *PloS One* 2013; 8: e70434. <https://doi.org/10.1371/journal.pone.0070434> PMID: 23936430
12. Lilley R, Owens D, Horrocks J, House A, Noble R, Bergen H, et al. Hospital care and repetition following self-harm: multicentre comparison of self-poisoning and self-injury. *Br J Psychiatry* 2008; 192: 440–445. <https://doi.org/10.1192/bjp.bp.107.043380> PMID: 18515895
13. Carroll R, Corcoran P, Griffin E, Perry IJ, Arensman E, Gunnell D, et al. Variation between hospitals in inpatient admission practices for self-harm patients and its impact on repeat presentation. *Soc Psychiatry Psychiatr Epidemiol* 2016; 51: 1485–1493. <https://doi.org/10.1007/s00127-016-1247-y> PMID: 27300340
14. Carroll R, Metcalfe C, Gunnell D. Hospital management of self-harm patients and risk of repetition: Systematic review and meta-analysis. *J Affect Disord* 2014; 168: 476–483 PMID: 25128754
15. Gunnell D, Hawton K, Ho D, Evans J, O'Connor S, Potokar J, et al. Hospital admissions for self harm after discharge from psychiatric inpatient care: Cohort study. *BMJ* 2008; 337: a2278. <https://doi.org/10.1136/bmj.a2278> PMID: 19018041
16. Jimenez-Trevino L, Saiz PA, Corcoran P, Burón P, García-Portilla MP, Chinea ER, et al. Factors associated with hospitalization after suicide spectrum behaviors: results from a multicentre study in Spain. *Arch Suicide Res* 2015; 19: 17–34. <https://doi.org/10.1080/13811118.2013.824841> PMID: 24810383
17. Suominen K, Lönnqvist J. Determinants of psychiatric hospitalization after attempted suicide. *Gen Hosp Psychiatry* 2006; 28: 424–430. <https://doi.org/10.1016/j.genhosppsych.2006.03.009> PMID: 16950379
18. Platt S, Bille-Brahe U, Kerkhof A, Schmidtke A, Bjerke T, Crepet P, et al. Parasuicide in Europe: the WHO/EURO multicentre study on parasuicide. I. Introduction and preliminary analysis for 1989. *Acta Psychiatr Scand* 1992; 85: 97–104 PMID: 1543046
19. World Health Organisation. Practice manual for establishing and maintaining surveillance systems for suicide attempts and self-harm. Geneva: World Health Organisation, 2016.
20. Corcoran P, Griffin E, Arensman E, Fitzgerald AP, Perry IJ. Impact of the economic recession and subsequent austerity on suicide and self-harm in Ireland: an interrupted time series analysis. *Int J Epidemiol* 2015; 44(3): 969–977. <https://doi.org/10.1093/ije/dyv058> PMID: 26082406
21. Thomas S, Burke S, Barry S. The Irish health-care system and austerity: sharing the pain. *Lancet* 2014; 383: 1545–1546. [https://doi.org/10.1016/S0140-6736\(14\)60744-3](https://doi.org/10.1016/S0140-6736(14)60744-3) PMID: 24792851
22. Sheehy C. Beaumont hospital declared 'unsafe' by hospital CEO. *Irish Independent*; 2014.
23. Health Service Executive. National clinical care programme for the assessment and management of patients presenting to emergency departments following self-harm. Dublin, Health Service Executive, 2014.
24. Mehlum L, Ramberg M. Continuity of care in the treatment of suicide attempters—current challenges. *Arc Suicide Res* 2010; 14: 105–108.
25. Hawton K, Witt KG, Salisbury TLT, Arensman E, Gunnell D, Hazell P, et al. Psychosocial interventions following self-harm in adults: a systematic review and meta-analysis. *Lancet Psychiatry* 2016; 3: 740–750. [https://doi.org/10.1016/S2215-0366\(16\)30070-0](https://doi.org/10.1016/S2215-0366(16)30070-0) PMID: 27422028
26. Kapur N, Murphy E, Cooper J, Bergen H, Hawton K, Simkin S, et al. Psychosocial assessment following self-harm: results from the Multi-Centre Monitoring of Self-Harm Project. *J Affect Disord* 2008; 106: 285–293 PMID: 17761308