


CORRECTION

Correction: Big data and analysis of risk factors for gallbladder disease in the young generation of Korea

Hyung Sun Kim, Seong Kyung Cho, Chang Soo Kim, Joon Seong Park

In the Results subsection of the Abstract, there is an error in the fourth sentence. The correct sentence is: The prevalence rate of GB stones was 1.9% (2,979/154,463) in the young age group.

There is an error in [Table 3](#). Please see the correct [Table 3](#) here.



OPEN ACCESS

Citation: Kim HS, Cho SK, Kim CS, Park JS (2019) Correction: Big data and analysis of risk factors for gallbladder disease in the young generation of Korea. PLoS ONE 14(3): e0213950. <https://doi.org/10.1371/journal.pone.0213950>

Published: March 13, 2019

Copyright: © 2019 Kim et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 3. Anthropometric indices and laboratory data of the young age group with abnormal GB finding (stones, sludge, polyps, and adenomyomatosis) and normal GB finding.

	Men (n = 37301)			Women (n = 22098)		
	Abnormal GB (n = 7631)	Normal GB (n = 29670)	p-value ^a	Abnormal GB (n = 3042)	Normal GB (n = 19056)	p-value ^a
Age, yr	33.5±3.9	32.9±4.2	<0.01	32.7±4.3	31.3±4.8	<0.01
Height, cm	174.8±5.4	174.7±5.6	0.41	162.0±4.9	162.0±5.1	0.85
Weight, kg	74.9±9.6	75.0±10.1	0.11	57.1±8.2	56.5±7.8	<0.01
BMI	24.5±2.8	24.6±2.9	0.03	21.7±2.9	21.5±2.8	<0.01
Waist circumference, cm	82.8±7.1	82.9±7.3	0.08	71.3±7.0	70.7±6.7	<0.01
Thigh circumference, cm	53.9±4.2	53.9±4.4	0.83	50.4±4.4	50.2±4.3	0.01
Hb [12–16g/dl]	15.4±0.8	15.5±0.8	<0.01	13.1±0.8	13.0±13.0	<0.01
Hct [37–47%]	44.8±2.2	44.9±2.2	<0.01	38.9±2.2	39.1±2.2	<0.01
AST [16–37U/L]	22.3±7.3	23.3±7.9	<0.01	17.9±5.1	18.1±5.1	0.14
ALT [11–46U/L]	26.7±15.4	28.6±17.3	<0.01	14.7±8.3	14.3±7.9	0.01
Bilirubin [0.3–1.8mg/dl]	0.9±0.3	0.9±0.3	0.56	0.8±0.2	0.8±0.3	0.15
Albumin [3.4–5.3g/dl]	4.4±0.2	4.5±0.2	<0.01	4.3±0.2	4.4±0.2	<0.01
r-GTP [8–46U/L]	39.0±27.8	40.3±28.3	<0.01	18.3±12.0	18.1±11.4	0.24
ALP [35–83U/L]	53.9±11.8	54.4±11.7	<0.01	43.6±10.4	43.7±9.9	0.72
Cholesterol [<200mg/dl]	189.6±29.9	190.7±30.3	<0.01	178.5±27.3	177.6±26.8	0.08
HDL [40–99mg/dl]	54.7±12.3	54.9±12.4	0.07	67.7±13.7	68.4±13.7	0.02
LDL [<100mg/dl]	110.3±27.1	110.3±27.5	0.94	96.3±24.3	94.7±21.2	<0.01
Triglyceride [<150mg/dl]	124.1±70.8	128.2±72.8	<0.01	72.8±39.3	72.5±38.6	0.65

Abbreviations: GB, gallbladder; yr, years; BMI, body mass index; Hb, hemoglobin; Hct, hematocrit; AST, aspartate aminotransferase; ALT, alanine aminotransferase; r-GTP, gamma-glutamyl transpeptidase; ALP, alkaline phosphatase; HDL, high-density lipoprotein; LDL, low-density lipoprotein

^a p values were calculated using the t test.

<https://doi.org/10.1371/journal.pone.0213950.t001>

Reference

1. Kim HS, Cho SK, Kim CS, Park JS (2019) Big data and analysis of risk factors for gallbladder disease in the young generation of Korea. PLoS ONE 14(2): e0211480. <https://doi.org/10.1371/journal.pone.0211480> PMID: 30794560