

RESEARCH ARTICLE

Open Access

# The genetic association study between polymorphisms in uncoupling protein 2 and uncoupling protein 3 and metabolic data in dogs

Chihiro Udagawa<sup>1</sup>, Naomi Tada<sup>1</sup>, Junzo Asano<sup>1</sup>, Katsumi Ishioka<sup>2</sup>, Kazuhiko Ochiai<sup>1</sup>, Makoto Bonkobara<sup>3</sup>, Shuichi Tsuchida<sup>4</sup> and Toshinori Omi<sup>1\*</sup>

## Abstract

**Background:** The uncoupling proteins (UCPs) in the mitochondrial inner membrane are members of the mitochondrial anion carrier protein family that play an important role in energy homeostasis. Genetic association studies have shown that human *UCP2* and *UCP3* variants (SNPs and indels) are associated with obesity, insulin resistance, type 2 diabetes mellitus, and metabolic syndrome. The aim of this study was to examine the genetic association between polymorphisms in *UCP2* and *UCP3* and metabolic data in dogs.

**Results:** We identified 10 SNPs (9 intronic and 1 exonic) and 4 indels (intronic) in *UCP2*, and 13 SNPs (11 intronic and 2 exonic) and one indel (exonic) in *UCP3*, by DNA sequence analysis of 11 different dog breeds (n = 119). An association study between these *UCP2* and *UCP3* variants and the biochemical parameters of glucose, total cholesterol, lactate dehydrogenase and triglyceride in Labrador Retrievers (n = 50) showed that none of the *UCP2* polymorphisms were significantly associated with the levels of these parameters. However, four *UCP3* SNPs (intron 1) were significantly associated with total cholesterol levels. In addition, the allele frequencies of two of the four SNPs associated with higher total cholesterol levels in a breed that is susceptible to hypercholesterolemia (Shetland Sheepdogs, n = 30), compared with the control breed (Shiba, n = 30).

**Conclusion:** The results obtained from a limited number of individuals suggest that the *UCP3* gene in dogs may be associated with total cholesterol levels. The examination of larger sample sizes and further analysis will lead to increased precision of these results.

**Keywords:** Dog, SNP, Indel polymorphism, UCP2, UCP3, Association study

## Background

The uncoupling proteins (UCPs) in the mitochondrial inner membrane are members of the mitochondrial anion carrier protein family [1,2]. Mammals have five UCP homologs, of which UCP1, UCP2, and UCP3 are closely related, while UCP4 and UCP5 are more divergent from the other UCPs [3].

Based on genetic association studies, *UCP2*, *UCP3*, or both are reportedly associated with obesity, insulin resistance, type 2 diabetes mellitus, and metabolic syndrome in humans [4-11]. For example, a SNP in the 5'

untranslated region in human *UCP3*, the UCP3 -55CT SNP, is known to be a genetic marker associated with mRNA expression [12], elevated high density lipoprotein cholesterol levels, a reduced body mass index (BMI), weight, waist circumference, waist to hip ratio, fat mass, low density lipoprotein (LDL) cholesterol, and total cholesterol (T-Cho) [13-15].

The treatment and prevention of obesity and metabolic-related diseases are also clinically important in dogs [16-25]. Our previous report showed that the nucleotide sequences, predicted amino acid sequences and the genomic structures of human *UCP2* and *UCP3* are highly homologous to the canine orthologs [26,27]. In this study, we investigate whether the dog *UCP2* and *UCP3* genes are associated with alterations in metabolism.

\* Correspondence: t.omi@nvl.u.ac.jp

<sup>1</sup>Department of Basic Science, School of Veterinary Nursing and Technology, Faculty of Veterinary Science, Nippon Veterinary and Life Science University, 1-7-1 Kyonan-cho, Musashino, Tokyo 180-8602, Japan

Full list of author information is available at the end of the article

## Results and discussion


Figure 1 shows a schematic representation of the canine *UCP2* and *UCP3* genes and the identified DNA polymorphisms from 119 animals from 11 breeds. For analysis of the dog *UCP2* gene, six regions were individually amplified from genomic DNA and sequenced. We then identified 10 SNPs (9 intronic and 1 exonic) and 4 indels (intronic) in *UCP2* (Figure 1, Additional file 1). In the dog *UCP3* gene, 13 SNPs (11 intronic and 2 exonic) and 1 indel (exonic) were revealed by sequencing nine regions of this gene (Figure 1, Additional file 1).

To test the association between the dog *UCP2* and *UCP3* genes and metabolic data, we determined the genotype of 50 Labrador Retrievers for each of 14 polymorphic sites (10 SNPs and 4 indels) in the *UCP2* gene, and examined whether any of the genotypes were associated with biochemical measurements of glucose (GLU), total cholesterol (T-Cho), lactate dehydrogenase (LDH), or triglyceride (TG). To exclude any contamination by disease of the animals, we selected Labrador Retrievers that had undergone a health examination for breeding for guide dogs by the Kyushu Guide Dog Association.

The average of measurements was calculated with respect to the genotype group. Nine of the 14 loci in the *UCP2* gene were polymorphic in this population of Labrador Retrievers. None of these DNA polymorphisms in the *UCP2* gene were significantly associated with any of the biochemical parameters in this study (Additional file 2). We also subjected the 14 polymorphic sites (13 SNPs and 1 indel) in the *UCP3* gene to this association

analysis. Ten of the 14 sites were polymorphic in this population of Labrador Retrievers. There were no significant differences between genotype and GLU, LDH, or TG measurements for any polymorphic site. However, the T-Cho levels differed significantly among the genotype groups at four sites: -4399C/T, -4339T/C, -930T/C and -803C/T in intron 1 of the *UCP3* gene (*UCP3* intron1). The average T-Cho levels in dogs carrying CC or CT at -4399 C/T were  $273.5 \pm 49.0$  and  $237.2 \pm 53.3$ , respectively. The average T-Cho levels for the TT, TC, or CC genotypes at -4339T/C and -930T/C were  $264.3 \pm 49.6$ ,  $276.9 \pm 49.5$ , and  $233.5 \pm 51.2$ , respectively. Those for CC or CT at -803C/T were  $271.6 \pm 49.5$  and  $239.1 \pm 54.5$ , respectively (Table 1). The genotype distributions were in a Hardy-Weinberg equilibrium.

Shetland Sheepdogs are considered to have a predisposition to primary hyperlipidemia as determined by the levels of cholesterol, triglycerides, and free fatty acids [28,29]. Therefore, we investigated the distribution of genotypes for SNPs and indels of the *UCP2* and *UCP3* genes in a population of Shetland Sheepdogs (n = 30). Shiba (n = 30) were also tested as a comparative contrast breed in this study. Statistically significant differences in allele frequency between the two breeds were found in five of the 14 polymorphic sites in *UCP2* (-3629C/G, -2931A/T, -748G/A, -636A/G and IVS6-133delTCTCCCC, Additional file 3). Four SNPs (-4339T/C, -930T/C, 143A/C and IVS3+121T/C) of the 14 *UCP3* polymorphic sites were significantly different in allele frequency between the two breeds (Table 2). Despite the



**Table 1 Association analysis of UCP3 DNA polymorphisms with biochemical parameters among healthy Labrador Retrievers**

DNA polymorphism	Genotype	GLU	T-Cho	LDH	TG
UCP3 -4399C/T	CC (34)	97.1 ± 8.4	273.5 ± 49.0	55.9 ± 18.0	44.8 ± 21.1
	CT (16)	98.8 ± 14.5	237.2 ± 53.3	55.7 ± 13.5	49.4 ± 24.8
	TT (0)	-	-	-	-
	CC vs CT + TT	0.597	<b>0.021*</b>	0.965	0.504
UCP3 -4339T/C	TT (8)	94.5 ± 5.4	264.3 ± 49.6	55.3 ± 13.7	40.0 ± 13.5
	TC (27)	96.3 ± 11.5	276.9 ± 49.5	58.2 ± 20.9	50.4 ± 24.8
	CC (15)	101.7 ± 10.4	233.5 ± 51.2	51.9 ± 5.3	42.3 ± 20.5
	TT vs TC + CC	0.366	0.890	0.914	0.388
	TT + TC vs CC	0.079	<b>0.011*</b>	0.279	0.408
UCP3 -4010C/T	CC (15)	94.4 ± 11.9	255.5 ± 52.6	56.9 ± 16.1	51.2 ± 23.2
	CT (29)	98.7 ± 10.2	270.2 ± 52.2	56.5 ± 18.5	43.2 ± 21.5
	TT (6)	100.7 ± 8.6	237.7 ± 54.7	50.0 ± 0.0	48.7 ± 24.6
	CC vs CT + TT	0.159	0.580	0.763	0.310
	CC + CT vs TT	0.462	0.234	0.362	0.782
UCP3 -930T/C	TT (8)	94.5 ± 5.4	264.3 ± 49.6	55.3 ± 13.7	40.0 ± 13.5
	TC (27)	96.3 ± 11.5	276.9 ± 49.5	58.2 ± 20.9	50.4 ± 24.8
	CC (15)	101.7 ± 10.4	233.5 ± 51.2	51.9 ± 5.3	42.3 ± 20.5
	TT vs TC + CC	0.366	0.890	0.914	0.388
	TT + TC vs CC	0.079	<b>0.011*</b>	0.279	0.408
UCP3 -803C/T	CC (35)	96.9 ± 8.4	271.6 ± 49.5	55.7 ± 17.7	44.5 ± 20.9
	CT (15)	99.5 ± 14.8	239.1 ± 54.5	56.1 ± 13.9	50.5 ± 25.2
	TT (0)	-	-	-	-
	CC vs CT + TT	0.431	<b>0.045*</b>	0.950	0.388
UCP3 IVS3+26T/C	TT (15)	94.4 ± 11.9	255.5 ± 52.6	56.9 ± 16.1	51.2 ± 23.2
	TC (29)	98.7 ± 10.2	270.2 ± 52.2	56.5 ± 18.5	43.2 ± 21.5
	CC (6)	100.7 ± 8.6	237.7 ± 54.7	50.0 ± 0.0	48.7 ± 24.6
	TT vs TC + CC	0.159	0.580	0.763	0.310
	TT + TC vs CC	0.462	0.234	0.362	0.782
UCP3 IVS3+69G/A	GG (15)	94.4 ± 11.9	255.5 ± 52.6	56.9 ± 16.1	51.2 ± 23.2
	GA (29)	98.7 ± 10.2	270.2 ± 52.2	56.5 ± 18.5	43.2 ± 21.5
	AA (6)	100.7 ± 8.6	237.7 ± 54.7	50.0 ± 0.0	48.7 ± 24.6
	GG vs GA + AA	0.159	0.580	0.763	0.310
	GG + GA vs AA	0.462	0.234	0.362	0.782
UCP3 IVS5-115G/C	GG (15)	94.4 ± 11.9	255.5 ± 52.6	56.9 ± 16.1	51.2 ± 23.2
	GC (29)	98.7 ± 10.2	270.2 ± 52.2	56.5 ± 18.5	43.2 ± 21.5
	CC (6)	100.7 ± 8.6	237.7 ± 54.7	50.0 ± 0.0	48.7 ± 24.6
	GG vs GC + CC	0.159	0.580	0.763	0.310
	GG + GC vs CC	0.462	0.234	0.362	0.782
UCP3 IVS5-100T/C	TT (15)	94.4 ± 11.9	255.5 ± 52.6	56.9 ± 16.1	51.2 ± 23.2
	TC (29)	98.7 ± 10.2	270.2 ± 52.2	56.5 ± 18.5	43.2 ± 21.5
	CC (6)	100.7 ± 8.6	237.7 ± 54.7	50.0 ± 0.0	48.7 ± 24.6
	TT vs TC + CC	0.159	0.580	0.763	0.310
	TT + TC vs CC	0.462	0.234	0.362	0.782

**Table 1 Association analysis of UCP3 DNA polymorphisms with biochemical parameters among healthy Labrador Retrievers (Continued)**

UCP3	II (15)	94.4 ± 11.9	255.5 ± 52.6	56.9 ± 16.1	51.2 ± 23.2
1106delAAG	ID (29)	98.7 ± 10.2	270.2 ± 52.2	56.5 ± 18.5	43.2 ± 21.5
	DD (6)	100.7 ± 8.6	237.7 ± 54.7	50.0 ± 0.0	48.7 ± 24.6
	II vs ID + DD	0.159	0.580	0.763	0.310
	II + ID vs DD	0.462	0.234	0.362	0.782

Data are expressed as the mean ± SD.

*p*-values were calculated by ANOVA. \* and bold: *p* < 0.05.

I: insertion, D: deletion, IVS: intervening sequence.

Loci which were not observed polymorphism in Labrador retriever, or were not detected *p*-value are not shown.

different genetic background in each of the dog breeds [30-32], the different allele frequencies in the *UCP2* and *UCP3* polymorphic site between the two breeds may result from the susceptibility of Shetland Sheepdogs to hypercholesterolemia in a limited number of individuals.

The T allele at -4339T/C and -930T/C located in the *UCP3* intron 1 is associated with higher T-Cho levels, as shown by two different experiments: the association between polymorphisms and metabolic data (Table 1), and the distribution of allele of genotype in the breed that is susceptible to hypercholesterolemia (Table 2). These results suggest that the dog *UCP3* gene might be associated with T-Cho levels in a limited number of individuals.

It is known that the peroxisome proliferator activated receptors (PPAR) ligands activate *UCP3* expression [33,34]. The *UCP3* intron 1 contains that the putative binding elements of MyoG/MyoD, PPAR $\gamma$ /RXR $\alpha$  and SP1/SP3 that enhanced the *UCP3* gene transcription mainly regulated by PPARs in hamster, rat, and mouse [33]. Recently, we find the similar nucleotide sequences of the PPAR $\gamma$ /RXR $\alpha$  element in the dog *UCP3* intron 1 (Canine Genome Draft, NC\_006603.3). These findings imply that the dog *UCP3* intron 1 may be associated with regulation of *UCP3* gene expression. Further studies will be needed to demonstrate whether PPAR ligands bind or not this intronic region in dog.

With each genetic study, a different sample size is used to identify the candidate gene associating with genotypes and phenotypes in common diseases (multifactorial diseases) and/or single gene disorders. For instance, genome-wide association studies (GWAS) have reported the candidate gene associated with a mild form of disproportionate dwarfism using 23 cases and 37 controls [35], atopic dermatitis using 91 cases and 88 controls [36], and the chromosomal region of Patellar Luxation using 45 cases and 40 controls [37]. Some of the candidate genes were also tested using more than a hundred samples. The examination of larger sample sizes and further analysis will lead to increased precision of our

results. In addition, because the association analysis in this study was performed using only polymorphisms within the *UCP2* and *UCP3* genes, we cannot exclude the possibility that a gene that is closely linked to *UCP3* is causal.

## Conclusions

A genetic association study between polymorphisms in the dog uncoupling protein 2 and 3 genes and metabolic data showed that the SNPs of the *UCP3* intron 1 were associated with T-Cho levels in Labrador Retrievers. Alleles associated with high T-Cho levels of these polymorphisms were also present at higher frequencies in a breed that is susceptible to hypercholesterolemia (Shetland Sheepdogs), than in the control group (Shiba). The results obtained from a limited number of individuals suggest that the *UCP3* gene in dogs may be associated with total cholesterol levels. Therefore, the *UCP3* gene could be an interesting target, not only for lipid metabolism, but also for the treatment and prevention of obesity and metabolic-related diseases in dogs.

## Methods

### Animals and DNA

All animal experiments were approved by The Experimental Animal Ethics Committee in Nippon Veterinary and Life Science University. The blood samples were originally collected at the Veterinary Medical Teaching Hospital at NVLU with the written consent of each owner or the Kyushu Guide Dog Association. The collection of samples was handled by licensed veterinarians only.

Panel 1, for the first SNP discovery, was collected from 11 dogs that represented 11 different breeds: Miniature Dachshund, Welsh Corgi, Labrador Retriever, Shetland Sheepdog, Beagle, Yorkshire Terrier, Dobermann, Whippet, Weimaraner, Papillon, and Shiba. Panel 2 was used for SNP discovery and a study of associations between SNP variants and biochemical parameters; these samples were collected from 50 Labrador Retrievers. Panel 3 was used for SNP discovery and an interbreed analysis was

**Table 2 Genotyping data and interbreed analysis of DNA polymorphisms in UCP3**

UCP3 DNA polymorphism	Genotype	Number of samples		p	Allele	Allele frequency	
		Shiba	Shetland sheepdog			Shiba	Shetland sheepdog
-4399C/T	CC	29	30	NS	C	0.98	1.00
	CT	1	0	CC vs CT + TT	T	0.02	0.00
	TT	0	0				
-4339T/C	TT	0	4	p < 0.05	T	0.07	0.37
	TC	4	14	TT + TC vs CC	C	0.93	0.63
	CC	26	12				
-4160G/A	GG	30	30	ND	G	1.00	1.00
	GA	0	0		A	0.00	0.00
	AA	0	0				
-4010C/T	CC	18	13	NS	C	0.75	0.70
	CT	9	16	CC vs CT + TT	T	0.25	0.30
	TT	3	1				
-930T/C	TT	0	4	p < 0.05	T	0.07	0.37
	TC	4	14	TT + TC vs CC	C	0.93	0.63
	CC	26	12				
-803C/T	CC	30	30	ND	C	1.00	1.00
	CT	0	0		T	0.00	0.00
	TT	0	0				
143A/C	AA	20	30	p < 0.05	A	0.82	1.00
	AC	9	0	AA vs AC + CC	C	0.18	0.00
	CC	1	0				
IVS3+26T/C	TT	10	13	NS	T	0.57	0.70
	TC	14	16	TT vs TC + CC	C	0.43	0.30
	CC	6	1				
IVS3+69G/A	GG	18	13	NS	G	0.75	0.70
	GA	9	16	GG vs GA + AA	A	0.25	0.30
	AA	3	1				
IVS3+121T/C	TT	29	14	p < 0.05	T	0.98	0.67
	TC	1	12	TT vs TC + CC	C	0.02	0.33
	CC	0	4				
IVS5-115G/C	GG	10	4	NS	G	0.55	0.37
	GC	13	14	GG + GC vs CC	C	0.45	0.63
	CC	7	12				
IVS5-100T/C	TT	18	13	NS	T	0.75	0.70
	TC	9	16	TT vs TC + CC	C	0.25	0.30
	CC	3	1				
838T/C	TT	28	30	NS	T	0.97	1.00
	TC	2	0	TT vs TC + CC	C	0.03	0.00
	CC	0	0				
1106delAAG	ins ins	18	13	NS	ins	0.75	0.70
	ins del	9	16	II vs ID + DD	del	0.25	0.30
	del del	3	1				

I: insertion, D: deletion. IVS: intervening sequence.  
 p-values were calculated by Fisher's exact test. p < 0.05 NS: not significance. ND: not detection.

collected from 30 Shetland Sheepdogs and 30 Shibas containing each one animals from Panel 1. A list of breeds and number of individuals are presented in Table 3. Genomic DNA was extracted from whole blood with the Puregene kit (Qiagen, Valencia CA, USA).

### PCR

We used sequences of *UCP2* and *UCP3* (Canine Genome Draft, NC\_006603.3), to design 15 pairs of primers for amplification of each exon of the *UCP2* and *UCP3* genes (Table 4). Each PCR using TaKaRa Ex *Taq* was performed in a total volume of 25  $\mu$ l and contained 20 ng genomic DNA, 2.5  $\mu$ l 10 $\times$  Ex *Taq* Buffer (including 20 mM Tris-HCl, 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween 20, 0.5% Nonidet P-40, 50% Glycerol, 20 mM Mg<sup>2+</sup>), 0.4 mM of each primer, 200  $\mu$ M dNTP (dATP, dTTP, dCTP and dGTP), and 1U TaKaRa Ex *Taq* (TaKaRa, Shiga, Japan). Each PCR using FastStart *Taq* DNA polymerase (Roche, Basel, Switzerland) was performed in a total volume of 25  $\mu$ l and contained 20 ng genomic DNA, 2.5  $\mu$ l 10 $\times$  reaction Buffer (including 500 mM Tris-HCl, 100 mM KCl, 50 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 mM MgCl<sub>2</sub>), 0.4 mM of each primer (F12: 0.2 mM of each primer), 200  $\mu$ M dNTP (dATP, dTTP, dCTP and dGTP), and 1U FastStart *Taq* DNA polymerase. If

necessary, we used FastStart *Taq* for primer pairs that did not work with TaKaRa Ex *Taq*. The PCR reactions were performed on TaKaRa PCR Thermal Cycler Dice TP600 (TaKaRa). The conditions for PCR are shown in Table 5.

### Sequencing and SNP detection

The PCR products were purified with High Pure PCR Product Purification Kit (Roche). Cycle sequencing was then performed with the Big Dye Terminator v3.1 kit (Applied Biosystems, Foster City CA, USA); each reaction was run in a 10  $\mu$ l reaction volume containing 1  $\mu$ l purified PCR amplification product, 1  $\mu$ l Ready Reaction Premix, 1.5  $\mu$ l 5 $\times$  Big Dye Sequence Buffer, 1  $\mu$ l primer (1.6 pmol/ $\mu$ l), and 5.5  $\mu$ l sterile water. Cycle sequencing reactions were performed with the following conditions: 60 s at 96°C followed by 25 cycles of 10 s at 96°C, 5 s at 50°C and 4 min at 60°C. BigDye Xterminator Purification kits were used according to the manufacturer's instructions (Applied Biosystems) to purify dye-labeled fragments. Samples were analyzed on an ABI PRISM 310 genetic analyzer (Applied Biosystems). We identified DNA polymorphisms by comparing each sequence with the reference sequence (Canine Genome Draft, NC\_006603.3) by BLAST in NCBI (National Center for Biotechnology Information) and GENETYX program Ver. 11 (GENETYX Corporation, Tokyo, Japan). The position of identified DNA polymorphism was numbered from the A of the initiator methionine ATG codon as the +1 revealed in exon. In case of intron, a positive number indicates the number of nucleotides away from the previous exon, while a negative number indicates the number of nucleotide away from the next exon.

**Table 3 List of 119 DNA samples from 11 breeds**

DNA samples	Breeds	N	Sex	
			Male	Female
Panel 1 <sup>a</sup>	Miniature Dachshund	1		1
	Welsh Corgi	1	1	
	Labrador Retriever	1	1	
	Shetland Sheepdog	1	1	
	Beagle	1		1
	Yorkshire Terrier	1	1	
	Dobermann	1		1
	Whippet	1		1
	Weimaraner	1		1
	Papillon	1		1
	Shiba	1	1	
Panel 2 <sup>a,b,d</sup>	Labrador Retriever	50	27	23
Panel 3 <sup>a,b,c</sup>	Shetland Sheepdog	30 <sup>e</sup>	15	15
	Shiba	30 <sup>e</sup>	15	15
Total		119 <sup>f</sup>	60	59

N. Number of samples.

a. SNP discovery.

b. SNP genotyping.

c. Interbreed analysis.

d. Association analysis of DNA polymorphisms with biochemical parameters.

e. Include one individual of panel 1.

f. Total numbers of independent individuals.

### Measurement of biochemical parameters

Blood samples were collected into heparinized plastic tubes at least 12 h postprandial. Plasma was separated by centrifugation at 1500 $\times$  g for 10 min. Glucose (GLU), triglyceride (TG), total cholesterol (T-Cho), and lactate dehydrogenase (LDH) were measured using a Spotchem EM SP-4430 (Arkray, Kyoto, Japan) with the manufacturer's reagents.

### Statistical analysis

Deviation from the Hardy-Weinberg equilibrium was assessed by the Chi-squared test. SNPalyze (Dynacom, Chiba, Japan) was used to estimate haplotype frequencies. Genotype frequencies were compared using the Fisher's exact test. Differences of  $p < 0.05$  were considered statistically significant. Associations between genotype frequencies and metabolic data were analyzed by one-way analysis of variance (ANOVA).

**Table 4 Sequences of primers for PCR**

Gene	Fragment	Primer	Primer sequences (5'-3')	Range of PCR amplification <sup>a</sup>	size (bp)	Region <sup>b</sup>	
<i>UCP2</i>	F1	UCP2F1-F	CAGCTCTCGGCTTGTGAGC	24304468-24305048	581	Exon 1, Intron 1	
		UCP2F1-R	CACAACAGTCAGCAGACTGG				
	F2	UCP2F2-F	CCTTGCTGGAGTGAATCTG	24305288-24306125	838	Intron 1, Exon 2, Intron 2	
		UCP2F2-R	TGGGTTTGCCAGGTCTTTC				
	F3	UCP2F3-F	TACCAACTCTCCATACCTC	24307315-24308410	1096	Intron 2, Exon 3	
		UCP2F3-R	ATGCAGGCAGCTGTGCCAG				
	F4	UCP2F4-F	TGAGCAGGACAGGACTGTT	24308186-24308944	759	Exon 3, Intron 3, Exon 4, Intron 4	
		UCP2F4-R	AAAGGAGCTATACAGCAAATCA				
	F5	UCP2F5-F	TCTCAGAGCATTACTCTGCT	24309392-24310367	976	Intron 4, Exon 5, Intron 5, Exon 6, Intron 6	
		UCP2F5-R	AGAAAAGGCAGTCAGGACTC				
	F6	UCP2F6-F	TCCTCCCCCTCAAACCATCA	24310274-24311183	910	Intron 6, Exon 7, Intron 7, Exon 8	
		UCP2F6-R	GAAAGGGAGGTGGTGGGAA				
	<i>UCP3</i>	F7	UCP3F7-F	ATAGTACTTACCTCATAGGGT	24277647-24278722	1076	5'Fl, Exon 1, Intron 1
			UCP3F7-R	TATCTGTTCTCCATGGCAGC			
		F8	UCP3F8-F	CTAAGGAGCCTTAAGGGAAC	24278114-24278825	712	Exon 1, Intron 1
UCP3F8-R			TTCAGGGAGAGCTCAGGATC				
F9		UCP3F9-F	ACGCTACAGGTATGTGTGAG	24281537-24282266	730	Intron 1	
		UCP3F9-R	CCTGAAGGTACAGAGAGCC				
F10		UCP3F10-F	TAACTAACAGTTTAGGTGAGTC	24282174-24282933	760	Intron 1, Exon 2, Intron 2	
		UCP3F10-R	TGCTCAGAGTTCTGTGTGAAG				
F11		UCP3F11-F	CAGGTCCTTCTGCACCCAG	24283244-24284111	868	Intron 2, Exon 3, Intron 3, Exon 4, Intron 4	
		UCP3F11-R	TCATTCTGGGAGTTCCTCC				
F12		UCP3F12-F	CCTGTGGCCTTGCAACCAGA	24285138-24285396	259	Intron 4, Exon 5, Intron 5	
		UCP3F12-R	TGTTACCTCTGAGTGGTGCC				
F13		UCP3F13-F	GGCACCCTCAGAGGTAACA	24285377-24286088	712	Intron 5, Exon 6, Intron 6	
		UCP3F13-R	TGGGAAGGGATGTTGGATGC				
F14		UCP3F14-F	GCACTATCGTTACTCAAGG	24285748-24286088	341	Intron 5, Exon 6, Intron 6	
	UCP3F14-R	TGGGAAGGGATGTTGGATGC					
F15	UCP3F15-F	TAACTGCCTAACACAGAACC	24288288-24289004	717	Intron 6, Exon 7		
	UCP3F15-R	TTCAGCCTTCTGTACACA					

a. Number of nucleotide position is from canine genome draft (CGD) NC\_006603.3.

b. Fl: Flanking region Start codon is located in Exon 3 in *UCP2* and Exon 2 in *UCP3*. Stop codon is located in Exon 8 in *UCP2* and Exon 7 in *UCP3*.

## Additional files

**Additional file 1: Description and localization of identified DNA polymorphisms in dog *UCP2* and *UCP3* genes.**

**Additional file 2: Association analysis of *UCP2* DNA polymorphisms with biochemical parameters among healthy Labrador Retrievers.**

**Additional file 3: Genotyping data and interbreed analysis of DNA polymorphisms in *UCP2*.**

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

CU contributed to designing the study, genotypes, performing the statistical analysis, and drafting the manuscript. NT and JA participated in collection of data and clinical test. KI, KO, MB, and ST participated in study design and the manuscript editing. TO participated in experimental design, data collection, data analysis, and drafting of the manuscript. All authors read and approved the final manuscript.

## Acknowledgement

We are thankful to the Department of Veterinary Clinical Pathology and the medical staff at Nippon Veterinary and Life Science University (NVLU) in Musashino Tokyo, and to the Kyushu Guide Dog Association in Itoshima, Fukuoka for biological samples.

**Table 5 Conditions for PCR**

Fragment	Taq <sup>a</sup>	Initial denature (°C/m) <sup>b</sup>	Denature (°C/m) <sup>b</sup>	Annealing (°C/s) <sup>b</sup>	Extention (°C/m) <sup>b</sup>	Cycle	Final extension (°C/m) <sup>b</sup>
F1	F	95/4	95/1	57/30	72/1	35	72/7
F2	E	95/1	95/1	60/30	72/1	35	72/7
F3	F	95/4	95/1	60/10	72/1	30	72/7
F4	E	95/1	95/1	60/30	72/1	35	72/7
F5	F	95/4	95/1	60/30	72/1	35	72/7
F6	E	95/1	95/1	62/30	72/1	35	72/7
F7	E	95/1	95/1	60/15	72/1	34	72/7
F8	F	95/4	95/1	62/30	72/1	35	72/7
F9	F	95/4	95/1	62/30	72/1	35	72/7
F10	E	95/1	95/1	60/30	72/1	35	72/7
F11	F	95/4	95/1	62/15	72/1	32	72/7
F12	E	95/1	95/1	60/30	72/1	35	72/7
F13	E	95/1	95/1	60/30	72/1	35	72/7
F14	E	95/1	95/1	60/30	72/1	35	72/7
F15	E	95/1	95/1	60/30	72/1	35	72/7

a. Taq polymerase: E = ExTaq (TaKaRa), F = Fast start Taq (Roche).  
b. m: minutes s: seconds.

#### Author details

<sup>1</sup>Department of Basic Science, School of Veterinary Nursing and Technology, Faculty of Veterinary Science, Nippon Veterinary and Life Science University, 1-7-1 Kyonan-cho, Musashino, Tokyo 180-8602, Japan. <sup>2</sup>Department of Veterinary Nursing, School of Veterinary Nursing and Technology, Faculty of Veterinary Science, Nippon Veterinary and Life Science University, 1-7-1 Kyonan-cho, Musashino, Tokyo 180-8602, Japan. <sup>3</sup>Department of Veterinary Clinical Pathology, Nippon Veterinary and Life Science University, 1-7-1 Kyonan-cho, Musashino, Tokyo 180-8602, Japan. <sup>4</sup>Laboratory of Comparative Cellular Biology, Nippon Veterinary and Life Science University, 1-7-1 Kyonan-cho, Musashino, Tokyo 180-8602, Japan.

Received: 26 September 2014 Accepted: 25 November 2014  
Published: 11 December 2014

#### References

- Nicholls DG, Locke RM: Thermogenic mechanisms in brown fat. *Physiol Rev* 1984, **64**:1–64.
- Boss O, Muzzin P, Giacobino JP: The uncoupling proteins, a review. *Eur J Endocrinol* 1998, **139**:1–9.
- Ricquier D, Bouillaud F: The uncoupling protein homologues: UCP1, UCP2, UCP3, StUCP and AtUCP. *Biochem J* 2000, **345**:161–179.
- Schrauwen P, Walder K, Ravussin E: Human uncoupling proteins and obesity. *Obes Res* 1999, **7**:97–105.
- Kagawa Y, Yanagisawa Y, Hasegawa K, Suzuki H, Yasuda K, Kudo H, Abe M, Matsuda S, Ishikawa Y, Tsuchiya N, Sato A, Umetsu K, Kagawa Y: Single nucleotide polymorphisms of thrifty genes for energy metabolism: evolutionary origins and prospects for intervention to prevent obesity-related diseases. *Biochem Biophys Res Commun* 2002, **295**:207–222.
- Reis AF, Dubois-Laforgue D, Bellanné-Chantelot C, Timsit J, Velho G: A polymorphism in the promoter of UCP2 gene modulates lipid levels in patients with type 2 diabetes. *Mol Genet Metab* 2004, **82**:339–344.
- Yoon Y, Park BL, Cha MH, Kim KS, Cheong HS, Choi YH, Shin HD: Effects of genetic polymorphisms of UCP2 and UCP3 on very low calorie diet-induced body fat reduction in Korean female subjects. *Biochem Biophys Res Commun* 2007, **359**:451–456.
- Chan CB, Harper ME: Uncoupling proteins: role in insulin resistance and insulin insufficiency. *Curr Diabetes Rev* 2006, **2**:271–283.
- Lee HJ, Ryu HJ, Shin HD, Park BL, Kim JY, Cho YM, Park KS, Song J, Oh B: Associations between polymorphisms in the mitochondrial uncoupling proteins (UCPs) with T2DM. *Clin Chim Acta* 2008, **398**:27–33.
- Oktavianthi S, Trimarsanto H, Febinia CA, Suastika K, Saraswati MR, Dwipayana P, Arindarto W, Sudoyo H, Malik SG: Uncoupling protein 2 gene polymorphisms are associated with obesity. *Cardiovasc Diabetol* 2012, **11**:41.
- De Souza BM, Brondani LA, Bouças AP, Sortica DA, Kramer CK, Canani LH, Leitão CB, Crispim D: Associations between UCP1–3826A/G, UCP2–866G/A, Ala55Val and Ins/Del, and UCP3–55C/T polymorphisms and susceptibility to type 2 diabetes mellitus: case–control study and meta-analysis. *PLoS One* 2013, **8**:e54259.
- Schrauwen P, Xia J, Walder K, Snitker S, Ravussin E: A novel polymorphism in the proximal UCP3 promoter region: effect on skeletal muscle UCP3 mRNA expression and obesity in male non-diabetic Pima Indians. *Int J Obes Relat Metab Disord* 1999, **23**:1242–1245.
- Hamada T, Kotani K, Fujiwara S, Sano Y, Domichi M, Tszuzaki K, Sakane N: The common –55 C/T polymorphism in the promoter region of the uncoupling protein 3 gene reduces prevalence of obesity and elevates serum high-density lipoprotein cholesterol levels in the general Japanese population. *Metabolism* 2008, **57**:410–415.
- De Luis DA, Aller R, Izaola O, De La Fuente B, Conde R, Eiros Bouza JM: Relation of -55CT polymorphism of UCP3 gene with weight loss and metabolic changes after a high monounsaturated fat diet in obese non diabetic patients. *Eur Rev Med Pharmacol Sci* 2013, **17**:2810–2815.
- Salopuro T, Pulkkinen L, Lindström J, Kolehmainen M, Tolppanen AM, Eriksson JG, Valle TT, Aunola S, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Tuomilehto J, Laakso M, Uusitupa M: Variation in the UCP2 and UCP3 genes associates with abdominal obesity and serum lipids: the Finnish Diabetes Prevention Study. *BMC Med Genet* 2009, **10**:94.
- Edney AT, Smith PM: Study of obesity in dogs visiting veterinary practices in the United Kingdom. *Vet Rec* 1986, **118**:391–396.
- Jeusette IC, Lhoest ET, Istasse LP, Diez MO: Influence of obesity on plasma lipid and lipoprotein concentrations in dogs. *Am J Vet Res* 2005, **66**:81–86.
- Laflamme DP: Understanding and managing obesity in dogs and cats. *Vet Clin North Am Small Anim Pract* 2006, **36**:1283–1295.
- Ishioka K, Omachi A, Sagawa M, Shibata H, Honjoh T, Kimura K, Saito M: Canine adiponectin: cDNA structure, mRNA expression in adipose tissues and reduced plasma levels in obesity. *Res Vet Sci* 2006, **80**:127–132.



20. German AJ, Hervera M, Hunter L, Holden SL, Morris PJ, Biourge V, Trayhurn P: **Improvement in insulin resistance and reduction in plasma inflammatory adipokines after weight loss in obese dogs.** *Domest Anim Endocrinol* 2009, **37**:214–226.
21. Zoran DL: **Obesity in dogs and cats: a metabolic and endocrine disorder.** *Vet Clin North Am Small Anim Pract* 2010, **40**:221–239.
22. Tvarijonavičiute A, Ceron JJ, Holden SL, Cuthbertson DJ, Biourge V, Morris PJ, German AJ: **Obesity-related metabolic dysfunction in dogs: a comparison with human metabolic syndrome.** *BMC Vet Res* 2012, **8**:147.
23. Axelsson E, Ratnakumar A, Arendt ML, Maqbool K, Webster MT, Perloski M, Liberg O, Arnemo JM, Hedhammar A, Lindblad-Toh K: **The genomic signature of dog domestication reveals adaptation to a starch-rich diet.** *Nature* 2013, **495**:360–364.
24. Park HJ, Lee SE, Oh JH, Seo KW, Song KH: **Leptin, adiponectin and serotonin levels in lean and obese dogs.** *BMC Vet Res* 2014, **10**:113.
25. Kawasumi K, Kashiwado N, Okada Y, Sawamura M, Sasaki Y, Iwazaki E, Mori N, Yamamoto I, Arai T: **Age effects on plasma cholesterol and triglyceride profiles and metabolite concentrations in dogs.** *BMC Vet Res* 2014, **10**:57.
26. Ishioka K, Kanehira K, Sasaki N, Kitamura H, Kimura K, Saito M: **Canine mitochondrial uncoupling proteins: structure and mRNA expression of three isoforms in adult beagles.** *Comp Biochem Physiol B Biochem Mol Biol* 2002, **131**:483–489.
27. Udagawa C, Chong YH, Shito M, Kawakami T, Tada N, Ochiai K, Ishioka K, Tsuchida S, Omi T: **cDNA cloning and expression analysis of canine uncoupling protein 2 and 3 genes.** *J Pet Anim Nutr* 2011, **14**:68–75.
28. Sato K, Agoh H, Kaneshige T, Hikasa Y, Kagota K: **Hypercholesterolemia in Shetland sheepdogs.** *J Vet Med Sci* 2000, **62**:1297–1301.
29. Mori N, Lee P, Muranaka S, Sagara F, Takemitsu H, Nishiyama Y, Yamamoto I, Yagishita M, Arai T: **Predisposition for primary hyperlipidemia in Miniature Schnauzers and Shetland sheepdogs as compared to other canine breeds.** *Res Vet Sci* 2010, **88**:394–399.
30. **American Kennel Club Home Page.** [http://www.akc.org/breeds/shiba\\_inu/index.cfm](http://www.akc.org/breeds/shiba_inu/index.cfm).
31. Sugiyama S, Chong YH, Shito M, Kasuga M, Kawakami T, Udagawa C, Aoki H, Bonkobara M, Tsuchida S, Sakamoto A, Okuda H, Nagai A, Omi T: **Analysis of mitochondrial DNA HVR1 haplotype of pure-bred domestic dogs in Japan.** *Leg Med (Tokyo)* 2013, **15**:303–309.
32. Parker HG, Kim LV, Sutter NB, Carlson S, Lorentzen TD, Malek TB, Johnson GS, DeFrance HB, Ostrander EA, Kruglyak L: **Genetic structure of the purebred domestic dog.** *Science* 2004, **304**:1160–1164.
33. Hoffmann C, Zimmermann A, Hinney A, Volckmar AL, Jarrett HW, Fromme T, Klingenspor M: **A novel SP1/SP3 dependent intronic enhancer governing transcription of the UCP3 gene in brown adipocytes.** *PLoS One* 2013, **8**: e83426.
34. Villarroya F, Iglesias R, Giralt M: **PPARs in the Control of Uncoupling Proteins Gene Expression.** *PPAR Res* 2007, **2007**:74364.
35. Frischknecht M, Niehof-Oellers H, Jagannathan V, Owczarek-Lipska M, Drögemüller C, Dietschi E, Dolf G, Tellhelm B, Lang J, Tiira K, Lohi H, Leeb T: **A COL11A2 mutation in Labrador retrievers with mild disproportionate dwarfism.** *PLoS One* 2013, **8**:e60149.
36. Tengvall K, Kierczak M, Bergvall K, Olsson M, Frankowiack M, Farias FH, Pielberg G, Carlborg Ö, Leeb T, Andersson G, Hammarström L, Hedhammar Å, Lindblad-Toh K: **Genome-wide analysis in German shepherd dogs reveals association of a locus on CFA 27 with atopic dermatitis.** *PLoS Genet* 2013, **9**:e1003475.
37. Lavrijsen IC, Leegwater PA, Wangdee C, van Steenbeek FG, Schwencke M, Breur GJ, Meutstege FJ, Nijman IJ, Cuppen E, Heuven HC, Hazewinkel HA: **Genome-wide survey indicates involvement of loci on canine chromosomes 7 and 31 in patellar luxation in Flat-Coated Retrievers.** *BMC Genet* 2014, **15**:64.

doi:10.1186/1756-0500-7-904

**Cite this article as:** Udagawa et al.: The genetic association study between polymorphisms in uncoupling protein 2 and uncoupling protein 3 and metabolic data in dogs. *BMC Research Notes* 2014 **7**:904.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

