



**Figure 1.** Example configuration of the finite element method lung model (upper part) and representation of the resulting lung aeration in 0°, 15°, and 30° bed inclination (lower part).

Eline Oppersma, Ph.D.\*  
 Stijn Geraats, B.Sc.  
 Nicolien H. Heikens, B.Sc.  
 Arjen Welleweerd, B.Sc.  
 Chiel A. Woerts, B.Sc.  
 Libera Fresiello, Ph.D.  
 University of Twente  
 Enschede, the Netherlands

Christiaan L. Meuwese, M.D., Ph.D.  
 Erasmus Medical Center  
 Rotterdam, the Netherlands

Dirk W. Donker, M.D., Ph.D.  
 University of Twente  
 Enschede, the Netherlands  
 and

University Medical Center Utrecht  
 Utrecht, the Netherlands

ORCID IDs: 0000-0002-0150-306X (E.O.); 0000-0003-0972-8340 (L.F.);  
 0000-0002-1901-1653 (C.L.M.); 0000-0001-6496-2768 (D.W.D.).

\*Corresponding author (e-mail: e.mos-oppersma@utwente.nl).

4. Samimian S, Ashrafi S, Khaleghdoost Mohammadi T, Yeganeh MR, Ashraf A, Hakimi H, *et al.* The correlation between head of bed angle and intra-abdominal pressure of intubated patients; a pre-post clinical trial. *Arch Acad Emerg Med* 2021;9:e23.

Copyright © 2022 by the American Thoracic Society



## Reply to Oppersma *et al.*



*From the Authors:*

We thank Dr. Oppersma and colleagues for their interest in our study, in which we described the effects of changes in trunk inclination on respiratory mechanics in mechanically ventilated patients with COVID-associated acute respiratory distress syndrome (ARDS) (1). Reducing trunk inclination from semi-recumbent (40° head-of-bed elevation) to supine-flat (0° head-of-bed elevation) increased markedly (and reversibly) the compliance of the respiratory system, both due to an increase in chest wall and lung compliance. We did not measure changes in lung aeration (end-expiratory lung volume [EELV]); however, we think that the most relevant underlying mechanisms is the decrease in EELV in supine-flat position, favoring a reduction of

Ⓐ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.202205-0987LE on June 2, 2022

## References

- Marrazzo F, Spina S, Forlini C, Guarnieri M, Giudici R, Bassi G, *et al.* Effects of trunk inclination on respiratory mechanics in patients with COVID-19-associated acute respiratory distress syndrome: let's always report the angle! [letter] *Am J Respir Crit Care Med* 2022; 205:582–584.
- de Ryk J, Thiesse J, Namati E, McLennan G. Stress distribution in a three dimensional, geometric alveolar sac under normal and emphysematous conditions. *Int J Chron Obstruct Pulmon Dis* 2007;2:81–91.
- Lee HJ, Lee HJ, Lee JS, Kang YN, Koo JC. A built-up-type deformable phantom for target motion control to mimic human lung respiration. *Rev Sci Instrum* 2020;91:054106.

overdistension of some, most likely ventral non-dependent, lung regions. Of note, similar observations regarding changes in respiratory mechanics were made in patients with “classical” ARDS (2, 3). Moreover, a reduction in EELV caused by a change in trunk inclination from semi-recumbent to supine-flat position has been described in “classical” ARDS (2, 3), mechanically ventilated patients with normal lungs (4) and in spontaneously breathing subjects (5).

Dr. Oppersma and colleagues developed a mathematical model aimed at simulating the changes in lung aeration due to variations in trunk inclination. The model consisted of 15 blocks of homogenous material, with specific properties of either edematous, fibrotic or emphysematous lung tissue. In addition, the simulated pressure applied to the diaphragm (intra-abdominal pressure) changed according to body position, and the gravitational forces (likely proxies of pleural and abdominal pressure gradients) were adapted according to the applied angle. The authors simulated several combinations of mechanical properties of the 15 blocks, having as output variable lung aeration. They found that the combination of an edematous lung with apical emphysema and basal fibrosis predicted the largest increase in lung aeration changing position from supine-flat to semi-recumbent (in their experiment up to 30° head-of-bed elevation).

As stated above, the change in EELV associated with variations in trunk inclination is well-established (2–5). However, we agree with the authors that, in the specific context of ARDS, it might be of interest to understand the pathophysiologic mechanisms underlying the variable change in EELV induced by the variation of trunk inclination (2, 3). Heterogeneity of lung lesions/density is a key feature of ARDS and the distribution of lesions might play a role also in this context. In our study, patients were enrolled early, i.e., a median of 2.5 days after intubation. The timing of our observations makes it therefore difficult to ascertain the relative role of fibrosis and emphysema.

Another aspect worth discussing is that the authors focus on lung aeration, while they did not model (or present) information regarding  $V_T$  distribution. Besides changing lung aeration, i.e., the starting pulmonary volume, variations in trunk inclination most likely affect the pleural pressure and abdominal pressure gradients. As a result, it is conceivable that also the distribution of  $V_T$  changes with trunk inclination, ultimately affecting ventilation homogeneity.

In conclusion, we look forward to seeing a study in which their interesting computational model is integrated with patient data regarding lung aeration and  $V_T$  distribution (e.g., electric impedance tomography). We agree with the authors that this approach could improve our understanding regarding the effects of trunk inclination on respiratory mechanics in mechanically ventilated patients with ARDS. ■

**Author disclosures** are available with the text of this letter at [www.atsjournals.org](http://www.atsjournals.org).

Francesco Marrazzo, M.D.  
Stefano Spina, M.D.  
ASST Grande Ospedale Metropolitano Niguarda  
Milan, Italy

Roberto Fumagalli, M.D.  
Thomas Langer, M.D.\*  
ASST Grande Ospedale Metropolitano Niguarda  
Milan, Italy  
and  
University of Milan-Bicocca  
Milan, Italy

ORCID IDs: 0000-0001-5407-6615 (F.M.); 0000-0003-3210-1714 (S.S.); 0000-0003-0398-1329 (R.F.); 0000-0002-9725-8520 (T.L.).

\*Corresponding author (e-mail: [thomas.langer@unimib.it](mailto:thomas.langer@unimib.it)).

## References

1. Marrazzo F, Spina S, Forlini C, Guarnieri M, Giudici R, Bassi G, *et al*. Effects of trunk inclination on respiratory mechanics in patients with COVID-19-associated acute respiratory distress syndrome: let's always report the angle! *Am J Respir Crit Care Med* 2022;205:582–584.
2. Dellamonica J, Lerolle N, Sargentini C, Hubert S, Beduneau G, Di Marco F, *et al*. Effect of different seated positions on lung volume and oxygenation in acute respiratory distress syndrome. *Intensive Care Med* 2013;39:1121–1127.
3. Mezidi M, Guérin C. Effect of body position and inclination in supine and prone position on respiratory mechanics in acute respiratory distress syndrome. *Intensive Care Med* 2019;45:292–294.
4. Spooner AJ, Corley A, Sharpe NA, Barnett AG, Caruana LR, Hammond NE, *et al*. Head-of-bed elevation improves end-expiratory lung volumes in mechanically ventilated subjects: a prospective observational study. *Respir Care* 2014;59:1583–1589.
5. Behrakis PK, Baydur A, Jaeger MJ, Milic-Emili J. Lung mechanics in sitting and horizontal body positions. *Chest* 1983;83:643–646.

Copyright © 2022 by the American Thoracic Society



## Erratum: Optimizing the Design of Latent Tuberculosis Treatment Trials: Insights from Mathematical Modeling



A funding statement is missing from the opening footnote section in the article by Stout and colleagues (1), published in the March 1, 2020, issue of the *Journal*. There should have been a footnote for Dr. Nicholas A. Turner that read:

N.A.T. was funded by an Antibacterial Resistance Leadership Group fellowship from the National Institute of Allergy and Infectious Diseases (UM1 AI104681). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. ■

Ⓐ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).