



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Detection of SARS-CoV-2 in human breastmilk

It remains unclear whether severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can be shed into breastmilk and transmitted to a child through breastfeeding. Recent investigations have found no evidence of SARS-CoV-2 in human breastmilk, but sample sizes were small.¹⁻³ We examined milk from two nursing mothers infected with SARS-CoV-2. Both mothers were informed about the study and gave informed consent. Ethical approval for this case study was waived by the Ethics Committee of Ulm University and all samples were anonymised.

Clinical data and the timecourse of infection in the two mothers is shown in figure 1. After feeding and nipple disinfection, milk was

collected with pumps and stored in sterile containers at 4°C or -20°C until further analysis. We determined viral loads using RT-qPCR for SARS-CoV-2 *N* and *ORF1b-nsp14* genes⁴ in both whole and skimmed milk (obtained after removal of the lipid fraction). Further details of sample storage and processing are provided in the appendix. Following admission and delivery (day 0), four samples from Mother 1 tested negative (figure 2). By contrast, SARS-CoV-2 RNA was detected in milk from Mother 2 at days 10 (left and right breast), 12, and 13. Samples taken subsequently were negative (figure 2). Ct values for SARS-CoV-2 *N* peaked at 29.8 and 30.4 in whole milk and skimmed milk, respectively, corresponding to 1.32×10^5 copies per mL and 9.48×10^4 copies per mL (mean of both isolations). Since

milk components might affect RNA isolation and quantification, viral RNA recovery rates in milk spiked with serial dilutions of a SARS-CoV-2 stock were determined. We observed up to 89.2% reduced recovery rate in whole milk and 51.5% in skimmed milk (appendix), suggesting that the actual viral loads in whole milk of Mother 2 could be even higher than detected.

We detected SARS-CoV-2 RNA in milk samples from Mother 2 for 4 consecutive days. Detection of viral RNA in milk from Mother 2 coincided with mild COVID-19 symptoms and a SARS-CoV-2 positive diagnostic test of the newborn (Newborn 2). Mother 2 had been wearing a surgical mask since the onset of symptoms and followed safety precautions when handling or feeding the neonate (including proper hand and breast disinfection, strict washing,



Published Online
May 21, 2020
[https://doi.org/10.1016/S0140-6736\(20\)31181-8](https://doi.org/10.1016/S0140-6736(20)31181-8)

This online publication has been corrected. The corrected version first appeared at thelancet.com on September 10, 2020

See Online for appendix




Figure 1: Timecourse of SARS-CoV-2 infection of two mothers with newborn children

After delivery, Mother 1 developed mild COVID-19 symptoms and tested positive for SARS-CoV-2. Following spatial isolation of Mother 1 with her newborn (Newborn 1), Newborn 1 subsequently tested positive and developed respiratory problems, but both Mother 1 and Newborn 1 recovered. Mother 2 was admitted to the same hospital and room as Mother 1 and Newborn 1. Upon delivery, Mother 2 and Newborn 2 were brought back to the same room as Mother 1 and Newborn 1, and they stayed in the same room until Mother 1 tested positive for SARS-CoV-2 and isolated. Mother 2 and Newborn 2 were discharged on day 4. Mother 2 developed mild COVID-19 symptoms shortly thereafter and began wearing a surgical mask at all times of the day. Mother 2 tested positive for SARS-CoV-2 on day 8. 3 days later, Newborn 2 tested positive for SARS-CoV-2 and was readmitted to hospital because of newborn icterus and severe breathing problems. The child received phototherapy with blue light and ventilation therapy. Newborn 2 tested positive for RSV and SARS-CoV-2 at later timepoints. Mother 1 tested positive for SARS-CoV-2 again on day 22, 13 days after first being diagnosed. RT-qPCR analysis of breastmilk samples from both mothers revealed SARS-CoV-2 RNA in the milk of Mother 2 on days 10–13 (red bottles), whereas samples from Mother 1 were negative (white bottles). Dark shading indicates time from first SARS-CoV-2 positive oropharyngeal and nasopharyngeal swabs. Brackets indicate duration of COVID-19 symptoms. SARS-CoV-2=severe acute respiratory syndrome coronavirus 2. RSV=respiratory syncytial virus.

Submissions should be made via our electronic submission system at <http://ees.elsevier.com/thelancet/>




Figure 2: Detection of SARS-CoV-2 in breastmilk from an infected mother

SARS-CoV-2 RNA was isolated from whole and skimmed breastmilk obtained at different timepoints and analysed by RT-qPCR, using primer sets targeting SARS-CoV-2 N and ORF1b genes. Samples and viral RNA standard were run in duplicates, and isolation and RT-qPCR were repeated in two independent assays. RNA in breastmilk from Mother 2 on day 25 was only isolated once and only analysed by RT-qPCR for SARS-CoV-2 N. Symbols at baseline indicate no amplification (or Ct > 36.5 and no amplification in one replicate). Blue dashed line denotes quantification threshold for N (160 copies per reaction; Ct 34.2) and red dotted line for ORF1b (32 copies per reaction; Ct 35.9). Values below these lines but above baseline indicate amplification in both replicates, but no reliable quantification. Values shown represent mean (SD) from duplicates. SARS-CoV-2=severe acute respiratory syndrome coronavirus 2. Ct=cycle threshold.

and sterilisation of milk pumps and tubes). However, whether Newborn 2 was infected by breastfeeding or other modes of transmission remains unclear. Further studies of milk samples from lactating women and possible virus transmission via breastfeeding are needed to develop recommendations on whether mothers with COVID-19 should breastfeed.

We declare no competing interests. RG, CC, and JAM contributed equally. This work was supported by the EU's Horizon 2020 research and innovation programme (Fight-nCoV, 101003555 to JM) and the German Research Foundation (CRC 1279 to SS, FK, and JM; and MU 4485/1-1 to JAM)



Published Online
May 25, 2020
[https://doi.org/10.1016/S0140-6736\(20\)31209-5](https://doi.org/10.1016/S0140-6736(20)31209-5)

Rüdiger Groß, Carina Conzelmann,
Janis A Müller, Steffen Stenger,
Karin Steinhart, Frank Kirchhoff,
*Jan Münch
jan.muench@uni-ulm.de

Institute of Molecular Virology (RG, CC, JAM, FK, JM) and Institute for Microbiology and Hygiene (SS), Ulm University Medical Center, Ulm 89081, Germany; and Administrative District Heidenheim, Public Health Office, Heidenheim, Germany (KS)

- 1 Lackey KA, Pace RM, Williams JE, et al. SARS-CoV-2 and human milk: what is the evidence? *medRxiv* 2020; published online April 20. DOI:10.1101/2020.04.07.20056812 (preprint).
- 2 Yang N, Che S, Zhang J, et al. Breastfeeding of infants born to mothers with COVID-19: a rapid review. *medRxiv* 2020; published April 19. DOI:10.1101/2020.04.13.20064378 (preprint).
- 3 Chen L, Li Q, Zheng D, Jiang H, et al. Clinical characteristics of pregnant women with Covid-19 in Wuhan, China. *N Engl J Med* 2020; published April 17. DOI:10.1056/NEJMc2009226.
- 4 Chu DKW, Pan Y, Cheng SMS, et al. Molecular diagnosis of a novel coronavirus (2019-nCoV) causing an outbreak of pneumonia. *Clin Chem* 2020; **66**: 549–55.

COVID-19-associated hyperviscosity: a link between inflammation and thrombophilia?

Reports of thrombotic complications in patients with COVID-19 are increasingly prominent, and these reports include patients receiving

therapeutic anticoagulation.^{1,2} At our institution, multiple occurrences of anticoagulation failure prompted us to search for alternative aetiologies contributing to refractory hypercoagulability. Here we describe COVID-19-associated hyperviscosity, a potentially severe consequence of infection with severe acute respiratory syndrome coronavirus 2, in 15 patients tested to date. This work was done ethically in accordance with institutional review board approval.

All patients were critically ill with COVID-19 pneumonia and admitted to the medical intensive care unit. 14 patients had acute respiratory distress syndrome requiring intubation, 14 patients were encephalopathic, 12 patients had shock requiring vasopressors, and 11 patients had renal failure requiring continuous renal replacement therapy (CRRT). All patients received anticoagulation according to an institutional protocol based on