

RESEARCH

Open Access

A study of a population of *Nyssomyia trapidoi* (Diptera: Psychodidae) caught on the Pacific coast of Ecuador

Zapata S^{1,2}, Mejía L², Le Pont F³, León R², Pesson B^{1,4}, Ravel C⁵, Bichaud L⁶, Charrel R⁶, Cruaud C⁷, Trueba G² and Depaquit J^{1,7*}

Abstract

Background: Cutaneous leishmaniasis is endemic to the Pacific coast of Ecuador, and *Nyssomyia trapidoi* is considered to be its main vector. Dujardin *et al.* [1] recorded some differences in body pigmentation and isoenzymatic profiles in sympatric populations of *Ny. trapidoi* from the Pacific coast of Ecuador and suggested the existence of two cryptic species.

Methods: Entomological collections were performed in November 2008 and March 2011 in the locality of Paraíso Escondido using CDC miniature light traps and human bait. Morphological, isoenzymatic and molecular (sequencing of cytochrome *b* and cytochrome *c* oxidase 1 of the mitochondrial DNA) analyses, such as detection of *Leishmania* DNA and phlebovirus RNA in some females, were performed.

Results: Neighbor-joining trees from mitochondrial sequences grouped all of Ecuadorian *Ny. trapidoi* (including the two color variants) in one cluster, except for two specimens which clustered separately in both genes. Isoenzymatic characterization confirmed that the color variants belong to the same population. Additionally, 11.5% of females were found by PCR to contain *Endotrypanum monterogeii* kinetoplastid DNA. All pools of *Ny. trapidoi* were negative for phlebovirus RNA.

Conclusion: Analysis of mitochondrial gene sequences and isoenzymes was unable to support the existence of two sibling species within *Ny. trapidoi*, which is a probable vector of *Endotrypanum monterogeii*.

Keywords: Cryptic species, Isoenzymes, Mitochondrial DNA, Sympatry, *Endotrypanum*, Phlebovirus

Background

Currently, 76 species of phlebotomine sand flies belonging to 17 genera have been reported for Ecuador [2]; of these, *Nyssomyia trapidoi*, *Lutzomyia hartmanni*, *Lutzomyia gomezi* and *Lutzomyia ayacuchensis* are proven vectors of *Leishmania* parasites [3]. Leishmaniasis was first reported in Ecuador in 1920 [4] and is now endemic in the coastal region. It is present in 22 of the country's 24 provinces and purportedly has an incidence of

approximately 1,650 cases per year [5] (likely unknown) in various clinical forms: cutaneous (CL), mucocutaneous (MCL), diffuse cutaneous (DCL) and recidiva cutis (LRC) [3,6,7].

In the Pacific region of Ecuador, the sand fly species *Nyssomyia trapidoi* is distributed throughout the western foothills of the Andes and along the coast and is an important vector of *Leishmania* (*Viannia*) *panamensis* [3,8-10]. *Ny. trapidoi* is highly anthropophilic [11-13] and is commonly collected in secondary forests, crop plantations and close to human dwellings. The same features were observed in the original description of *Ny. trapidoi* by Fairchild & Hertig [14] who analyzed specimens from Panama (type-locality) and from the Ecuadorian Coast. A recent study in Panama found a high prevalence (43.3%) of *Leishmania naiffi* in *Ny. trapidoi* [15].

* Correspondence: jerome.depaquit@univ-reims.fr

¹ Université de Reims Champagne-Ardenne, ANSES, EA 4688 USC Transmission vectorielle et épidémiologie de maladies parasitaires, VECPAR, France

⁷ Genoscope, Centre National de Séquençage, 2 rue Gaston Crémieux, CP5706, 91057 Evry, Cedex, France

Full list of author information is available at the end of the article

In 1996, an entomological collection carried out in two communities on the north coast of Ecuador, obtained specimens of *Ny. trapidoi* showing a slight color variation. Additionally, isoenzymatic analysis revealed a possible presence of two species living in sympatry [1], although they did not correspond to the color variants. The purpose of the present study was to confirm the existence of two species under the name *Ny. trapidoi* at these locations. This report describes the results of an isoenzymatic and mitochondrial DNA analysis carried out on specimens of *Ny. trapidoi* collected in 2008 and 2011 at one of the locations described previously by Dujardin *et al.* [1]. Moreover, we investigated the vectorial role of *Ny. trapidoi* for Trypanosomatids and phleboviruses.

Methods

Sand fly collection

Sand flies were captured in November 2008 and March 2011 in the locality of Paraíso Escondido (00° 85' 03" N, 79° 17' 49" W), Pichincha Province, using CDC miniature light traps. Moreover, female sand flies were collected manually on human bait (captured on the skin of the authors). An out-group was analyzed consisting of specimens from Nicaragua (Musun), a place close to the type-locality of the species. Ecuadorian specimens were identified based on color phenotypes: light (B), dark (G) and indistinguishable (T). Specimens were killed using carbon dioxide and immediately stored in 96% ethanol for molecular studies and liquid nitrogen for isoenzymatic analysis. Once in the laboratory, the thorax of each specimen was separated and stored at -20°C for subsequent DNA extraction. The specimens used for isoenzymatic study and virus detection were processed as described below. The head, wings and genitalia of each specimen processed for morphology, molecular biology or isoenzyme analysis were cleared in boiling Marc-André [16] solution and mounted between slide and cover slide. The specimens selected for virus detection and isolation were individually identified in a drop of sterile saline solution under a stereomicroscope and pooled in groups of 50 belonging to the same species and of the same genus. The specimens we were unable to identify according to the latter method were stored at 80°C for future studies.

Isoenzyme analysis

Isoelectrofocusing was carried out in ultrathin agarose gels (MultiphorTM II Electrophoresis system, GE Healthcare Life Sciences) with the ampholyte at pH 4.0-6.5 in accordance with the protocols described by Pesson *et al.* [17]. The following isoenzyme systems were tested: malate dehydrogenase (MDH, E.C.1.1.1.37), isocitrate dehydrogenase (ICD, E.C.1.1.1.42), glycerol-3-phosphate

dehydrogenase (α GPDH, E.C.1.1.1.8), glucose-6-phosphate dehydrogenase (6PGD, E.C.1.1.1.44), hexokinase (HK, E.C.2.7.1.1), phosphoglucosmutase (PGM, E.C.5.4.2.2), fumarase (FUM, E.C.4.2.1.2), and glucose phosphate isomerase (GPI, E.C.5.3.1.9). The alleles for each locus were visualized as colored bands on the gels and numbered from the lowest to the highest pHi.

Allele frequencies, tests for deviation from Hardy-Weinberg equilibrium and Nei's [18] genetic distance between the two groups of sand flies were calculated using BIOSYS-2 [19]. Genotypic linkage disequilibrium and genotypic differentiation were tested using GENEPOP 3.3 [20].

Molecular taxonomy of phlebotomine sand flies

DNA was extracted using the QIAmp[®]DNA Mini Kit (Qiagen, Germany) and following the protocol used by Depaquit *et al.* [21]. Polymerase chain reactions (PCR) were performed for 30 *Ny. trapidoi* specimens. Each PCR was carried out in a 50 μ l volume using 5 μ l of DNA extracted solution and 50 pmol of the primers LepF and LepR [22] and C3B-PDR / NIN-PDR [23], as described previously, to amplify, respectively, cytochrome *c* oxidase 1 (COI) and cytochrome *b* genes from sand fly mitochondrial DNA. Amplification conditions for COI were as follows: an initial denaturation step at 94°C for 3 min followed by 5 cycles of 94°C denaturation for 30 s, 45°C annealing for 90 s, and 68°C extension for 60 s followed by 35 cycles of 94°C denaturation for 30 s, 51°C annealing for 30 s, 68°C extension for 60 s and, finally, a 68°C extension for 10 min [24,25]. For cytochrome *b*, an initial denaturation step at 94°C for 3 min was followed by 5 cycles (94°C for 30 s, 40°C for 30 s and 68°C for 60 s) then 35 cycles (94°C for 30 s, 44°C for 30 s and 68°C for 60 s), with a final extension at 68°C for 10 min [23].

PCR products were analyzed by electrophoresis in 1.5% agarose gels stained with 0.1% ethidium bromide. Amplicons were sequenced in both directions using the primers for DNA amplification.

DNA sequences were edited using the Pregap and Gap programs from the Staden Package software [26]. Sequence alignments were done using ClustalW software [27] and checked by eye. We selected a Neighbor-joining (NJ) analysis, which is a suitable method for intraspecific characterization of populations [28,29]. NJ was performed using MEGA 4.0 software [30]. Node support carried out using the NJ method was assessed by bootstrapping with 500 replications.

Trypanosomatid DNA detection

Detection of trypanosomatid DNA was carried out by PCR using two segments: a 120 bp kDNA corresponding to the conserved region of kinetoplast minicircles

(primers JW11, 5'-CCT ATT TTA CAC CAA CCC CCA GT-3' and JW12, 5'-GGG TAG GGG CGT TCT GCG AAA-3') [31] and the ribosomal small sub-units (SSU) (primers: KRD5, 5'-GATCTGGTTGATTCTGCCAGTAG-3' and KRD3, 5'-GATCCAGCTGCAGGTTACCTAC-3') [32]. As positive controls, DNA samples from *Leishmania* reference strains belonging to *L. panamensis*, *L. guyanensis*, *L. braziliensis* and *L. peruviana* were used. The conditions for these PCR reactions were the same as those described by Nicolas *et al.* [31] and Clark *et al.* [32].

Detection of phleboviruses in sandfly pools by RT-PCR

Twelve pools, each containing 50 *Ny. trapidoi* sand flies (a total of 550 females and 50 males) were ground using a Mixer Mill MM300 (Qiagen) with a 3-mm tungsten bead at a frequency of 30 cycles s⁻¹ for 3 min in the presence of 600 µl Eagle's minimal essential medium supplemented with 5% decomplexed calf serum, 1% L-Glutamine and 100 IU penicillin G ml⁻¹, 100 mg kanamycin ml⁻¹, 100 mg streptomycin ml⁻¹ and 7.5 µg amphotericin B ml⁻¹. The resulting mixture was clarified by centrifugation at 5800 g for 10 min and the supernatant fluid aliquoted (three aliquots of 200 µl for each pool) and stored at -80°C. One 200 µl aliquot was used for viral DNA and RNA purification using a BioRobot EZ1XL and the EZ1 Virus Mini kit (Qiagen), eluted into 90 µl and stored at -80°C. 10 µl was subsequently used in each RT-PCR reaction. Two sets of primers targeting different genes were used in independent reactions: (i) phlebovirus consensus primers targeting the polymerase gene in the L RNA segment [33], (ii) primers specific for the nucleoprotein gene in the S RNA segment of phleboviruses within the Sandfly fever Naples virus complex [34]. The RT-PCR cycling conditions consisted of 48°C for 45 min and 94°C for 2 min, followed by 40 cycles at 94°C for 30 s, 45°C for 1 min and 68°C for 45 s, with a

final elongation step at 68°C for 7 min. Nested PCRs were performed using the same conditions with 1.25 U Taq DNA polymerase (Invitrogen, USA). PCR products were visualized in a 2% TAE/agarose electrophoresis gel and sequenced in both directions either directly or after cloning in a TA cloning vector.

Virus isolation

Pooled sandfly homogenates stored at -80°C were used to inoculate Vero cells. Briefly, 100 µl each homogenate was diluted with 900 µl Eagle's minimal essential medium without fetal bovine serum (FBS), but enriched with antibiotics (100 IU penicillin G ml⁻¹, 100 mg streptomycin ml⁻¹, 100 mg kanamycin ml⁻¹ and 7.5 µg amphotericin B ml⁻¹) and used to seed Vero monolayers in a 12.5-cm² flask. After incubation at room temperature for 1 hr, 4 ml of fresh 5% FBS medium was added. The flask was incubated at 37°C in an atmosphere containing 5% CO₂. Flasks were examined daily for the presence of a cytopathic effect, and 400 µl supernatant was removed and tested by RT-PCR after viral RNA/DNA extraction as aforementioned.

Results and Discussion

Entomological captures using human baits resulted in 93% *Ny. trapidoi*, 4.2% *Psychodopygus panamensis* and 2.8% *Lutzomyia (Helcocyrtomyia) hartmanni*. Captures with CDC light traps yielded, in order of abundance: *Ny. trapidoi*, *Trichophoromyia reburra*, *Pressatia dysponeta*, *Psathyromyia aragoi*, *Psychodopygus panamensis*, *Lutzomyia hartmanni*, *Psathyromyia abunaensis* and *Psychodopygus carrerai thula* respectively (Table 1). Sixty-four *Ny. trapidoi* females were used for isoenzyme analysis and 550 *Ny. trapidoi* females and 50 *Ny. trapidoi* males were processed for virus detection. Of the males captured with CDC light traps, a small proportion was

Table 1 Phlebotomine sand fly species collected using two methods of capture

Genus	Species	Method of capture	Sex	Number of specimens	Percentage (%)
<i>Nyssomyia</i>	<i>trapidoi</i>	Human baits	♀	266	93
<i>Psychodopygus</i>	<i>panamensis</i>	Human baits	♀	12	4.2
<i>Helcocyrtomyia</i>	<i>hartmanni</i>	Human baits	♀	8	2.8
<i>Nyssomyia</i>	<i>trapidoi</i>	CDC	♀	108	38.7
<i>Trichophoromyia</i>	<i>reburra</i>	CDC	♀	102	36.6
<i>Pressatia</i>	<i>dysponeta</i>	CDC	♀	18	6.5
<i>Psathyromyia</i>	<i>aragoi</i>	CDC	♀	15	5.4
<i>Psychodopygus</i>	<i>panamensis</i>	CDC	♀	12	4.3
<i>Helcocyrtomyia</i>	<i>hartmanni</i>	CDC	♀	12	4.3
<i>Psathyromyia</i>	<i>abunaensis</i>	CDC	♀	6	2.2
<i>Psychodopygus</i>	<i>carrerai thula</i>	CDC	♀	6	2.2

subsequently studied. They belonged to the same species as the females and most were *Ny. trapidoi*. They were stored at -80°C .

Analysis of sand fly mitochondrial DNA

Mitochondrial COI and cytochrome *b* sequences from 30 *Ny. trapidoi* females were obtained and submitted to Genbank (accession numbers JQ 322908 through JQ 322968). Nucleotide sequences from COI amplicons (683 bp) revealed 31 variable sites, 10 of which were informative; cytochrome *b* (511 bp) contained 16 variable sites, two of them informative. The nucleotide composition was 37% T, 19% C, 28% A and 17% G for COI and 38% T, 15% C, 39% A and 9% G for cytochrome *b*. The alignment showed no indel in either sequence. COI sequences showed an average base substitution per site of 0.6% and cytochrome *b* sequences showed an average of 0.25%. There are more COI haplotypes than cyt *b* haplotypes. (Figure 1).

Analysis of isoenzyme polymorphism

Since MDH and ICD each showed two distinct loci, a total of 10 loci were studied. Six of these gave clearly interpretable polymorphic patterns: GPI, PGM, HK, FUM, 6PGD and MDH1. Samples tested for αGPDH were monomorphic. Table 2 shows the number of sand flies and allelic frequencies at each polymorphic locus.

Both populations of light and dark sand flies were in Hardy-Weinberg equilibrium ($P > 0.05$). When combined as one group, only PGM deviated ($P = 0.033$); this disequilibrium was due to a deficiency of heterozygotes. There was no significant linkage disequilibrium for either populations tested alone or grouped together ($1 > P > 0.180$). Nei's genetic distance value was 0.004 and there was no genotypic differentiation between light and dark *Ny. trapidoi* ($\chi^2 = 12.309$, $P = 0.421$).

Detection of trypanosomatid DNA

Out of 78 *Ny. trapidoi* females tested by PCR for the presence of trypanosomatid kDNA, 9 (11.5%) were




Figure 1 Neighbor joining trees obtained from nucleotide analysis of: A) COI mtDNA and B) cytochrome *b* mtDNA sequences of the 30 Ecuadorian specimens of *Nyssomyia trapidoi* and one from Nicaragua. Bootstrap values are shown in nodes (500 replicates).

Table 2 Allelic frequencies at the six polymorphic isoenzyme loci characterized in light and dark populations of *Nyssomyia trapidoi*

Locus	Populations	
	<i>Nyssomyia trapidoi</i> LIGHT	<i>Nyssomyia trapidoi</i> DARK
PGI		
n	33	31
1	0.879	0.806
2	0.076	0.065
3	0.045	0.113
4	0.000	0.016
P	1.000	0.558
PGM		
n	33	32
1	0.803	0.891
2	0.015	0.000
3	0.182	0.109
P	0.077	0.306
HK		
n	33	31
1	0.955	1.000
2	0.015	0.000
3	0.030	0.000
P	1.000	-
FUM		
n	22	23
1	0.932	1.000
2	0.068	0.000
P	1.000	-
6 PGD		
n	22	23
1	0.932	0.848
2	0.023	0.022
3	0.023	0.109
4	0.023	0.022
P	1.000	1.000
MDH 2		
n	21	22
1	0.976	1.000
2	0.024	0.000
P	1.000	-

n : sample size; P: probability of χ^2 value occurring by chance, when testing for deviation from Hardy-Weinberg expectations of genotypes frequencies.

positive. BLAST analysis of the 120 bp PCR products failed to reveal any significant score. However, 8 out of 78 kDNA PCR positive females were also PCR positive for the SSU rDNA locus. All the 2,135 bp sequences obtained from our samples were similar and showed

100% identity to the *Endotrypanum monterogei* sequence (X53911) described by Fernandes *et al.* [35]. One sequence was deposited in Genbank (JQ863389).

Detection of phleboviruses in sandfly pools by RT-PCR and by viral isolation

No pools were positive for phleboviruses using RT-PCR or by isolation on vero cells


Conclusion

The results of this study confirmed that *Ny. trapidoi* is the most common species captured [9] in the Paraíso Escondido community in the central coastal region of Ecuador. Two other anthropophilic species, *Psychodopygus panamensis* and *Lutzomyia hartmanni*, were recorded at this locality. Other non-anthropophilic species, i.e. *Trichophoromyia reburra*, *Pressatia dysponeta*, *Psathyromyia aragoi*, *Lutzomyia hartmanni*, *Psathyromyia abunaensis* and *Psychodopygus carrerai thula* were also collected there (Table 1).

Molecular (mtDNA sequencing) and isoenzymatic analyses did not support the existence of two populations (possible cryptic species) at Paraíso Escondido as previously hypothesized by Dujardin *et al.* [1]. The topology of the NJ tree obtained using cytochrome *b* sequences (Figure 1) grouped all Ecuadorian *Ny. trapidoi* into two clusters. The first cluster comprised the specimens B001 and B010 and the second cluster grouped the other specimens. This dichotomy is supported by only two nucleotide substitutions (positions 222 and 482). The *Ny. trapidoi* from Nicaragua differed from the Ecuadorian specimens in 12 nucleotide positions (Figure 2).

On the other hand, the NJ tree of COI sequences showed 5 clusters (Figure 1). The main branch included many of the specimens sharing the main cytochrome *b* haplotype. Another branch includes the B001 and B010 specimens, which grouped exactly as in the cytochrome *b* NJ clustering, thus perhaps warranting additional studies. These two specimens differ from the main haplotype by two nucleotide positions: 35 and 642. Specimens B007, B033 and G309 make up a third cluster that is supported mainly by one mutation. The fourth cluster (specimens B070, B071, G092 and T025) is strongly supported by seven positions (Figure 2). It is interesting to note that there is no correlation between haplotypes and color phenotypes. In this case, the COI gene provided a greater level of stratification than the cytochrome *b* gene.

Isozyme analysis suggested that the two color variants are conspecific *Ny. trapidoi*, and we did not find any variation of α GPDH profiles in our samples. However, the electrophoresis protocol used by Dujardin *et al.* [1] was different from that used in this study.



Ny. trapidoi is known to be an efficient vector for the vesicular stomatitis virus (Vesiculovirus) [36,37]. According to our study, there is no evidence for the circulation of any phlebovirus in *Ny. trapidoi* in the locality of Paraíso Escondido.

In this work, we processed 78 *Ny. trapidoi* females, 2 engorged on human blood and the rest unfed. Nine (11.5%) of the unfed sand flies had identical trypanosomatid kDNA sequences and showed no homology to *Leishmania panamensis* and *L. guyanensis*, species commonly found in *N. trapidoi* [3]. However, SSU rDNA sequences (2135 bp) from the same specimens showed 100% homology with the *Endotrypanum monterogeii* sequence (GenBank X53911) obtained from a sloth (*Choloepus hoffmanni*) and described by Fernandes et al. [35]. Moreover, our sequences showed a high homology (over more than 2100 bp) to sequences described as *Endotrypanum* sp. by Ferreira et al. [38]: 99.7%, homology to *Endotrypanum* isolated from *Lutzomyia gomezi* (EU21238) and 99.8-99.9% to isolates from *Psathyromyia dendrophyla* (EU21239, EU21240). If we consider this variability as intraspecific, it might suggest that *E. monterogeii* has a wide

spectrum of vectors, including the genera *Lutzomyia*, *Psathyromyia* and *Nyssomyia*. Although sloths were common in study sites some time ago, they are now rare, and other vertebrates may play a role as reservoirs. Further studies are needed to learn more about the vertebrate hosts of *E. monterogeii*, since to date our knowledge is limited to *C. hoffmanni* [35,39].

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SZ, FLP, RL & JD were responsible for the conception and design of the work and have been involved in drafting the manuscript. SZ & BP carried out the isoenzymatic analysis and have given final approval of the version to be published. SZ, JD and CR did the parasitologic study and have given final approval of the version to be published. CC did the sequencing of the samples through the project @speed-ID and have given final approval of the version to be published. LM & GT helped with collecting and processing the samples, revising the manuscript critically and have given final approval of the version to be published. LB and RC carried out the viral study, critically reviewed the manuscript and have given final approval of this version to be submitted for publication.

Acknowledgements

The authors are grateful to Anita Freire, Soledad Tello and Moisés Gualapuro for their invaluable support in entomological collection and field work.

This work is part of the @SPEED-ID "Accurate SPECies Delimitation and IDentification of Eukaryotic Biodiversity using DNA Markers" project proposed by F-BoL, the French Barcode of Life initiative, funded by the French National Center of Sequencing (Genoscope) and supported by the National Secretariat for Higher Education, Science, Technology and Innovation (SENESCYT) of Ecuador.

Author details

¹Université de Reims Champagne-Ardenne, ANSES, EA 4688 USC Transmission vectorielle et épidémiologie de maladies parasitaires, VECPAR, France. ²Instituto de Microbiología, Universidad San Francisco de Quito, Quito, Ecuador. ³72 Rue de la Colonie, Paris, France. ⁴Faculté de Pharmacie, Université de Strasbourg, Strasbourg, France. ⁵French Reference Centre on Leishmaniasis, University of Montpellier, UMR5290, 39 Av. Charles Flahault, F34295, Montpellier, France. ⁶Aix Marseille Univ, IRD French Institute of Research for Development, EHESP French School of Public Health, UMR_D 190 "Emergence des Pathologies Virales, and IHU Mediterranee Infection, APHM Public Hospitals of Marseille, 13005, Marseille, France. ⁷Genoscope, Centre National de Séquençage, 2 rue Gaston Crémieux, CP5706, 91057 Evry, Cedex, France.

Received: 12 April 2012 Accepted: 23 July 2012
Published: 23 July 2012

References

- Dujardin JP, Le Pont F, Cruz M, Leon R, Tarrieu F, Guderian R, Tibayrenc M: **Cryptic speciation in *Lutzomyia* (*Nyssomyia*) *trapidoi* (Fairchild and Hertig) (Diptera: Psychodidae) detected by multilocus enzyme electrophoresis.** *Am J Trop Med Hyg* 1996, **54**:42–45.
- Galati EAB: *Phlebotominae (Diptera, Psychodidae) Classificação, Morfologia, Terminologia e Identificação de Adultos*. Apostila Disciplina HEP 5752 *Bioecologia e Identificação de Phlebotominae Vol. I*. Departamento de Epidemiologia Faculdade de Saúde Pública Universidade de São, São Paulo, SP: Brasil. Press; 2010.
- Calvopiña M, Armijos RX, Hashiguchi Y: **Epidemiology of leishmaniasis in Ecuador: current status of knowledge - a review.** *Mem Inst Oswaldo Cruz* 2004, **99**(7):663–672.
- Rodríguez JD: **Lecciones de Parasitología Humana**. In *Género Leishmania*. 5th edition: Departamento de publicaciones de la Universidad de Guayaquil; 1974:170–185.
- Servicio nacional de control de la malaria y otras enfermedades transmitidas por vectores artrópodos "Juan Montalván Cornejo" (SNEM)*. 2010.
- Hashiguchi Y, Gomez EA: **A review of leishmaniasis in Ecuador.** *Bull Pan Am Health Organ* 1991, **25**(1):64–76.
- Calvopiña M, Uezato H, Gomez EA, Korenaga M, Nonaka S, Hashiguchi Y: **Leishmaniasis recidiva cutis due to *Leishmania* (Viannia) *panamensis* in subtropical Ecuador: isoenzymatic characterization.** *Int J Dermatol* 2006, **45**(2):116–120.
- Armijos RX, Chico ME, Cruz ME, Guderian RH, Kreutzer RD, Berman JD, Rogers MD, Grögl M: **Human cutaneous leishmaniasis in Ecuador: identification of parasites by enzyme electrophoresis.** *Am J Trop Med Hyg* 1990, **42**(5):424–428.
- Le Pont F, Leon R, Guerrini F, Gantier JC, Mouchet J, Echeverría R, Guderian RH: **Leishmaniasis in Ecuador. 3. *Lutzomyia trapidoi*, vector of *Leishmania panamensis*.** *Ann Soc Belg Med Trop* 1994, **74**(1):23–28.
- Dantas-Torres F: **Canine leishmaniasis in South America.** *Parasites & Vectors*. 2009, **26**(2 Suppl 1):S1.
- Le Pont F, Leon R, Mouchet R, Echeverría R, Guderian RH: **Leishmaniose en Equateur. 2. Contacts Homme/vecteurs de Leishmaniose: Cas de *Lutzomyia trapidoi* et *Lutzomyia gomezi*.** *Ann Soc Belg Med Trop* 1994, **74**(1):13–21.
- Mouchet J, Le Pont F, Leon R, Echeverría R: **Leishmaniasis in Ecuador. 5. Leishmaniasis and anthropization on the Pacific coast.** *Ann Soc Belg Med Trop* 1994, **74**(1):35–41.
- Barrera C, Herrera M, Martínez F, Leon R, Richard A, Guderian RH, Mouchet R, Echeverría R, Le Pont F: **Leishmaniose en Equateur. 1. Incidence de la Leishmaniose Tegumentaire sur la Façade Pacifique.** *Ann Soc Belg Med Trop* 1994, **74**(1):1–12.
- Fairchild GB, Hertig M: **Notes on the *Phlebotomus* of Panama (Diptera, Psychodidae). IX. Description of seven new species.** *Ann Entomol Soc Amer* 1952, **45**:505–528.
- Azpuruja J, De La Cruz D, Valderama A, Windsor D: ***Lutzomyia* Sand Fly Diversity and Rates of Infection by *Wolbachia* and an Exotic *Leishmania* Species on Barro Colorado Island, Panama.** *PLoS Negl Trop Dis* 2010, **4**(3):e627.
- Abonnenc E: **Les phlébotomes de la région éthiopienne (Diptera: Phlebotomidae).** *Mémoires de l'ORSTOM* 1972, **55**:1–289.
- Pesson B, Wallon M, Floer MT, Kistensen AR: **Étude isoenzymatique de populations méditerranéennes de phlébotomes du sous-genre *Larrousius*.** *Parassitologia* 1991, **33**(1):471–476.
- Nei M: **Genetic distance between populations.** *Am Nat* 1972, **106**(949):283–292.
- Swofford DL, Selander RB: **BIOSYS-1: a FORTRAN program for the comprehensive analysis of electrophoretic data in population genetics and systematics.** *J Hered* 1981, **72**(4):281–283.
- Raymond M, Rousset F: **GENEPOP (version 12): population genetics software for exact tests and ecumenicism.** *J Hered* 1995, **86**(3):248–249.
- Depaquit J, Lienard E, Verzeaux-Griffon A, Ferté H, Bounamous A, Gantier JC: **Molecular homogeneity in diverse geographical populations of *Phlebotomus papatasi* (Diptera, Psychodidae) inferred from ND4 mtDNA and ITS2 rDNA epidemiological consequences.** *Infect Genet Evol* 2008, **8**(2):159–170.
- Hajjibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN: **DNA barcodes distinguish species of tropical Lepidoptera.** *Proc Natl Acad Sci USA* 2006, **103**(4):968–997.
- Esseghir S, Ready PD, Killick-Kendrick R, Ben-Ismaïl R: **Mitochondrial haplotypes and phylogeography of *Phlebotomus* vectors of *Leishmania major*.** *Insect Molec Biol* 1997, **6**(3):211–225.
- Costa FO, DeWaard JR, Boutillier J, Ratnasingham S, Dooh RT, Hajjibabaei M, Hebert PDN: **Biological identifications through DNA barcodes: the case of the Crustacea.** *Can J Fish Aquat Sci* 2007, **64**:272–295.
- Boudabous R, Bounamous A, Jouet D, Depaquit J, Augot D, Ferté H, Berchi S, Veuille M, Babba H: **Mitochondrial DNA differentiation between two closely related species *Phlebotomus chabaudi* Croset, Abonnenc & Rioux, 1970 and *P. (Pa.) riouxi* Depaquit, Killick-Kendrick & Léger, 1998 based on direct sequencing and PCR-RFLP.** *Ann Entomol Soc Am* 2009, **102**(1):347–353.
- Bonfield JK, Staden R: **Experiment files and their application during large-scale sequencing projects.** *DNA Seq* 1996, **6**:109–117.
- Thompson JD, Higgins DG, Gibson TJ: **CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice.** *Nucleic Acids Res* 1994, **22**(22):4673–4680.
- Lecointre G: **Que représente-t-on avec un arbre ?** *Biosystema* 2011, **28**:9–40.
- Tassy P, Barriol V: **L'homologie, l'arbre généalogique et le cladogramme: un apologue.** *Bull Soc Zool Fr* 1995, **120**:361–378.
- Kumar S, Tamura K, Nei M: **MEGA3: Integrated software for Molecular Evolutionary Genetics Analysis and sequence alignment.** *Brief Bioinform* 2004, **5**(2):150–163.
- Nicolas L, Prina E, Lang T, Milon G: **Real-Time PCR for Detection and Quantitation of *Leishmania* in Mouse Tissues.** *J Clin Microbiol* 2002, **40**(5):1666–1669.
- Clark CG, Martin DS, Diamond LS: **Phylogenetic relationships among anuran trypanosomes as revealed by riboprinting.** *J Eukaryot Microbiol* 1995, **42**(1):92–96.
- Sánchez-Seco MP, Echeverría JM, Hernández L, Estévez D, Navarro-Marí JM, Tenorio A: **Detection and identification of Toscana and other phleboviruses by RT-nested-PCR assays with degenerated primers.** *J Med Virol* 2003, **71**(1):140–149.
- Charrel RN, Izri A, Temmam S, Delaunay P, Toga I, Dumon H, Marty P, de Lamballerie X, Parola P: **Cocirculation of 2 genotypes of Toscana virus, southeastern France.** *Emerg Infect Dis* 2007, **13**:465–468.
- Fernandes AP, Nelson K, Beverley SM: **Evolution of nuclear ribosomal RNAs in kinetoplastid protozoa: perspectives on the age and origins of parasitism.** *PNAS* 1993, **90**(24):11608–11612.
- Tesh RB, Chaniotis BN, Johnson KM: **Vesicular stomatitis virus, Indiana serotype: multiplication in and transmission by experimentally infected phlebotomine sandflies (*Lutzomyia trapidoi*).** *Am J Epidemiol* 1971, **93**(6):491–495.
- Tesh RB, Chaniotis BN, Johnson KM: **Vesicular stomatitis virus (Indiana serotype): transovarial transmission by phlebotomine sandflies.** *Science* 1972, **175**(4029):1477–1479.

38. Ferreira RC, De Souza AA, Freitas RA, Campaner M, Takata CS, Barrett TV, Shaw JJ, Teixeira MM: **A Phylogenetic Lineage of Closely Related Trypanosomes (Trypanosomatidae, Kinetoplastida) of Anurans and Sand Flies (Psychodidae, Diptera) Sharing the Same Ecotopes in Brazilian Amazonia.** *J Eukaryot Microbiol* 2008, **55**(5):427–435.
39. Katakura K, Mimori T, Furuya M, Uezato H, Nonaka S, Okamoto M, Gomez L EA, Hashiguchi Y: **Identification of *Endotrypanum* species from a sloth, a squirrel and *Lutzomyia* sandflies in Ecuador by PCR amplification and sequencing of the mini-exon gene.** *J Vet Med Sci* 2003, **65**(5):649–653.

doi:10.1186/1756-3305-5-144

Cite this article as: Zapata *et al.*: A study of a population of *Nyssomyia trapidoi* (Diptera: Psychodidae) caught on the Pacific coast of Ecuador. *Parasites & Vectors* 2012 **5**:144.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

