

Research article

Open Access

Impact of imatinib on the pharmacokinetics and *in vivo* efficacy of etoposide and/or ifosfamide

Keyvan Rezaï¹, François Lokiec¹, Isabelle Grandjean², Sophie Weill¹, Patricia de Cremoux³, Vincent Bordier², Richard Ekue², Mickael Garcia², Marie-France Poupon⁵ and Didier Decaudin^{*4,6}

Address: ¹Department of Pharmacology Oncology, Centre René Huguenin, Saint-Cloud, France, ²Unit of Animal experiments, Research Section, Institut Curie, Paris, France, ³Department of tumor Biology, Institut Curie, Paris, France, ⁴Department of Clinical Hematology, Institut Curie, Paris, France, ⁵FRE 2584, Section de Recherche, Institut Curie, Paris, France and ⁶UMR144 CNRS/Institut Curie, Paris, France

Email: Keyvan Rezaï - k.rezai@stcloud-huguenin.org; François Lokiec - lokiec@crh1.org; Isabelle Grandjean - isabelle.grandjean@curie.fr; Sophie Weill - s.weill@stcloud-huguenin.org; Patricia de Cremoux - decremoux@curie.net; Vincent Bordier - vincent.bordier@curie.fr; Richard Ekue - richard.ekue@curie.fr; Mickael Garcia - mickael.garcia@curie.fr; Marie-France Poupon - marie-france.poupon@curie.fr; Didier Decaudin* - didier.decaudin@curie.net

* Corresponding author

Published: 27 October 2007

Received: 12 February 2007

BMC Pharmacology 2007, 7:13 doi:10.1186/1471-2210-7-13

Accepted: 27 October 2007

This article is available from: <http://www.biomedcentral.com/1471-2210/7/13>

© 2007 Rezaï et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Using a human small cell lung cancer (SCLC) xenografted in *nude* mice, we have previously reported enhanced tumor growth inhibition following chemotherapy in combination with imatinib (STI571). We therefore investigated the *in vivo* impact of imatinib on the pharmacokinetics and efficacy of chemotherapy.

Methods: Two different human tumors were used: SCLC6 small cell lung cancer xenografted in *nude* mice, and LY-3 EBV-associated human B-cell lymphoma xenografted in SCID mice. Plasma, urine, and fecal concentrations of etoposide (VP16) were determined by a validated high performance liquid chromatography method. Plasma concentrations of ifosfamidewere determined by a validated gas chromatography assay with nitrogen-phosphorus detection.

Results: Slight tumor growth inhibition was induced by imatinib administered alone in one *in vivo* EBV-associated B-cell lymphomatous xenograft. In contrast, an increase of the chemotherapy-induced antitumor effect was observed in the lymphoma model but not in a small cell lung cancer model when mice bearing human xenografted tumors were treated concomitantly by imatinib and chemotherapy. This antitumor effect was not influenced by concomitant administration of fluconazole. The AUC0-3 h (Area Under the concentration-time Curve) of etoposide was increased when mice were treated with etoposide + imatinib due to decreased fecal excretion. In contrast, imatinib did not appear to influence the urinary excretion of etoposide, and concomitant administration of the CYP3A4 inhibitor, fluconazole, with imatinib did not modify the pharmacokinetics of etoposide plus imatinib alone.

Conclusion: Altogether, these results therefore justify further prospective phase I and II clinical trials with combinations of etoposide-based chemotherapy and imatinib in patients with certain cancers, such as malignant lymphoma, with careful toxicologic monitoring.

Background

The tyrosine kinase inhibitor imatinib (STI571), belonging to the 2-phenylaminopyrimidine class, selectively inhibits BCR/ABL [1], PDGFR ("Platelet-Derived Growth Factor receptor"), c-kit [2], and c-fms ("Macrophage colony-stimulating factor receptor")[3] kinase activity. As imatinib is commonly used without chemotherapeutic agents, few reports have evaluated the therapeutic effect of concomitant administration of imatinib and chemotherapy either in mice xenografted tumors or cancer patients. Using a human small cell lung cancer (SCLC) xenograft in *nude* mice, we previously reported enhanced tumor growth inhibition following chemotherapy (etoposide + ifosfamide or topotecan) in combination with imatinib and showed that this effect was not dependent on c-kit expression level [4]. The increase of conventional antineoplastic agent-induced tumor growth inhibition was also exclusively observed when imatinib and chemotherapy were administered concomitantly. Various mechanisms can be proposed to explain this effect, namely (1) an increase of drug uptake by imatinib-induced reduction of tumor interstitial fluid pressure and an increase of transcapillary transport [5], (2) inhibition of tumor angiogenesis combined with the antitumor effect of chemotherapy [6-8], and (3) certain intracellular events induced by the two concomitant therapeutic modalities that remain to be determined.

However, our *in vivo* experiments showed a higher toxicity of combined imatinib and chemotherapy than for imatinib or chemotherapy alone, exclusively when both treatments were administered concomitantly. Various mechanisms for the toxicity of combined therapy can be proposed, particularly pharmacokinetic interactions between imatinib and chemotherapeutic agents that could also explain the increase of chemotherapy-induced tumor growth inhibition, as numerous data have shown that imatinib induces cytochrome p450 inhibition and that this inhibition decreases elimination of chemotherapeutic agents such as etoposide, ifosfamide, and topotecan [9-12]. In order to explore these pharmacokinetic interactions, we therefore investigated the *in vivo* impact of imatinib on the pharmacokinetics of chemotherapy and its efficacy on two different human cancers xenografted into immunodeficient mice: a SCLC [4] and an EBV-associated B-cell lymphoproliferation [13]. We showed that imatinib significantly increases the AUC (Area Under the concentration-time Curve) of etoposide in mice via a decrease of its fecal excretion. Finally, in one of the two xenografted models used, we confirmed the imatinib-induced increase of tumor growth inhibition after chemotherapy with agents such as etoposide (VP16) and gemcitabine. These results therefore support the initiation of further prospective phase I and II clinical trials combining etoposide-based chemotherapy and imatinib

in cancer patients, such as malignant lymphoma patients, with careful toxicologic monitoring.

Methods

In vivo experiments in immunodeficient mice bearing human tumors

Female *nude* or SCID mice, weighing 20 g to 30 g, 6–8 weeks old, were bred in the animal facilities (Institut Curie, Paris, France), maintained under specific pathogen-free conditions with artificial lighting (12-hour light/12-hour dark cycle) and fed with a regular diet and water *ad libitum*. The care, housing, and handling of the mice were performed in accordance with the recommendations of the French Ethics Committee and under the supervision of authorized investigators. For curative therapeutic trials, the tumor-bearing mice were randomly divided into equivalent groups of 6 to 8 animals and mice were treated at different times after transplantation.

Two different human tumors were used: the SCLC6 small cell lung cancer xenografted in *nude* mice [4], and the LY-3 EBV-associated human B-cell lymphoma xenografted in SCID mice [13]. STI571 (gift from Novartis Pharma SAS, Rueil-Malmaison, France) was diluted in 150 μ l of H₂O and administered at a total dosage of 70 mg/kg per day in one intraperitoneal injection on different days, as indicated. Etoposide (VP16)(Pierre Fabre, Boulogne, France) and ifosfamide (Baxter, Paris, France) were diluted in 200 μ l of 0.9% sodium chloride and administered at a dose of 12 mg/kg and 90 mg/kg, respectively, in one daily intraperitoneal injection on days 1 to 3 of treatment. Fluconazole (PFIZER Paris, France) was diluted in 400 μ l of water and administered at a total dosage of 40 mg/kg per day in one intraperitoneal injection on days 1 to 3 of treatment. Gemcitabine (Lilly France SAS, Suresnes, France) was diluted in 150 μ l of 0.9% sodium chloride and administered in one weekly intraperitoneal injection at a dosage of 60 mg/kg per day. The control group received 0.9% sodium chloride injections according to the same schedule as experimentally treated mice.

All mice were weighed once weekly. Tumor growth was monitored by measuring two perpendicular diameters with calipers. Tumor volume (V) and relative tumor volume (RTV) were calculated as follows:

$$V = a^2 \times b/2,$$

where *a* is the width (large diameter) and *b* the length (small diameter) of the tumor in millimeters.

$$RTV = V_x/V_i,$$

where V_x is the mean tumor volume in cubic millimeters at any given time and V_i is the mean initial tumor volume

in cubic millimeters at the start of treatment [14]. Mice were ethically sacrificed when the tumor volume reached 2,500 mm³ in the control group.

Pharmacokinetics of etoposide (VP16) and/or ifosfamide in mice

In order to evaluate pharmacokinetic interactions between STI571 (imatinib) and chemotherapeutic agents (etoposide and ifosfamide), various concentrations of the two cytotoxic agents were measured. Plasma, urine, and fecal concentrations of etoposide were determined by a validated high performance liquid chromatography (HPLC) method with U.V. detection [15]. Standard samples were prepared from a stock solution of 1000 µg/ml etoposide which was added to drug-free pooled plasma or urine. The analytes were extracted from plasma, urine and feces by liquid-liquid extraction into aqueous phase with dichloromethane. All 24-hour fecal samples were weighed and homogenized with 1 ml of water. Plasma concentrations of ifosfamide were determined by a validated gas chromatography assay (GC) with nitrogen-phosphorus detection [16]. The analytes were extracted from plasma and urine by liquid-liquid extraction into aqueous phase with ethyl acetate.

Evaluation of c-kit mRNA expression

Total RNA extraction and cDNA synthesis were performed as previously described [17]. Briefly, total RNA was extracted from crushed tumor samples (SCLC 6, SCLC 61, SCLC 74 and SCLC 108) by RNA plus[®] kit (Bioprobe, France). One µg of total RNA was reverse transcribed in a final volume of 20 µL containing 1× reverse transcriptase buffer [500 mM of each deoxynucleotide triphosphate, 3 mM MgCl₂, 75 mM KCl, and 50 mM Tris-HCl (pH 8.3)], 10 units of RNase inhibitor (Promega, Madison, WI), 10 mM DTT, 50 units of Superscript II RNase H-reverse-transcriptase (Life Technology, Inc), and 1.5 mM random hexamers (Pharmacia, Uppsala, Sweden). The reaction mix was then incubated at 42 °C for 30 min.

C-kit transcripts were quantified using real-time quantitative reverse transcription-PCR assays. Primers and probes were chosen with the assistance of Primer Express software (Applied Biosystems, Foster City, CA-IC). The nucleotide sequence was then blasted against dbEST and nr (the non-redundant set of the GenBank, EMBL and DDBJ database sequences) to confirm the total gene specificity of the nucleotide sequences chosen as primers at probes. Each primer couple was positioned in different exons (i.e. exons 20–21 for c-kit gene) of the gene in order to avoid amplification of contaminating genomic DNA. PCR reactions were performed using an ABI Prism 7700 Sequence Detection System (Applied Biosystems, Foster City, CA-IC) and Core Reagent Kit (Eurogenetec, Belgium). Real-time detection was performed using oligonucleotide

probes containing a fluorescent dye at its 5'-end and a quencher at its 3'-end (for c-kit mRNA quantification).

Fluorescent probes were synthesized by Applied Biosystems, and primers were synthesized by Invitrogen (Paisley, UK). The nucleotide and probe sequences are as follows: c-kit: upper primer 5'-aagcagatttcagagaccac-3', lower primer 5'-gctgccgacagaattgatcc-3' and probe 5'-actc-caactagcaactgcagcccaa-3'. Transcripts of TBP (TATA box-binding protein)[18] were also quantified as endogenous RNA of reference genes to normalize c-kit expression.

The thermal cycling conditions comprised an initial denaturation step at 95 °C for 10 min, then 40 cycles at 95 °C for 15 sec and an annealing temperature depending upon the target. For c-kit quantification, a calibration curve for mRNA expression was generated using serial dilutions of L174 human colon carcinoma cell line mRNA expressing high levels of c-kit mRNA to assess PCR efficiency. All measurements were performed under blind conditions and in duplicate. Two negative "non-template" controls were included in each amplification run.

Results were expressed as N-fold differences in target gene expression relative to the reference gene (either TBP) and the calibrator was called "Ntarget". It was determined as follows: $N_{\text{target}} = E^{(\Delta C_t \text{ sample} - \Delta C_t \text{ calibrator})}$, where the ΔC_t values of the sample and calibrator were determined by subtracting the average Ct values of the target gene from the average value of the reference gene (TBP) and E is the efficiency of PCR measured using the slope of the calibration curve. For c-kit mRNA quantification, the positive control, corresponding to a case of uveal melanoma exhibiting strong c-kit overexpression, was assessed with SCLC samples.

Statistical analysis

Statistical analysis of variance (ANOVA test) was performed to assess the difference between AUC_{0–3 h} (Area Under the concentration-time Curve) of etoposide in the various treatments. A Mann-Whitney U-test was used to assess the *in vivo* effect of the various treatments on the growth of xenografted tumors in *nude* or SCID mice. U values were considered significant when the probability of a difference was less than or equal to 0.05. Values were considered significant when the probability of a difference was less than or equal to 0.05.

Results

Impact of STI571 on the *in vivo* xenografted tumor growth of etoposide (VP16) chemotherapy

SCLC6-bearing *nude* mice received intraperitoneal injections of etoposide (12 mg/kg) on days 1–3, with or without STI571 administered on days 1–3 by intraperitoneal

injections at a total dose of 70 mg/kg daily, and with or without intraperitoneal injections of fluconazole (40 mg/kg/d) on days 1–3. Each treatment was initiated on day +6 after tumor transplantation. We observed a brief but non-significant inhibition of SCLC6 tumor growth after

etoposide administration compared to the control group ($p = NS$), and this inhibition was not increased by concomitant injections of etoposide + STI571 or etoposide + STI571 + fluconazole (Figure 1A).




Figure 1

Antitumor activity of imatinib with or without chemotherapy in two human xenografted tumors. (A)(C). Both xenografted SCLC6 (A) and LY-3 (C) tumors were treated by etoposide (VP16) alone (E) at a dosage of 12 mg/kg in one daily intraperitoneal injection on days 1 to 3 (▲), etoposide (VP16) and STI571 (E + S) administered by one daily intraperitoneal injection at a dose of 70 mg/kg on days 1 to 3 (■), etoposide (VP16) and fluconazole (E + F) at a dosage of 40 mg/kg in one daily intraperitoneal injection on days 1 to 3 (○), or etoposide (VP16) with STI571 and fluconazole (E + S + F)(◇). All other groups included STI571 alone (S)(□), fluconazole alone (F)(△), and STI571 + fluconazole (S + F)(◆) and 0.9% NaCl (●). (B) Mice bearing LY-3 tumors were treated by one (□) daily intraperitoneal injection of STI571 (S) at a dose of 70 mg/kg from day 1 until sacrifice of the animals. (D) Xenografted LY-3 tumors were treated by gemcitabine at a dosage of 60 mg/kg by one weekly intraperitoneal injection, with (G + S)(■) or without (G)(▲) STI571 administered by one daily intraperitoneal injection at a dose of 70 mg/kg from day 1 until sacrifice of the animals. Mice treated by STI571 alone (S) are indicated by (□). All control groups received injections of 0.9% NaCl (Control)(●). Tumor growth was evaluated by measuring the relative tumor volume (RTV), as described in "Materials and Methods". A Mann-Whitney *U* test was used to assess the effects of treatments on xenografted tumor growth.

As previously performed for the SCLC6 tumor (Decaudin IJC), we first evaluated the antitumor effect of STI571 alone in the SCID mice xenografted LY-3 lymphomatous tumors. Mice bearing tumors measuring approximately 60 mm³ were treated from day 1 until sacrifice of the animals by an intraperitoneal injection of STI571 at a dose of 70 mg/kg. A slight but non-significant tumor growth inhibition was temporarily observed for a few days after initiation of treatment (Figure 1B). Thereafter, SCID mice bearing LY-3 tumors were treated by intraperitoneal injections of etoposide (12 mg/kg) on days 1–3, with or without intraperitoneal injection of STI571 at a dose of 70 mg/kg on days 1–3, and with or without intraperitoneal injections of fluconazole (40 mg/kg/d) on days 1–3. Each treatment was initiated at day + 6 after tumor transplantation. We observed a spectacular enhancement of etoposide-induced tumor growth inhibition by concomitant administration of STI571 ($p < 0.05$), and this inhibition was not enhanced by concomitant administration of fluconazole (Figure 1C). Moreover, we observed three early deaths in the combined etoposide + STI571 group, but not in the etoposide + STI571 + fluconazole group.

Finally, in order to confirm the value of combining STI571 with other chemotherapy modalities, mice bearing LY-3 tumors measuring approximately 60 mm³ received weekly intraperitoneal injections of gemcitabine (60 mg/kg), with or without intraperitoneal injections of STI571 at a total dose of 70 mg/kg daily from day 1 until sacrifice of the animals. We observed an enhancement of gemcitabine-induced tumor growth inhibition by concomitant and continuous administration of STI571 ($p < 0.01$)(Figure 1D).

Impact of STI571 on the pharmacokinetics of combined etoposide (VP16) and ifosfamide chemotherapy regimen in mice

To study the impact of STI571 on the *in vivo* pharmacokinetics of chemotherapy, C57 Black6 mice were treated on days 1 to 3 by concomitant intraperitoneal injections of etoposide (12 mg/kg/d) and ifosfamide (90 mg/kg/d), with or without STI571 (70 mg/kg/d), and serum was collected from an eye vein on days 1 and 3, 0, 1, 3 and 6 hours after injections. We observed an increase of the Cmax of both etoposide and ifosfamide on days 1 and 3 when mice were treated by chemotherapy combined with STI571 compared to chemotherapy alone (Table 1). To confirm this result and to focus on etoposide pharmacokinetics, CD1 mice were treated on days 1–3 by intraperitoneal injections of 12 mg/kg/d etoposide, with or without intraperitoneal 70 mg/kg/d of STI571. Samples were collected on days 1 and 3, 0.5, 1, 2 and 3 hours after injections. We observed a significant increase of the mean AUC_{0–3 h} (Area Under the concentration-time Curve) of etoposide (17900 µg/l × h) when mice were treated with

Table 1: Mean Cmax of etoposide (VP16) and ifosfamide administered alone or with imatinib

	<i>Ifosfamide</i>	<i>Ifosfamide + imatinib</i>
Mean Cmax	9 +/- 4.5 µg/ml	14.4 +/- 6.4 µg/ml
	Etoposide	Etoposide + imatinib
Mean Cmax	8.6 +/- 4.1 µg/ml	42.7 +/- 37.6 µg/ml

etoposide + STI571 (60800 µg/l × h)($p = 0.025$)(Figure 2A).

In order to evaluate the role of CYP3A4 on the pharmacokinetic interaction between etoposide and STI571, CD1 mice were treated on days 1–3 by intraperitoneal injections of etoposide (12 mg/kg/d), with or without STI571 (70 mg/kg/d), and with or without the CYP3A4 inhibitor, fluconazole (40 mg/kg/d). The samples were collected on days 1 and 3, 0.5, 1, 2 and 3 hours after injections. A summary of etoposide pharmacokinetic parameters (mean) is




Figure 2
(A) Mean AUC_{0–3 h} of etoposide (VP16) in CD1 mice. Area Under Curve concentration versus time of etoposide (VP16) was calculated between base time (0) and 3 hours after administration of etoposide alone or etoposide plus fluconazole or etoposide plus imatinib or etoposide with imatinib and fluconazole. **(B) Concentration-time curve of etoposide (VP16) administered with STI571.** After administration of etoposide (VP16) with STI571, concentrations of VP16 peaked at 30.9 ± 2.1 ng/mL, followed by a bi-exponential decline. **(C) Graphical analysis between calculated concentrations and observed concentrations of etoposide (VP16).**

Table 2: Pharmacokinetic parameters of Etoposide (VP16) administered alone or in combination

	Etoposide	Etoposide + fluconazole	Etoposide + imatinib	Etoposide + fluconazole + imatinib
T 1/2 (hours)	0.24	0.23	0.35	0.49
AUC ($\mu\text{g/L} \times \text{h}$)	19970	23680	38380	48960
Clearance (ml/h)	0.62	0.51	0.32	0.25

presented in Table 2. We observed a highly significant increase of the mean $\text{AUC}_{0-3\text{h}}$ of etoposide ($19970 \mu\text{g/l} \times \text{h}$) when mice were treated with etoposide + STI571 + fluconazole ($48960 \mu\text{g/l} \times \text{h}$), as compared to either etoposide + STI571 ($38380 \mu\text{g/l} \times \text{h}$) or etoposide + fluconazole ($23680 \mu\text{g/l} \times \text{h}$) combinations ($p = 0.0003$) (Figure 2A). In both experiments, after administration of VP16 with STI571, VP16 concentrations peaked at $30.9 \pm 2.1 \text{ ng/mL}$, followed by a bi-exponential decline (Figure 2B). VP16 concentrations reached $46.5 \pm 6.4 \text{ ng/mL}$ when VP16 was administered with STI571 and fluconazole. In both cases, the pharmacokinetic curve was best described as bi-exponential as demonstrated by the concentration-time curve of VP16 administered with STI571 (Figure 2C).

To explain the impact of STI571 on the pharmacokinetics of etoposide, we studied the liver and kidney clearance of etoposide by assaying the chemotherapeutic agent in urine and feces of CD1 mice treated on day 1 by one intraperitoneal injection of etoposide (12 mg/kg/d) with or without STI571 (70 mg/kg/d). Samples were collected for 24 hours after injection. We observed a significant decrease of fecal excretion of etoposide when mice were treated with either etoposide + STI571 ($p = 0.005$) (Figure 3A) or etoposide + STI571 + Fluconazole ($p = 0.003$) (Figure 3B). In contrast, no significant impact of STI571 was observed on the renal clearance of etoposide, with (Figure 3C) or without fluconazole (Figure 3D).

Evaluation of c-kit mRNA expressions

The level of c-kit mRNA expression was quantified and compared to the level of mRNA expression of a known c-kit protein overexpressing melanoma. As previously reported [4], the two SCLC6 and LY-3 xenografts express very low levels of c-kit mRNA (data not shown).

Discussion

In conclusion, we have shown that imatinib increased the AUC of etoposide in mice and that this effect was mediated by a reduction of its fecal excretion. In contrast, imatinib did not appear to influence the urinary excretion of etoposide, probably due to sample collection conditions, and concomitant administration of the CYP3A4 inhibitor fluconazole with imatinib did not modify the pharmacokinetics of etoposide plus imatinib alone. We also demonstrated a slight tumor growth inhibition induced by imatinib administered alone in one *in vivo* EBV-associated

B-cell lymphomatous xenograft, and this effect was observed despite the absence of c-kit receptor mRNA expression. Finally, when mice bearing human xenografted tumors were treated concomitantly by imatinib and chemotherapy, an increase of the chemotherapy-induced antitumor effect was observed in the lymphoma model but not in a small cell lung cancer model, and this antitumor effect was not influenced by concomitant administration of fluconazole.

Our *in vivo* B-cell lymphoma studied by RT-PCR was negative for c-kit mRNA expression. This result was concordant with that reported in the literature, where c-kit expression, studied by RT-PCR or immunohistochemical methods, was mainly observed in multiple myeloma, CD30-positive anaplastic large cell lymphoma, and Hodgkin's disease, as shown in Table 3 [19-31]. Moreover, very few data have been published on the therapeutic efficacy of imatinib on lymphoproliferative diseases. Imatinib demonstrated a minimal effect in anaplastic large cell lymphoma with NPM-ALK fusion protein [32] and Hodgkin lymphoma [33] cell lines. In contrast, despite an *in vitro* antiproliferative effect of STI571 on a multiple myeloma cell line [34], no response was observed in a phase II trial of imatinib in patients with refractory/relapsed myeloma [35].

The combination of imatinib and chemotherapy has not yet been reported in the literature. However, despite a large number of *in vitro* studies, very few data are available on *in vivo* effect of combined imatinib and chemotherapeutic agents. These reports evaluated combinations of imatinib and zoledronate [36], troxacitabine [37], epothilone B [38], and paclitaxel [39]. In our study, we showed that the imatinib-induced increase of etoposide AUC enhances the tumor growth inhibition of a human xenografted lymphomatous tumor, but not that of an *in vivo* human small cell lung cancer tumor. This observation argues in favor of an increased *in vivo* efficacy of etoposide combined with imatinib, showing that chemotherapy resistance could be reversed by the combination with imatinib in some cancer situations, such as our EBV-associated B-cell lymphomatous model and several other small cell lung cancer xenografts previously reported [4].

A number of published clinical studies have used various chemotherapeutic regimens in various cancer indications.




Figure 3
Fecal and urine excretion of etoposide in CDI mice. Etoposide (VP16) levels were determined in feces (ng of etoposide/g of feces) when mice were treated with etoposide alone (A, B) or in combination with imatinib (B) or in combination with imatinib and fluconazole (B). Etoposide (VP16) levels were determined in urine (ng of etoposide/24 hours of urine) when mice were treated with etoposide alone (C, D) or in combination with imatinib (C) or in combination with imatinib and fluconazole (D).

All these data are summarized in Table 4[40-50]. Because of the heterogeneity of these series, it is impossible to clearly define the impact of these combinations on patient outcome. However, they provide a good overview of the toxicities. Some observed side effects can be mainly attributed to the use of antineoplastic drugs, namely hematologic toxicity, nausea and vomiting, infections, and diarrhea, while other side effects could be attributed to the use of imatinib, such as fatigue, peripheral edema, skin eruptions, neuropathy, and liver dysfunction. Only one phase I study of imatinib mesylate combined with doxo-

rubin and gemcitabine in patients with small cell lung carcinoma showed a high rate of dose-limiting toxicity that required early discontinuation of the trials [47].

Conclusion

In summary, imatinib potentiates the effects of etoposide in cancer cells such as lymphoma cell lines. These results therefore show that further prospective phase I and II clinical trials with combinations of etoposide-based chemotherapy and imatinib are therefore warranted in some

Table 3: c-kit expression in lymphoid malignancies

Type of lymphoma	Methods	Positive c-kit expression/N (%)	References
Multiple myeloma	Flow cytometry	49/158 (31%)	Kraj 2004
Multiple myeloma	Flow cytometry	17/48 (35%)	Li 2004
Multiple myeloma	Flow cytometry	18/56 (32%)	Ocqueteau 1996
Multiple myeloma	Immunohistochemistry	5/31 (16%)	Potti 2002
Multiple myeloma	Immunohistochemistry	2/72 (3%)	Lugli 2004
Cutaneous plasmacytoma	Immunohistochemistry	13/13 (100%)	Bayer-Garner 2003
Total MM	/	104/378 (27%)	/
Lymphoplasmacytic	Immunohistochemistry	0/10 (0%)	Lugli 2004
Lymphoplasmacytic	Flow cytometry	0/7 (0%)	Kraj 2004
Mantle cell L.	Immunohistochemistry	2/17 (1%)	Potti 2002
Lymphomatoid papulosis	Immunohistochemistry	0/18 (0%)	Rassidakis 2004
DLBCL	Immunohistochemistry	24/65 (37%)	Vakiani 2005
Total B-cell L.	/	26/117 (22%)	/
CD30+ anaplastic large cell L.	Immunohistochemistry	7/18 (39%)	Brauns 2004
CD30+ anaplastic large cell L.	Immunohistochemistry	11/16 (69%)	Pinto 1994
CD30+ anaplastic large cell L.	Immunohistochemistry	1/78 (1%)	Rassidakis 2004
Lymphomatoid papulosis	Immunohistochemistry	0/18 (0%)	Rassidakis 2004
Total CD30+ L.	/	19/130 (15%)	/
Nasal NK/T-cell L.	Immunohistochemistry	0/36 (0%)	Li 2006
Cutaneous T-cell L.	Immunohistochemistry	2/8 (25%)	Brauns 2004
Mycosis fungoides	Immunohistochemistry	6/18 (33%)	Brauns 2004
Sezary's syndrome	Immunohistochemistry	3/5 (60%)	Brauns 2004
Total T-cell L.	/	11/67 (16%)	/
Hodgkin's disease	Immunohistochemistry	11/21 (52%)	Pinto 1994
Hodgkin's disease	Immunohistochemistry	0/87 (0%)	Rassidakis 2004
Hodgkin's disease	Immunohistochemistry	0/342 (0%)	Zimpfer 2004
Total HD	/	11/450 (2%)	/

Abbreviations: L., lymphoma; MM, multiple myeloma; DLBCL, diffuse large B-cell lymphoma; HD, Hodgkin's disease.

Table 4: Toxicities of imatinib combined with chemotherapy in human cancers

References	N	Treatment protocol	Main toxicities
40	30	STI 400 mg/d + AraC 20 mg/m ² /d ₁₅₋₂₈ .	* Hematologic. * Nausea-vomiting. * Fatigue. * Abdominal pain. * Edema.
41	22	STI 300–800 mg/d + anagrelide 0.5–4 mg/d.	* Hematologic. * Peripheral edema. * Fatigue. * Skin rash.
42	28	STI 600 mg/d + docetaxel 20–45 mg/m ² /d _{1,8,15,22} .	* Hematologic. * Nausea. * Fatigue.
43	20	STI 400 mg/d + hyper-CVAD.	* Hematologic. * Infections.
44	30	STI 400 mg/d + hydroxyurea 1 g/d.	* Edema. * Abdominal pain.
45	20	STI 400/600 mg/d + daunorubicin 50 mg/m ² /d ₁₋₃ + vincristine 2 mg d _{1,8,15,22} + prednisolone 60 mg/m ² /d ₁₋₂₈ + L-asparaginase 4000 U/m ² /d ₁₇₋₂₈ .	* Hematologic. * Hyperbilirubinemia.
46	33	STI 400/500 mg/d + hydroxyurea 1 g/d.	* Hematologic.
47	9	STI 300–400 mg/d + doxorubicin 50–60 mg/m ² /d ₁ .	* Hematologic. * Nausea-vomiting. * Reduction of LVEF. * Fatigue.
47	7	STI 300–400 mg/d + gemcitabine 700–800 mg/m ² /d _{1,8,(15)} .	* Hematologic. * Fatigue.
48	9	* STI 300–600 mg/d + irinotecan 65 mg/m ² d _{1,8} + cisplatin 30 mg/m ² d _{1,8} . * STI 300–800 mg/d + irinotecan 60 mg/m ² d _{1,8,15} + cisplatin 60 mg/m ² d ₁ .	* Hematologic. * Diarrhea. * Thrombosis. * Electrolyte disorder.
49	31	STI 800 mg/d + vincristine 2 mg d _{1,8,15,22} + dexamethasone 40 mg d _{1,2,8,9,15,16,22,23} .	* Hematologic. * Infections. * Neuropathy.
50	80	STI 600 mg/d + cyclophosphamide 1200 mg/m ² /d ₁ + daunorubicin 50 mg/m ² /d ₁₋₃ + vincristine 1.3 mg/m ² /d _{1,8,15,22} + prednisolone 60 mg/m ² /d ₁₋₂₁ .	* Hematologic. * Nausea. * Liver dysfunction.

Abbreviations: Hyper-CVAD, cyclophosphamide + vincristine + doxorubicin + dexamethasone + methotrexate + cytarabine; LVEF, left ventricular ejection fraction.

cancer patients, such as malignant lymphoma patients, with careful toxicologic monitoring.

Authors' contributions

KR, FL, and SW contributed to the pharmacokinetics experiments. IG, VB, RE, and MG contributed to the *in vivo* experiments. PdC contributed to the c-Kit mRNA expression determination. MFP contributed to the correction of the manuscript. DD and FL contributed to the direction of the experiments. DD conceived of the study. All authors read and approved the final manuscript.

Acknowledgements

We are grateful to Novartis for imatinib gift.

References

- Druker BJ, Tamura S, Buchdunger E, Ohno S, Segal GM, Fanning S, Zimmermann J, Lydon NB: **Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of BCR-Abl positive cells.** *Nat Med* 1996, **2**:561-566.
- Carroll M, Ohno-Jones S, Tamura S, Buchdunger E, Zimmermann J, Lydon NB, Gilliland DG, Druker BJ: **CGP 57148, a tyrosine kinase inhibitor, inhibits the growth of cells expressing BCR-ABL, TEL-ABL, and TEL-PDGFR fusion proteins.** *Blood* 1997, **90**(12):4947-4952.
- Dewar AL, Cambareri AC, Zannettino AC, Miller BL, Doherty KV, Hughes TP, Lyons AB: **Macrophage colony-stimulating factor receptor c-fms is a novel target of imatinib.** *Blood* 2005, **105**:3127-3132.
- Decaudin D, de Cremoux P, Sastre X, Judde JG, Némati F, Tran-Perennou C, Fréneaux P, Livartowski A, Pouillart P, Poupon M-F: **In vivo efficacy of STI571 in xenografted human small cell lung cancer alone or combined with chemothe.** *Int J Cancer* 2005, **113**:849-856.
- Pietras K, Ostman A, Sjoquist M, Buchdunger E, Reed RK, Heldin CH, Rubin K: **Inhibition of platelet-derived growth factor receptor reduces interstitial hypertension and increases transcapillary transport in tumors.** *Cancer Res* 2001, **61**:2929-2934.
- Miyamoto T, Sasaguri Y, Sugama K, Azakami S, Morimatsu M: **Expression of the c-kit mRNA in human aortic endothelial cells.** *Biochem Mol Biol Int* 1994, **34**:513-520.
- Langley RR, Fan D, Tsan RZ, Rebhun R, He J, Kim SJ, Fidler IJ: **Activation of the platelet-derived growth factor-receptor enhances survival of murine bone endothelial cells.** *Cancer Res* 2004, **64**:3727-3730.
- Matsui J, Wakabayashi T, Asada M, Yoshimatsu K, Okada M: **Stem cell factor/c-kit signaling promotes the survival, migration, and capillary tube formation of human umbilical vein endothelial cells.** *J Biol Chem* 2004, **279**:18600-18607.
- Platzer P, Herzog W, Thalhammer T, Hamilton G, Haberl I, Jager W: **Formation of a novel topotecan metabolite in the hormone-independent human prostate carcinoma cell lines DU-145 and PC-3.** *Anticancer Res* 1998, **18**:2737-2741.
- Kerbusch T, Jansen RL, Mathot RA, Huitema AD, Jansen M, van Rijswijk RE, Beijnen JH: **Modulation of the cytochrome P450-mediated metabolism of ifosfamide by ketoconazole and rifampin.** *Clin Pharmacol Ther* 2001, **70**:132-141.
- Deininger MWN, O'Brien SG, Ford JM, Druker BJ: **Practical management of patients with chronic myeloid leukemia receiving imatinib.** *J Clin Oncol* 2003, **21**:1637-1647.
- Toffoli G, Corona G, Basso B, Boiocchi M: **Pharmacokinetic optimisation of treatment with oral etoposide.** *Clin Pharmacokinet* 2004, **43**:441-466.
- Decaudin D, Baran Marszak F, Némati F, Mathiot C, Martin A, Némati F, Lantz O, di Santo J, Arnaud P, Bordier V, Vincent-Salomon A, Poupon MF: **High efficacy of combined rituximab and gemcitabine on EBV-associated human B-cell lymphoma obtained after Hodgkin's xenograft in immunodeficient mice.** *Anti-Cancer drugs* 2006, **17**:685-695.
- Poupon M-F, Arvelo F, Goguel AF, Bourgeois Y, Jacrot M, Hanania N: **Response of small-cell lung cancer xenografts to chemotherapy: multidrug resistance and direct clinical correlates.** *J Natl Cancer Inst* 1993, **85**:2023-2029.
- Harvey VJ, Joel SP, Johnston A, Slevin ML: **High performance liquid chromatography of etoposide in plasma and urine.** *J Chromatogr* 1985, **339**:419-423.
- de Bruijn EA, Slee PH, Van Oosterom AT, Lameijer DW, Roozendaal KJ, Tjaden UR: **Pharmacokinetics of intravenous and oral cyclophosphamide in the presence of methotrexate and fluorouracil.** *Pharm Weekbl Sci* 1988, **10**:200-206.
- de Cremoux P, Tran-Perennou C, Brockdorff B, Boudou E, Brunner N, Magdelenat H, Lykkesfeldt AE: **Validation of real-time RT-PCR for analysis of human breast cancer cell lines resistant or sensitive to treatment with antiestrogens.** *Endocr Relat Cancer* 2003, **10**:409-184.
- Spyratos F, Bouchet C, Tozlu S, Labroquere M, Vignaud S, Becette V, Lidereau R, Bieche I: **Prognostic value of uPA, PAI-1 and PAI-2 mRNA expression in primary breast cancer.** *Anticancer Res* 2002, **22**:2997-3003.
- Pinto A, Gloghini A, Gattei V, Aldinucci D, Zagonel C, Carbone A: **Expression of the c-kit receptor in human lymphomas is restricted to Hodgkin's disease and CD30+ anaplastic large cell lymphomas.** *Blood* 1994, **83**:785-792.
- Ocqueteau M, Orfao A, Garcia-Sanz R, Almeida J, Gonzalez M, San Miguel JF: **Expression of the CD117 antigen (c-Kit) on normal and myelomatous plasma cells.** *Br J Haematol* 1996, **95**:489-493.
- Potti A, Ganti AK, Kargas S, Koch M: **Immunohistochemical detection of C-kit (CD117) and vascular endothelial growth factor (VEGF) overexpression in mantle cell lymphoma.** *Anticancer Res* 2002, **22**:2899-2901.
- Potti A, Ganti AK, Koch M, Levitt R, Mehdi SA: **Immunohistochemical identification of HER-2/neu overexpression and CD117 (c-kit) expression in multiple myeloma.** *Leuk Lymphoma* 2002, **43**:2427-2430.
- Bayer-Garner IB, Schwartz MR, Lin P, Smoller BR: **CD117, but not lysozyme, is positive in cutaneous plasmacytoma.** *Arch Pathol Lab Med* 2003, **127**:1596-1598.
- Brauns TC, Schultewolter T, Dissemmond J, Maschke J, Goos M: **C-KIT expression in primary cutaneous T-cell lymphomas. C-KIT expression in primary cutaneous T-cell lymphomas.** *J Cutan Pathol* 2004, **31**:577-582.
- Kraj M, Poglod R, Kopec-Szlezak J, Sokolowska U, Wozniak J, Kruk B: **C-kit receptor (CD117) expression on plasma cells in monoclonal gammopathies.** *Leuk Lymphoma* 2004, **45**:2281-2289.
- Li J, Luo SK, Zhang GC, Hong WD, Tong XZ: **Expression of CD117 antigen on multiple myeloma and its significance.** *Ai Zheng* 2004, **23**:951-954.
- Lugli A, Went P, Khanlari B, Nikolova Z, Dirnhofer S: **Rare KIT (CD117) expression in multiple myeloma abrogates the usefulness of imatinib mesylate treatment.** *Virchows Arch* 2004, **444**:264-268.
- Rassidakis GZ, Georgakis GV, Oyarzo M, Younes A, Medeiros LJ: **Lack of c-kit (CD117) expression in CD30+ lymphomas and lymphomatoid papulosis.** *Mod Pathol* 2004, **17**:946-953.
- Zimpfer A, Went P, Tzankov A, Pehrs AC, Lugli A, Maurer R, Terracciano L, Pileri S, Dirnhofer S: **Rare expression of KIT (CD117) in lymphomas: a tissue microarray study of 1166 cases.** *Histopathology* 2004, **45**:398-404.
- Vaniaki E, Cattoretti G, Colovai AI, Murty VV, Alobeid B, Bhagat G: **CD117 expression in diffuse large B-cell lymphomas: fact or fiction?** *Pathol Int* 2005, **55**:716-723.
- Li T, Zhang B, Ye Y, Yin H: **Immunohistochemical and genetic analysis of Chinese nasal natural killer/T-cell lymphomas.** *Hum Pathol* 2006, **37**:54-60.
- Ergin M, Denning MF, Izban KF, Amin HM, Martinez RL, Saeed S, Alkan S: **Inhibition of tyrosine kinase activity induces caspase-dependent apoptosis in anaplastic large cell lymphoma with NPM-ALK (p80) fusion protein.** *Exp Hematol* 2001, **29**:1082-1090.
- Re D, Wickenhauser C, Ahmadi T, Buchdunger E, Kochanek M, Diehl V, Wolf J: **Preclinical evaluation of the antiproliferative potential of STI571 in Hodgkin's disease.** *Br J Cancer* 2002, **86**:1333-1335.
- Pandiella A, Carvajal-Vergara X, Tabera S, Mateo G, Gutierrez N, San Miguel JF: **Imatinib mesylate (STI571) inhibits multiple myeloma cell proliferation and potentiates the effect of common antimyeloma agents.** *Br J Haematol* 2003, **123**:858-888.

35. Dispenzieri A, Gertz MA, Lacy MQ, Geyer SM, Greipp PR, Rajkumar SV, Kimminger T, Lust JA, Fonseca R, Allred J, Witzig TE: **A phase II trial of imatinib in patients with refractory/relapsed myeloma.** *Leuk Lymphoma* 2006, **47**:39-42.
36. Kuroda J, Kimura S, Segawa H, Kobayashi Y, Yoshikawa T, Urasaki Y, Ueda T, Enjo F, Tokuda H, Ottmann OG, Maekawa T: **The third-generation bisphosphonate zoledronate synergistically augments the anti-Ph+ leukemia activity of imatinib mesylate.** *Blood* 2003, **102**:2229-2235.
37. Orsolic N, Giles FJ, Gourdeau H, Golemovic M, Beran M, Cortes J, Freireich EJ, Kantarjian H, Verstovsek S: **Troxacitabine and imatinib mesylate combination therapy of chronic myeloid leukaemia: preclinical evaluation.** *Br J Haematol* 2004, **124**:727-738.
38. O'Reilly T, Wartmann M, Maira SM, Hattenberger M, Vaxelaire J, Muller M, Ferretti S, Buchdunger E, Altmann KH, McSheehy PM: **Patupilone (epothilone B, EPO906) and imatinib (ST1571, Glivec) in combination display enhanced antitumour activity in vivo against experimental rat C6 glioma.** *Cancer Chemother Pharmacol* 2005, **55**:307-317.
39. Corcoran NM, Costello AJ: **Combined low-dose imatinib mesylate and paclitaxel lack synergy in an experimental model of extra-osseous hormone-refractory prostate cancer.** *BJU Int* 2005, **96**(4):640-646.
40. Gardembas M, Rousselot P, Tulliez M, Vigier M, Buzyn A, Rigal-Huguet F, Legros L, Michallet M, Berthou C, Cheron N, Maloisel F, Mahon FX, Facon T, Berthaud P, Guilhot J, Guilhot F, CML French Group: **Results of a prospective phase 2 study combining imatinib mesylate and cytarabine for the treatment of Philadelphia-positive patients with chronic myelogenous leukemia in chronic phase.** *Blood* 2003, **102**:4298-4305.
41. Tsimberidou AM, Colburn DE, Welch MA, Cortes JE, Verstovsek S, O'Brien SM, Albitar M, Kantarjian HM, Giles FJ: **Anagrelide and imatinib mesylate combination therapy in patients with chronic myeloproliferative disorders.** *Cancer Chemother Pharmacol* 2003, **52**:229-234.
42. Mathew P, Thall PF, Jones D, Perez C, Bucana C, Troncso P, Kim SJ, Fidler IJ, Logothetis C: **Platelet-derived growth factor receptor inhibitor imatinib mesylate and docetaxel: a modular phase I trial in androgen-independent prostate cancer.** *J Clin Oncol* 2004, **22**:3323-3329.
43. Thomas DA, Faderl S, Cortes J, O'Brien S, Giles FJ, Kornblau SM, Garcia-Manero G, Keating MJ, Andreeff M, Jeha S, Beran M, Verstovsek S, Pierce S, Letvak L, Salvado A, Champlin R, Talpaz M, Kantarjian H: **Treatment of Philadelphia chromosome-positive acute lymphocytic leukemia with hyper-CVAD and imatinib mesylate.** *Blood* 2004, **103**:4396-4407.
44. Dresemann G: **Imatinib and hydroxyurea in pretreated progressive glioblastoma multiforme: a patient series.** *Ann Oncol* 2005, **16**:1702-1708.
45. Lee KH, Lee JH, Choi SJ, Lee JH, Seol M, Lee YS, Kim WK, Lee JS, Seo EJ, Jang S, Park CJ, Chi HS: **Clinical effect of imatinib added to intensive combination chemotherapy for newly diagnosed Philadelphia chromosome-positive acute lymphoblastic leukemia.** *Leukemia* 2005, **19**:1509-1516.
46. Reardon DA, Egorin MJ, Quinn JA, Rich JN, Gururangan S, Vredenburg JJ, Desjardins A, Sathornsumetee S, Provenzale JM, Herndon JE 2nd, Dowell JM, Badruddoja MA, McLendon RE, Lagattuta TF, Kicielewski KP, Dresemann G, Sampson JH, Friedman AH, Salvado AJ, Friedman HS: **Phase II study of imatinib mesylate plus hydroxyurea in adults with recurrent glioblastoma multiforme.** *J Clin Oncol* 2005, **23**:9359-9368.
47. George S, Desai J, Paul Eder J, Manola J, Ryan DP, Appleman LJ, Demetri GD: **Selective kinase inhibition with daily imatinib intensifies toxicity of chemotherapy in patients with solid tumours.** *Eur J Cancer* 2006, **42**:864-870.
48. Johnson FM, Krug LM, Tran HT, Shoaf S, Prieto VG, Tamboli P, Peeples B, Patel J, Glisson BS: **Phase I studies of imatinib mesylate combined with cisplatin and irinotecan in patients with small cell lung carcinoma.** *Cancer* 2006, **106**:366-374.
49. Rea D, Legros L, Raffoux E, Thomas X, Turlure P, Maury S, Dupriez B, Pigneux A, Choufi B, Reman O, Stephane D, Royer B, Vigier M, Ojeda-Urbe M, Recher C, Dombret H, Huguier F, Rousselot P, Intergroupe Francais des Leucemies Myeloides Chronique; Group for Research in Adult Acute Lymphoblastic Leukemia: **High-dose imatinib mesylate combined with vincristine and dexamethasone (DIV regimen) as induction therapy in patients with resistant Philadelphia-positive acute lymphoblastic leukemia and lymphoid blast crisis of chronic myeloid leukemia.** *Leukemia* 2006, **20**:400-403.
50. Yanada M, Takeuchi J, Sugiura I, Akiyama H, Usui N, Yagasaki F, Kobayashi T, Ueda Y, Takeuchi M, Miyawaki S, Maruta A, Emi N, Miyazaki Y, Ohtake S, Jinnai I, Matsuo K, Naoe T, Ohno R, Japan Adult Leukemia Study Group: **High complete remission rate and promising outcome by combination of imatinib and chemotherapy for newly diagnosed BCR-ABL-positive acute lymphoblastic leukemia: a phase II study by the Japan Adult Leukemia Study Group.** *J Clin Oncol* 2006, **24**:460-466.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

