

Table 2. Distribution of Microorganisms in Positive Sputum Cultures

| True pathogens | | | |
|---|----------|--|---|
| VAP N= 48 | HAP N=11 | | |
| <i>Staphylococcus aureus</i> (MSSA) | 21 | <i>Staphylococcus aureus</i> (MSSA) | 2 |
| <i>Pseudomonas aeruginosa</i> | 8 | <i>Pseudomonas aeruginosa</i> | 2 |
| <i>Klebsiella (enterobacter) aerogenes</i> | 6 | <i>Staphylococcus aureus</i> (MRSA) | 2 |
| <i>Klebsiella pneumoniae</i> | 5 | <i>Aspergillus fumigatus</i> | 2 |
| <i>Stenotrophomonas (Xanthomonas) maltophilia</i> | 3 | <i>Klebsiella pneumoniae</i> | 1 |
| <i>Klebsiella pneumoniae MDR</i> | 2 | <i>Klebsiella (enterobacter) aerogenes</i> | 1 |
| <i>Staphylococcus aureus</i> (MRSA) | 2 | <i>Candida albicans</i> | 1 |
| <i>Candida tropicalis</i> | 2 | | |
| <i>Escherichia coli</i> | 2 | | |
| <i>Candida albicans</i> | 1 | | |
| <i>Streptococcus agalactiae</i> (group B) beta hemolytic | 1 | | |
| <i>Streptococci</i> (group C) beta hemolytic | 1 | | |
| <i>Burkholderia cepacia</i> complex | 1 | | |
| <i>Acinetobacter baumannii</i> complex | 1 | | |
| <i>Proteus mirabilis</i> | 1 | | |
| <i>Klebsiella oxytoca</i> MDR | 1 | | |
| <i>Escherichia coli</i> MDR | 1 | | |
| <i>Citrobacter farmeri</i> MDR | 1 | | |
| <i>Citrobacter koseri</i> (<i>Citrobacter diversus</i>) | 1 | | |
| <i>Candida dubliniensis</i> | 1 | | |
| <i>Candida parapsilosis</i> | 1 | | |
| <i>Streptococcus pneumoniae</i> | 1 | | |
| <i>Pseudomonas putida</i> | 1 | | |
| Colonization | | | |
| <i>Candida albicans</i> | 34 | | |
| <i>Candida tropicalis</i> | 6 | | |
| <i>Candida parapsilosis</i> | 5 | | |
| <i>Candida krusei</i> | 1 | | |
| <i>Candida dubliniensis</i> | 1 | | |

MDR: Multidrug resistant; MRSA: Methicillin-resistant *Staphylococcus aureus*; MSSA: Methicillin-sensitive *Staphylococcus aureus*

Conclusion: Among high risk COVID-19 patients, NP is a common complication. MSSA and *Enterobacteriaceae* were the most frequent isolates. The risk increases with intubation, longer hospital stay and use of steroids but not tocilizumab.

Disclosures: All Authors: No reported disclosures

383. Increased Need for Antimicrobial Stewardship during a COVID-19 Outbreak in New York City

Maurice Policar, MD¹; Peter Barber, PharmD¹; Yesha Malik, MD¹; ¹Health + Hospitals / Elmhurst, Elmhurst, New York

Session: P-12. COVID-19 Complications, Co-infections, and Clinical Outcomes

Background: The impact of COVID-19 on the health care system in New York City (NYC) cannot be overstated. The first documented cases of COVID-19 in Queens NYC occurred in early March of 2020. The total number of patients with proven or suspected COVID-19 at Elmhurst Hospital peaked in early April. A dramatic increase in the use of antimicrobials occurred in April, and correlated with the increased number of intubated COVID-19 patients at Elmhurst Hospital.

Methods: Antimicrobial Stewardship Committee activities and meetings had been suspended for the months of March and April due to the increased clinical demands associated with the COVID-19 outbreak. In preparation for the May meeting, a retrospective analysis of antimicrobial use for March and April of 2020 was performed.

Results: The analysis revealed a 30% increase in the use of antimicrobials. The average total days of antimicrobials per 1000 patient days (TDA/TPD) was 445 for January through March of 2020. In April, this number climbed to 580. TDA/TPD increased from 57 to 90 (58%) for vancomycin, 25 to 35 (40%) for meropenem, and 31 to 89 (187%) for cefepime. The number of intervention by the Antibiotic Stewardship team remained low during this time period.

Total Days of Antimicrobials per 1000 Patient Days (TDA/TPD)

| ANTIBIOTIC | JANUARY | FEBRUARY | MARCH | Q1 AVERAGE | APRIL | PERCENT INCREASE |
|------------|---------|----------|-------|------------|-------|------------------|
| vancomycin | 63 | 57 | 52 | 57 | 90 | 58 |
| meropenems | 35 | 30 | 27 | 31 | 89 | 40 |
| cefepime | 35 | 30 | 27 | 31 | 89 | 187 |

Conclusion: A dramatic increase in the use of antimicrobials correlated with an increase in the number of intubated patients at Elmhurst Hospital during a COVID-19 outbreak. It is likely that the frequent appearance of fever and leukocytosis in intubated patients with COVID-19 prompted an increase in empiric antimicrobial use. The 48 hour time outs and prospective review of antimicrobial use may be necessary to maintain stewardship efforts during the COVID-19 epidemic. Further review of antibiotic usage in critically ill COVID-19 patients is needed to help define stewardship practices as we go forward in this pandemic.

Disclosures: All Authors: No reported disclosures

384. Invasive aspergillosis in COVID-19 patients in an intensive care unit in Mexico City

Mariana Velez-Pintado, n/a¹; Mercedes Aguilar-Soto, n/a¹; Antonio Camiro, MD, MSc²; Dalia Cuenca-Abruch, n/a¹; Renzo Alberto Pérez-Doramea, n/a¹; Brenda Crabtree-Ramirez, n/a¹; Moises Mercado-Atri, n/a¹; ¹Centro Médico ABC, Mexico City, Distrito Federal, Mexico; ²Centro Médico ABC, Mexico City, Distrito Federal, Mexico

ARMII study group

Session: P-12. COVID-19 Complications, Co-infections, and Clinical Outcomes

Background: An elevated incidence of invasive pulmonary aspergillosis (IPA) in patients with COVID-19 without traditional risk factors for IPA has been recently reported around the world. This co-infection has been described in patients requiring treatment in an intensive care unit. The risk factors for its development are still unclear.

Methods: We conducted a nested case-control study using the COVID-19 registry of the ARMII study group, based in the Centro Médico ABC, a private hospital in Mexico City. We included all patients that required admission to the intensive care unit (ICU) from March 12 to June 15, 2020, and excluded patients without serum galactomannan measurements or bronchial secretion cultures. We used the modified definition of IPA proposed by Schauwvlieghe et al for IPA in influenza patients. The control group was formed by patients with ruled-out IPA (negative galactomannan and secretion cultures). We compared both groups to identify risk factors for IPA using the chi-squared test or the Mann-Whitney U test as applicable.

Results: Out of a total 239 patients, 54 met the inclusion criteria. We identified 13 patients with IPA (24.07%) that met the definition of IPA (2 with positive cultures and 11 with positive galactomannan) and 41 without IPA. Only three patients with IPA had important comorbidities (COPD, chronic kidney disease, and HIV). Patients with IPA tended to have a higher median age (64.6 vs 53.59, p=0.075) and a higher serum glucose at their arrival (145 vs 119, p=0.028). All patients with IPA presented to the hospital with ARDS (100% vs 72.5%, p=0.034), but ultimately did not have a higher requirement for mechanical ventilation (100% vs 82.93%, p=0.110). There were no statistical significant differences in use of Tocilizumab, use of glucocorticoids, mortality (23.07% vs 17.50%, p=0.563) or length of stay.

Conclusion: It has been previously described that patients with acute respiratory disease syndrome triggered by viral infection, like the influenza virus, are prone to invasive aspergillosis even in the absence of underlying immunodeficiency. The use of antifungals to prevent aspergillosis in COVID-19 patients should be assessed because of the gravity presented in the patients with this co-infection.

Disclosures: All Authors: No reported disclosures

385. Kawasaki's Disease and Sars-Cov-2: an Unexpected Pediatric Global Crisis?

Lucca G. Giarola, Medical Student¹; Braulio Roberto Gonçalves Marinho Couto, n/a¹; Carlos Ernesto Ferreira Starling, n/a²; Handerson Dias Duarte de Carvalho, Medical Student³; ¹Centro Universitário de Belo Horizonte, Belo Horizonte, Minas Gerais, Brazil; ²Lifecenter Hospital, Belo Horizonte, Minas Gerais, Brazil; ³Centro Universitário de Belo Horizonte - UniBH, Belo Horizonte, Minas Gerais, Brazil

Session: P-12. COVID-19 Complications, Co-infections, and Clinical Outcomes

Background: Infection by SARS-CoV-2 can lead to dyspnea, edema, deposition of intra alveolar fibrin, thrombosis and hemorrhages. During the COVID-19 outbreak, several questions were raised about the risks for the pediatric population. Pediatric patients appeared to be relatively safe, with only minor symptoms and a quick recovery. However, there have been reports of a relationship between COVID 19 and a Kawasaki-like inflammatory disease in this population. Kawasaki's disease (KD) is a rheumatological vasculitis prevalent in childhood characterized mainly by diffuse inflammation of the arteries associated with skin rash, changes in the mucosa and its main complication is coronary aneurysms.

Methods: A systematic literature review was performed in the PubMed database using the keywords "Kawasaki disease", "COVID-19" and "Pediatrics". The selected filters were "Case reports", "Multicenter study", "Clinical Study", "Observational study", "Human" and "English". A total of 18 articles were selected.

Results: There seems to be a convergence between the literature published so far, pointing to a greater propensity for pediatric patients infected with Sars-Cov-2 to develop KD. The number of patients with KD symptoms seen at a specific center increased from 2 to 17 in 11 days (MOREIRA, 2020). In a sample space of 21 patients diagnosed with KD, 91% had previous contact with SARS-CoV-2 (TOUBIANA, 2020) whereas other studies point to a 30-fold increase in the prevalence of KD since the beginning of 2020 (VERDONI, 2020).

There is already an established relationship between KD and HCoV-NH, describing that 4.5% of patients with this infection develop KD. Therefore, it was suggested that infection with another Coronavirus strain could have a similar relationship.

Conclusion: Despite the relationship described between pediatric patients infected with COVID-19 being more likely to develop KD, further studies are needed to prove a statistical relationship between both condition.

Disclosures: All Authors: No reported disclosures

386. Long-term Complications Associated with COVID-19 Infection

Smitha Gudipati, MD¹; Judith L. Ranger, BSN²; Amit T. Vahia, MD MPH²; Tommy J. Parraga Acosta, MD²; Zachary W. Hanna³; Sashi N. Nair, MD²; Nicholas F. Yared, MD¹; Geehan Suleyman, MD²; Indira Brar, MD²; ¹Henry Ford Health System, Detroit, Michigan; ²Henry Ford Hospital, Harrison Township, Michigan; ³Michigan State University, Berkley, MI

Session: P-12. COVID-19 Complications, Co-infections, and Clinical Outcomes

Background: In Michigan, 44,964 (68%) of the 66,269 COVID-19 patients have recovered. However, there is concern that COVID-19 infection may lead to long-term sequelae, including pulmonary defects, cardiac complications, blood clots, and neurocognitive impairment. This study describes the 30-day outcomes of patients who had recovered.