

Research Article

R-Loop Formation *In Trans* at an AGGAG Repeat

Kazuya Toriumi,^{1,2} Takuma Tsukahara,¹ and Ryo Hanai¹

¹ Department of Life Science and Research Center for Life Science, Rikkyo (St. Paul's) University, 3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan

² Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kami-Kitazawa, Setagaya-ku, Tokyo 156-8506, Japan

Correspondence should be addressed to Ryo Hanai; hanai@rikkyo.ac.jp

Received 29 May 2013; Accepted 25 July 2013

Academic Editor: Luis A. Marky

Copyright © 2013 Kazuya Toriumi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Formation of RNA-DNA hybrid, or R-loop, was studied *in vitro* by transcribing an AGGAG repeat with T7 RNA polymerase. When ribonuclease T1 was present, R-loop formation *in cis* was diminished, indicating that the transcript was separated from the template and reassociated with it. The transcript was found to form an R-loop *in trans* with DNA comprising the AGGAG repeat, when the DNA was supercoiled. Results of chemical modification indicated that the duplex opened at the AGGAG repeat under negative supercoiling.

1. Introduction

The DNA-dependent RNA polymerase separates product RNA from its template DNA, thereby making the RNA available to ensuing cellular processes [1, 2]. The polymerase possesses structural elements for this separation and the RNA transcript is extruded through a hole of the polymerase molecule [3–6]. Despite this “separator” function of the polymerase, some transcripts have been known to anomalously form an RNA:DNA hybrid, or an R-loop, with their template, and R-loops have been implicated in a number of biological processes [7]. One classical example is the *colE1* replication origin, where the RNA in the R-loop serves as the primer of DNA replication [8]. A more recent example is the *FLOWERING LOCUS C* of *Arabidopsis thaliana*, where an R-loop at the promoter of the *COOLER* gene represses the transcription of the gene [9]. In addition to the “separator” function of the RNA polymerase, cellular processes ensuing transcription, such as splicing and RNA export, also serve to sequester RNA from the template DNA. Compromising such a function has been shown to result in hyperrecombination [7, 10].

R-loops have also been observed at G-rich repetitive sequences in immunoglobulin class-switch regions, although R-loop *per se* is not considered to trigger class-switching [11–16]. R-loop formation at a class-switch region was first shown

for the murine S_{α} region [11]. Transcription of supercoiled plasmid DNA containing a 2.3 kb fragment of the region resulted in relaxation of the DNA, and the relaxation was reversed by RNase H treatment, which indicated R-loop formation. The relaxation was dependent on the direction of transcription. Radiolabeling experiments showed that the R-loop formed also when the template was linearized and that the RNA content in the R-loop was solely A and G [12]. The latter result, together with the length of the RNase A-resistant RNA bound to the DNA, indicated that the R-loop formed at the 28 tandem repeats of AGGAG in the S_{α} region.

Despite the simple repetitive nature of the S_{α} sequence, the exact mechanism of the R-loop formation has not been known. In this report, we present experimental results *in vitro* on an AGGAG repeat to address the following two questions. Firstly, it is not known whether the RNA is separated from and reassociate with the template or a gross structural change takes place in the polymerase molecule for the transcript to stay on the template. The latter possibility cannot be ruled out *a priori*, because the R-loop was found to form in the presence of ribonuclease (RNase) A *in vitro* [12]. Secondly, it is not known what structure in the DNA and/or the RNA leads to the R-loop formation: it is not known what allows the transcript to reassociate with its template or what forces the polymerase into an altered conformation in which the transcript can stay on the template.

2. Materials and Methods

2.1. Plasmid Construction. Synthetic oligodeoxynucleotides (GGAGA)₅ and (TCTCC)₅ were used to synthesize (GGAGA)_n · (TCTCC)_n by the use of the nontemplate PCR method [17]. The product was cloned into the *EcoRV* site of pBluescript SK(-) (Stratagene). Nucleotide sequencing showed that one clone contained GAG(AGGAG)₂₂ in the *Kpn* I-to-*Sac* I direction and the clone was designated pSK-(AGGAG)₂₂. A fragment containing the repeat was cut out with *EcoR* I and *Hind* III and cloned in between the *EcoR* I and the *Hind* III sites of pHC624 [18] to make pHC624-(AGGAG)₂₂. Both plasmids were purified from *Escherichia coli* DH5 α by Cesium-Ethidium banding [19].

2.2. Transcription. Standard transcription reaction was carried out with 6 units of T7 RNA polymerase (Roche Diagnostics) on 0.3 μ g DNA (0.3 μ g for each DNA, when two DNA species were present) at 37°C for 20 min in 20 μ L of 50 mM NaCl, 40 mM Tris-HCl pH 8.0, 6 mM MgCl₂, 2 mM spermidine, 10 mM DTT, and 0.6 mM each of ATP, CTP, GTP, and UTP [11, 12]. One μ L of 0.5 M EDTA pH 8.0 was added to terminate transcription and 2.5 μ L of 1 μ g/ μ L RNase A (Sigma) was added [11, 12]. When the transcript was radiolabeled, [α -³²P] ATP was also included. After incubation at 37°C for 1 h, the reaction was extracted with phenol/chloroform/isoamylalcohol. When RNase T1 was included (Figure 1(b)), transcription was carried out at an NaCl concentration of 150 mM for 20 min. One μ L of 0.5 M EDTA pH 8.0 was added and the reactions were incubated at 37°C for 5 min and extracted with phenol/chloroform/isoamylalcohol. Samples were electrophoresed in 1% agarose in 50 mM Tris-borate and 1 mM EDTA, DNA was stained with ethidium bromide, and the gel was photographed on a UV transilluminator or scanned on a Typhoon analyzer (GE Healthcare Life Science). For detection of the radiolabel, the agarose gel was dried on Hybond-N⁺ membrane (GE Healthcare Life Science), the radioactivity was recorded on an Image Plate (Fuji Photo Film), and the Plate was scanned on a Typhoon analyzer.

2.3. Chemical Probing of DNA Structure. Supercoiled pSK-(AGGAG)₂₂ was relaxed with vaccinia topoisomerase I [20]. Five μ g of supercoiled or relaxed DNA was subjected to chemical modification in 200 μ L of 50 mM sodium cacodylate pH 8.0 and 6 mM MgCl₂. Oxidation with potassium permanganate [21] was carried out at 0.9 mM for 2 min at 37°C and terminated by adding 10 μ L of β -mercaptoethanol. Modification with diethylpyrocarbonate [22] was started by adding 3 μ L of the reagent and vortexing, the reaction was incubated at 37°C for 10 min with one more vortexing during incubation, and 10 μ L of β -mercaptoethanol was added to terminate the modification. The DNAs were purified by the use of glass milk (Q-Biogene). Permanganate-oxidized DNA was digested with *Xho* I and dephosphorylated with calf intestine phosphatase. The reaction was extracted with phenol/chloroform/isoamylalcohol, and the DNA purified with glass milk. The DNA was end-labeled with [γ -³²P]




FIGURE 1: R-loop formation on pSK-(AGGAG)₂₂ and effects of RNase T1 on the formation. A plasmid containing a (AGGAG)₂₂ repeat was transcribed with T7 RNA polymerase and the effects on the topology of the plasmid were examined by agarose gel electrophoresis. (a) Lane 1: mock transcribed; lane 2: transcribed with T7 RNA polymerase; lane 3: transcribed with T3 RNA polymerase. (b) Lane 1: mock transcribed, lane 2: transcribed with T7 RNA polymerase, lane 3: transcribed with T7 RNA polymerase in the presence of 3 units of RNase T1; lane 4: transcribed with T7 RNA polymerase and ethanol-precipitated; and lane 5: transcribed with T7 RNA polymerase, ethanol-precipitated, and then incubated with 3 units of RNase T1.

ATP and T4 polynucleotide kinase. After inactivating the kinase by heating at 65°C for 20 min, the DNA was digested with *Xho* I in the presence of 1 mg/mL spermidine [22]. After extraction with phenol/chloroform/isoamylalcohol, the samples were spun in S-400 HR columns (GE Healthcare Life Science) to remove unincorporated radiolabel and the small *Xho* I-*Kpn* I fragment. Diethylpyrocarbonate-treated DNA was similarly processed with *Not* I and *Sac* I. Recovered DNA was ethanol precipitated with 4 μ g of carrier DNA and air dried. The DNA was dissolved in 100 μ L of 1 M piperidine and 2 mM EDTA, heated at 90°C for 30 min, and vacuum-dried. The dried pellet was dissolved in 20 μ L of water and vacuum-dried. This process was repeated once, and the DNA was dissolved in formamide loading solution [19]. Maxam-Gilbert sequencing reactions were carried out by the standard procedures [19]. The samples were electrophoresed in 5% Long Ranger (Lonza)-urea sequencing gel. After washing-off of urea and drying of the gel, the radioactivity was recorded on an image plate and the plate was scanned on a Typhoon analyzer.

3. Results

3.1. Formation of an R-Loop at (AGGAG)₂₂. In order to create a simplified model of S α , an AGGAG repeat was synthesized by the nontemplate PCR method [17] and cloned into

the *EcoR* V site of pBluescript SK(-). One clone contained GAG(AGGAG)₂₂ at the site in the direction that transcription from the T7 promoter would produce an AGGAG-repeat transcript. The plasmid, designated pSK-(AGGAG)₂₂, purified from *E. coli* and negatively supercoiled, was transcribed with T7 RNA polymerase. Agarose gel electrophoresis showed extensive relaxation of the DNA (Figure 1(a), lane 2) indicative of R-loop formation. The relaxation was orientation-dependent (compare with transcription from T3 promoter; lane 3), as reported previously [11, 12]. In addition, when the repeat was moved into pBluescript KS(+) to reverse the orientation, transcription from the T3 promoter but not from the T7 promoter resulted in relaxation of the DNA (data not shown). These results showed that the AGGAG repeat was sufficient for R-loop formation, although the AGGAG repeat in the S_α region is preceded by a tantalizing GAGCT repeat (GenBank/EMBL/DDBJ entry D11468).

3.2. Inhibition of R-Loop Formation by RNase T1. In order to test whether the transcript was separated from and then reassociated with the template or the RNA stayed on the template, we included a ribonuclease (RNase) in the transcription mixture. If the RNA stays on the template, the RNase should not affect the R-loop formation. Although it was reported that inclusion of RNase A did not affect the formation [11, 12], RNase A is known to be specific to pyrimidines and thus would not be expected to digest the AGGAG-repeat RNA. Therefore, RNase T1, which is specific to purines, was chosen. When RNase T1 was present, the relaxation was diminished (Figure 1(b), lane 3). RNase T1 is known to cleave RNA in R-loops at lower salt concentrations. However, the R-loop was stable under the conditions used, as no change was observed when pre-formed R-loop was incubated with RNase T1 (lanes 4 and 5). Thus, the result indicated that the transcript was separated from and reassociated with the template and that the “separator” function of T7 RNA polymerase was not compromised on the CTCCT-repeat template, in agreement with results reported for other class-switch regions and a telomeric repeat [23].

3.3. R-Loop Formation in Trans and Its Dependence on DNA Supercoiling. The above result indicating reassociation suggested that the R-loop might form *in trans*. This possibility was explored by including in the transcription mixture a plasmid that contained the AGGAG repeat but no T7 promoter. The plasmid, pHC624-(AGGAG)₂₂, was constructed by inserting the AGGAG repeat in a small cloning vector, pHC624 [18]. An AGGAG-repeat transcript was produced by transcribing linearized pSK-(AGGAG)₂₂ with T7 RNA polymerase, and effects on the target pHC624-(AGGAG)₂₂ were examined. The result is shown in Figure 2. Change in DNA topology was observed for supercoiled pHC624-(AGGAG)₂₂ (Figure 2, lanes 4 and 5). The change was dependent on the presence of the AGGAG repeat both in the transcribed and the target plasmids (lanes 6–8). The change was reversed by treatment with RNase H (lanes 11 and 12), indicating that the change was due to the formation of an R-loop. No change in topology was observed for relaxed pHC624-(AGGAG)₂₂




FIGURE 2: R-loop formation *in trans* and its dependence on supercoiling of the target DNA. RNA containing an AGGAG repeat was produced from linearized pSK-(AGGAG)₂₂, and the effects on a T7 promoterless plasmid containing an AGGAG repeat, pHC624-(AGGAG)₂₂, were examined by agarose gel electrophoresis. Lane 1: pSK-(AGGAG)₂₂ linearized with *Xba* I; lane 2: linearized pSK-(AGGAG)₂₂ transcribed with T7 RNA polymerase; lane 3: supercoiled pHC624-(AGGAG)₂₂, which lacked T7 promoter; lane 4: supercoiled pHC624-(AGGAG)₂₂ incubated in transcription mixture with T7 RNA polymerase; lane 5: supercoiled pHC624-(AGGAG)₂₂ and linearized pSK-(AGGAG)₂₂ transcribed with T7 RNA polymerase; lane 6: supercoiled pHC624-(AGGAG)₂₂ and pBluescript SK(-) linearized with *Xba* I, transcribed with T7 RNA polymerase; lane 7: supercoiled pHC624; lane 8: supercoiled pHC624 and linearized pSK-(AGGAG)₂₂ incubated in transcription mixture with T7 RNA polymerase; lane 9: pHC624-(AGGAG)₂₂ relaxed with vaccinia topoisomerase I; lane 10: relaxed pHC624-(AGGAG)₂₂ and linearized pSK-(AGGAG)₂₂ transcribed with T7 RNA polymerase; lane 11: same as lane 5 and ethanol-precipitated; lane 12: same as lane 11 and treated with 90 units of *E. coli* RNase H (TaKaRa) in 40 mM Tris-HCl pH 7.7, 4 mM MgCl₂, 1 mM DTT, 4% glycerol, and 0.003% bovine serum albumin at 37°C for 1 h.

(lanes 9 and 10), indicating that the R-loop formation *in trans* depended on the superhelicity of the plasmid DNA.

This R-loop formation *in trans* was further examined by linearizing pHC624-(AGGAG)₂₂ and radiolabeling the transcript. Transcription of linearized pSK-(AGGAG)₂₂ generated an extra band with a slightly reduced mobility (Figure 2). Radioactivity was found to be associated with this extra band (Figure 3, lanes 4–6), indicating the mobility shift was due to the presence of the R-loop. Transcription of linearized pSK-(AGGAG)₂₂ resulted in relaxation of supercoiled pHC624-(AGGAG)₂₂ and radioactivity was associated with the relaxed species (lane 5). By contrast, little radioactivity was associated with pHC624-(AGGAG)₂₂ when the DNA was linearized (lane 6). These confirmed the dependence of the R-loop formation *in trans* on supercoiling.

3.4. Formation of an Alternative Structure at the AGGAG Repeat by DNA Supercoiling. The results obtained thus far suggested that negative supercoiling induced a structural




FIGURE 3: R-loop formation *in trans* detected by radiolabeling of the transcript. (a) Agarose gel stained with ethidium bromide and (b) autoradiogram of the same gel. Lane 1: pBluescript SK(-) linearized with *Xba* I; lane 2: supercoiled pHC624-(AGGAG)₂₂; lane 3: pHC624-(AGGAG)₂₂ linearized with *Sca* I; lane 4: pSK-(AGGAG)₂₂ linearized with *Xba* I; lane 5: linearized pSK-(AGGAG)₂₂ and supercoiled pHC624-(AGGAG)₂₂; and lane 6: linearized pSK-(AGGAG)₂₂ and linearized pHC624-(AGGAG)₂₂. All DNAs were incubated in the transcription mixture with T7 RNA polymerase.

change and the CTCCT-repeat strand became accessible to the AGGAG-repeat transcript in the altered structure. Chemical modification was employed to detect the structural change. Figure 4(a) shows the result for oxidation of T residues in the CTCCT-repeat strand by permanganate [21]. T residues were found to be oxidized more readily when the DNA was supercoiled than when relaxed. Permanganate is known to oxidize T residues (and C residues to a lesser extent) preferentially in single-stranded forms [21]. Therefore, the result indicated that supercoiling induced, a transition to an alternative structure in which the CTCCT-repeat strand was single stranded. The AGGAG-repeat strand was probed with diethylpyrocarbonate. The chemical is known to readily react with A (and less so with G) preferentially in single-stranded forms [22]. As shown in Figure 4(b), A residues in the central AGGAG units, especially the second A residues in the units, were more reactive to diethylpyrocarbonate when the DNA was supercoiled.

4. Discussion

One possible mechanism that allows reassociation would be zippering of RNA and DNA by backtracking of RNA polymerase [24]. However, in the current model of the elongation complex, backtracking dissociates the 3' end of the transcript from the template [25]. This will limit the size of the R-loop to that of the transcription bubble, which is only several base pairs [2, 25]. A result on R-loop formation in *Saccharomyces cerevisiae* cells seems to suggest against

backtracking as well, if the mechanism is assumed to be the same. Placing of a ribozyme sequence downstream of a *PHO5* was shown to reduce R-loop formation at the *PHO5* [10]. The result indicated that at least some population of the R-loop formed after the RNA polymerase had transcribed the ribozyme region, which was 52 bp. This would require the backtracking to start further downstream and to continue through the 52 bp region until reaching the *PHO5* to start forming an R-loop, both of which seem very unlikely. The present result of R-loop formation *in trans* also indicates that backtracking is not necessary, although the result does not exclude backtracking.

The present work showed that an R-loop at the AGGAG repeat formed *in trans* and the formation *in trans* was facilitated by negative supercoiling of the target DNA. The results of chemical probing of the repeat DNA under supercoiling were consistent with a structure in which the 5' side in the CTCCT repeat was single-stranded (Figure 4). Thus, the R-loop formation *in trans* observed here is most likely to be the association of the AGGAG-repeat transcript with the CTCCT-repeat template that was made accessible by the structural conversion. Formation of alternative structures had previously been studied for (AGGAG)₆GA [26]. The previous result of diethylpyrocarbonate modification at low pH was consistent with formation of an intramolecular triple helix. The present result of diethylpyrocarbonate modification was different from the previous one and, therefore, the alternative structure observed here is likely to be different from the previous one, which was a canonical H-form. Another structure had also been observed at neutral pH




FIGURE 4: Supercoiling-induced structural change in the AGGAG repeat detected by chemical modification. Reactions of the Maxam-Gilbert type were carried out on relaxed and supercoiled pSK-(AGGAG)₂₂ and analyzed by sequencing gel. (a) Oxidation of T residues by permanganate. Lane 1: Maxam-Gilbert C reaction; lane 2: relaxed DNA; lane 3: supercoiled DNA. The large bracket indicates the repeat and the small one, the 5'-most CTCCT unit. (b) Modification of A residues by diethylpyrocarbonate. Lane 1: Maxam-Gilbert G reaction; lane 2: supercoiled DNA; lane 3: relaxed DNA. The large bracket indicates the repeat and the small one, the 5'-most AGGAG unit. (c) An H-r3 structure consistent with the modification results. Filled circles indicate Watson-Crick base pairing, and open circles indicate triplex formation. Triangles indicate the bases expected to be reactive to the chemicals.

[26]. The regions of enhanced chemical modification were different between the previous and the present results, and thus, the structures are likely to be different. The difference is likely to be due to the number of the repeats and/or the presence of Mg(II), which was absent in the previous study, as dependence of the formation of alternative forms on the size of the homopurine-homopyrimidine stretch, metal ions, and pH has been well documented [27–29]. One structure consistent with our chemical modification results is depicted in Figure 4(c), which is an H-r3 isoform of the *H-form [27, 28]. Alternative fold-back points would result in stronger modification sites in the middle of the AGGAG strand.

Another possible mechanism consistent with the present findings would be one that involves opening of the duplex in the wake of the advancing RNA polymerase and ensuing hybridization of the transcript RNA to the template DNA, which has been proposed by Grabczyk et al. [30]. They have posited that negative supercoiling behind the advancing RNA polymerase induces triplex formation at GAA repeats, however, with no data to show the formation. The present study demonstrated that extensive helix opening does occur

at an AGGAG repeat under negative supercoiling, indicating the repeat's potential of an alternative conformation. There is also circumstantial evidence linking R-loop formation at *E. coli* *rrnB* to supercoiling both *in vitro* and *in vivo* [31, 32]. It is debatable, however, whether the transient supercoiling behind the polymerase is strong enough to induce the conformation transition on small purified DNA molecules *in vitro*, especially because R-loop has been found to form on small linearized DNA that cannot be topologically constrained [12, 33]. Duquette et al. [23] studied G-loop formation in immunoglobulin class-switch regions and a telomeric repeat and have proposed that invasion of RNA into the duplex DNA initiates the formation. This general idea of RNA invasion into duplex is supported by the finding that a nick in the nontemplate strand can serve as an initiation site for R-loop at immunoglobulin S_{γ3} repeats *in vitro*: fraying of the nontemplate strand at the nick should increase the accessibility of the template strand to the transcript [33]. Finally, we would like to raise the possibility that the transient duplex separation by the RNA polymerase could be more persistent for some templates, which may involve interaction between

DNA single strands and the RNA polymerase, and also the possibility that the nontemplate strand may contribute to the stability of the R-loop by the base interactions involved in triple helix formation [27, 28, 34, 35].

5. Conclusions

In vitro experiments showed that R-loop formed *in trans* at an AGGAG repeat when the DNA comprising the repeat was negatively supercoiled. Chemical modifications indicated that the duplex at the repeat opened under negative supercoiling. These results were the first to connect structural conversion, duplex opening, and R-loop formation for the same sequence, supporting the importance of the potential of duplex opening in R-loop formation.

Conflict of Interests

The authors declare that they do not have any conflict of interests with the content of the paper.

Acknowledgment

This work was supported in part by the Strategic Research Foundation Grant-aided Project for Private Universities (S1201003) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

References

- [1] J. P. Richardson, "Attachment of nascent RNA molecules to superhelical DNA," *Journal of Molecular Biology*, vol. 98, no. 3, pp. 565–579, 1975.
- [2] M. Jiang, N. Ma, D. G. Vassilyev, and W. T. McAllister, "RNA displacement and resolution of the transcription bubble during transcription by R7 RNA polymerase," *Molecular Cell*, vol. 15, no. 5, pp. 777–788, 2004.
- [3] Y. W. Yin and T. A. Steitz, "Structural basis for the transition from initiation to elongation transcription in T7 RNA polymerase," *Science*, vol. 298, no. 5597, pp. 1387–1395, 2002.
- [4] T. H. Tahlrov, D. Temiakov, M. Anikin et al., "Structure of a T7 RNA polymerase elongation complex at 2.9 Å resolution," *Nature*, vol. 420, no. 6911, pp. 43–50, 2002.
- [5] T. Naryshkina, K. Kuznedelov, and K. Severinov, "The role of the largest RNA polymerase subunit lid element in preventing the formation of extended RNA-DNA hybrid," *Journal of Molecular Biology*, vol. 361, no. 4, pp. 634–643, 2006.
- [6] I. Touloukhonov and R. Landick, "The role of the lid element in transcription by *E. coli* RNA polymerase," *Journal of Molecular Biology*, vol. 361, no. 4, pp. 644–658, 2006.
- [7] A. Aguilera and T. García-Muse, "R loops: from transcription byproducts to threats to genome stability," *Molecular Cell*, vol. 46, no. 2, pp. 115–124, 2012.
- [8] H. Masukata and J.-I. Tomizawa, "A mechanism of formation of a persistent hybrid between elongating RNA and template DNA," *Cell*, vol. 62, no. 2, pp. 331–338, 1990.
- [9] Q. Sun, T. Csorba, K. Skourti-stathaki, N. J. Proudfoot, and C. Dean, "R-loop stabilization represses antisense transcription at the *Arabidopsis FLC* locus," *Science*, vol. 340, no. 6132, pp. 619–621, 2013.
- [10] P. Huertas and A. Aguilera, "Cotranscriptionally formed DNA:RNA hybrids mediate transcription elongation impairment and transcription-associated recombination," *Molecular Cell*, vol. 12, no. 3, pp. 711–721, 2003.
- [11] M. E. Reaban and J. A. Griffin, "Induction of RNA-stabilized DNA conformers by transcription of an immunoglobulin switch region," *Nature*, vol. 348, no. 6299, pp. 342–344, 1990.
- [12] M. E. Reaban, J. Lebowitz, and J. A. Griffin, "Transcription induces the formation of a stable RNA-DNA hybrid in the immunoglobulin α switch region," *Journal of Biological Chemistry*, vol. 269, no. 34, pp. 21850–21857, 1994.
- [13] G. A. Daniels and M. R. Lieber, "RNA:DNA complex formation upon transcription of immunoglobulin switch regions: implications for the mechanism and regulation of class switch recombination," *Nucleic Acids Research*, vol. 23, no. 24, pp. 5006–5011, 1995.
- [14] K. Kinoshita, J. Tashiro, S. Tomita, C.-G. Lee, and T. Honjo, "Target specificity of immunoglobulin class switch recombination is not determined by nucleotide sequences of S regions," *Immunity*, vol. 9, no. 6, pp. 849–858, 1998.
- [15] K. Yu, F. Chedin, C.-L. Hsieh, T. E. Wilson, and M. R. Lieber, "R-loops at immunoglobulin class switch regions in the chromosomes of stimulated B cells," *Nature Immunology*, vol. 4, no. 5, pp. 442–451, 2003.
- [16] R. Shinkura, M. Tian, M. Smith, K. Chua, Y. Fujiwara, and F. W. Alt, "The influence of transcriptional orientation on endogenous switch region function," *Nature Immunology*, vol. 4, no. 5, pp. 435–441, 2003.
- [17] N. Takahashi, N. Sasagawa, K. Suzuki, and S. Ishiura, "Synthesis of long trinucleotide repeats *in vitro*," *Neuroscience Letters*, vol. 262, no. 1, pp. 45–48, 1999.
- [18] P. T. Chan and J. Lebowitz, "Site-directed mutagenesis of the -10 region of the lacUV5 promoter. Introduction of dA₄-dT₄ tract suppresses open complex formation," *Journal of Biological Chemistry*, vol. 265, no. 7, pp. 4091–4097, 1990.
- [19] J. Sambrook and D. W. Russel, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA, 3rd edition, 2001.
- [20] S. Shuman, M. Golder, and B. Moss, "Characterization of vaccinia virus DNA topoisomerase I expressed in *Escherichia coli*," *Journal of Biological Chemistry*, vol. 263, no. 31, pp. 16401–16407, 1988.
- [21] S. Sasse-Dwight and J. D. Gralla, "Footprinting protein-DNA complexes *in vivo*," *Methods in Enzymology*, vol. 208, pp. 146–168, 1991.
- [22] T. Kohwi-Shigematsu and Y. Kohwi, "Detection of non-B-DNA structures at specific sites in supercoiled plasmid DNA and chromatin with haloacetaldehyde and diethyl pyrocarbonate," *Methods in Enzymology*, vol. 212, pp. 155–180, 1992.
- [23] M. L. Duquette, P. Handa, J. A. Vincent, A. F. Taylor, and N. Maizels, "Intracellular transcription of G-rich DNAs induces formation of G-loops, novel structures containing G4 DNA," *Genes and Development*, vol. 18, no. 13, pp. 1618–1629, 2004.
- [24] N. Komissarova and M. Kashlev, "RNA polymerase switches between inactivated and activated states by translocating back and forth along the DNA and the RNA," *Journal of Biological Chemistry*, vol. 272, no. 24, pp. 15329–15338, 1997.
- [25] N. Korzheva, A. Mustaev, E. Nudler, V. Nikiforov, and A. Goldfarb, "Mechanistic model of the elongation complex of *Escherichia coli* RNA Polymerase," *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 63, pp. 337–345, 1998.

- [26] D. A. Collier, J. A. Griffin, and R. D. Wells, "Non-B right-handed DNA conformations of homopurine-homopyrimidine sequences in the murine immunoglobulin C_α switch region," *Journal of Biological Chemistry*, vol. 263, no. 15, pp. 7397–7405, 1988.
- [27] S. M. Mirkin and M. D. Frank-Kamenetskii, "H-DNA and related structures," *Annual Review of Biophysics and Biomolecular Structure*, vol. 23, pp. 541–576, 1994.
- [28] M. D. Frank-Kamenetskii and S. M. Mirkin, "Triplex DNA structures," *Annual Review of Biochemistry*, vol. 64, pp. 65–95, 1995.
- [29] S. Burge, G. N. Parkinson, P. Hazel, A. K. Todd, and S. Neidle, "Quadruplex DNA: sequence, topology and structure," *Nucleic Acids Research*, vol. 34, no. 19, pp. 5402–5415, 2006.
- [30] E. Grabczyk, M. Mancuso, and M. C. Sammarco, "A persistent RNA-DNA hybrid formed by transcription of the Friedreich ataxia triplet repeat in live bacteria, and by T7 RNAP *in vitro*," *Nucleic Acids Research*, vol. 35, no. 16, pp. 5351–5359, 2007.
- [31] P. Phoenix, M.-A. Raymond, É. Massé, and M. Drolet, "Roles of DNA topoisomerases in the regulation of R-loop formation *in vitro*," *Journal of Biological Chemistry*, vol. 272, no. 3, pp. 1473–1479, 1997.
- [32] S. Broccoli, F. Rallu, P. Sanscartier, S. M. Cerritelli, R. J. Crouch, and M. Drolet, "Effects of RNA polymerase modifications on transcription-induced negative supercoiling and associated R-loop formation," *Molecular Microbiology*, vol. 52, no. 6, pp. 1769–1779, 2004.
- [33] D. Roy, Z. Zhang, Z. Lu, C.-L. Hsieh, and M. R. Lieber, "Competition between the RNA transcript and the nontemplate DNA strand during R-Loop formation *in vitro*: a nick can serve as a strong R-loop initiation site," *Molecular and Cellular Biology*, vol. 30, no. 1, pp. 146–159, 2010.
- [34] R. W. Roberts and D. M. Crothers, "Stability and properties of double and triple helices: dramatic effects of RNA or DNA backbone composition," *Science*, vol. 258, no. 5087, pp. 1463–1466, 1992.
- [35] H. Han and P. B. Dervan, "Sequence-specific recognition of double helical RNA and RNA-DNA by triple helix formation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 90, no. 9, pp. 3806–3810, 1993.