



Review

# Diversified Stimuli-Induced Inflammatory Pathways Cause Skin Pigmentation

Md Razib Hossain, Tuba M. Ansary, Mayumi Komine \*  and Mamitaro Ohtsuki

Department of Dermatology, Faculty of Medicine, Jichi Medical University, Tochigi 329-0498, Japan; razib@jichi.ac.jp (M.R.H.); tuba2020@jichi.ac.jp (T.M.A.); mamitaro@jichi.ac.jp (M.O.)

\* Correspondence: mkomine12@jichi.ac.jp; Tel.: +0285-58-7380

**Abstract:** The production of melanin pigments by melanocytes and their quantity, quality, and distribution play a decisive role in determining human skin, eye, and hair color, and protect the skin from adverse effects of ultraviolet radiation (UVR) and oxidative stress from various environmental pollutants. Melanocytes reside in the basal layer of the interfollicular epidermis and are compensated by melanocyte stem cells in the follicular bulge area. Various stimuli such as eczema, microbial infection, ultraviolet light exposure, mechanical injury, and aging provoke skin inflammation. These acute or chronic inflammatory responses cause inflammatory cytokine production from epidermal keratinocytes as well as dermal fibroblasts and other cells, which in turn stimulate melanocytes, often resulting in skin pigmentation. It is confirmed by some recent studies that several interleukins (ILs) and other inflammatory mediators modulate the proliferation and differentiation of human epidermal melanocytes and also promote or inhibit expression of melanogenesis-related gene expression directly or indirectly, thereby participating in regulation of skin pigmentation. Understanding of mechanisms of skin pigmentation due to inflammation helps to elucidate the relationship between inflammation and skin pigmentation regulation and can guide development of new therapeutic pathways for treating pigmented dermatosis. This review covers the mechanistic aspects of skin pigmentation caused by inflammation.

**Keywords:** melanosome; melanocytes; melanogenesis; skin pigmentation; inflammation; inflammatory cytokine



**Citation:** Hossain, M.R.; Ansary, T.M.; Komine, M.; Ohtsuki, M. Diversified Stimuli-Induced Inflammatory Pathways Cause Skin Pigmentation. *Int. J. Mol. Sci.* **2021**, *22*, 3970. <https://doi.org/10.3390/ijms22083970>

Academic Editor: Javier Conde Aranda

Received: 15 March 2021  
Accepted: 7 April 2021  
Published: 12 April 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Skin color is one of the topmost concerns for human beings as it provides uniformity, identity and represents the overall appearance of a person [1]. Production, quantity, quality, and distribution of melanin, a group of natural pigments, determines the color of human skin, eyes, and hair. It is crucially involved in the defense mechanism to protect skin from adverse effects of ultraviolet radiation (UVR) and oxidative stress of environmental pollutants [2]. Synthesis of melanin starts with formation of melanosomes, melanin-containing organelles, in epidermal melanocytes by a process called melanogenesis [3,4]. Melanocytes are specialized cells derived from unpigmented precursor cells called melanoblasts, originating from embryonic neural crest cells which can migrate towards the skin and other tissues during embryogenesis [5–8]. Skin pigmentation is a specific and complex mechanism that occurs due to accumulation of melanosomes in keratinocytes to protect skin from solar irradiation. Melanin pigment can be divided into two types—pheomelanin and eumelanin. Pheomelanin has a reddish to brownish color, causing reactive oxygen species production under UVR stimulation. Eumelanin has black color, which works mainly to protect the nucleus from UVR. Skin color differences can emerge from the quantity and quality (pheo/eumelanin ratio) of melanin produced as well as size, number, composition, mode of transfer, distribution, and degradation of melanosomes inside keratinocytes, whereas melanocytes numbers typically remain relatively constant [8]. Keratinocytes have a significant role in regulating adhesion, proliferation, survival, and morphology of melanocytes [8].

There are various nonspecific intrinsic factors (hormonal environment, inflammation) and extrinsic factors (solar irradiation such as ultraviolet irradiation, environmental pollution, drugs) modulating genetically determined melanin levels. Various stimuli including skin eruptions such as eczema, allergens, microbial infection, pathogens, chemical stimuli and physical damage, ultraviolet light exposure, and aging can all lead to skin inflammation [9–12], defined as a defensive reaction of living tissues with a vascular system in response to exogenous and endogenous stimuli [13]. In dermatology and cosmetology, hyperpigmentation or hypopigmentation after inflammation is a major problem and commonly seen due to acute or chronic inflammatory skin reactions [14]. Inflammatory mediators are chemical factors involved in mediating inflammatory reactions and mainly secreted by T helper cells (Th), monocytes and macrophages, dendritic cells, as well as epidermal keratinocytes. Recent studies have revealed a close relationship between inflammatory cytokines and skin pigmentation. Diversified interleukins (ILs) and varieties of inflammatory mediators can also participate in melanogenesis regulation by modulating proliferation and differentiation of human epidermal melanocytes, and also promote or inhibit melanogenesis-related gene expression directly or indirectly [14–16]. Many genes are involved in regulating pigmentation at various levels, as well as mutations in several cause pigmentary disorders.

The above studies draw attention to the relationship between inflammation and skin pigmentation regulation, as well as guide new therapeutic pathway development for treating pigmented dermatosis. In this review, we describe the mechanistic perspectives of skin pigmentation due to inflammation.

## 2. Melanogenesis Process


Skin epidermal units are responsible for melanin production by a complex multi-stage process called melanogenesis. These units are composed of melanocytes surrounded by keratinocytes. Melanocyte interacts with the endocrine, immune, inflammatory and central nervous systems, and its activity is also regulated by extrinsic factors such as UVR and drugs. Melanocytes produce melanin pigment in melanosomes, which are transferred to surrounding keratinocytes. The production of melanin in melanocytes and accumulation of melanin-containing melanosomes in keratinocytes leads to skin pigmentation and its dysregulation causes different types of pigmentation defects, which are classified as hypopigmentation (including oculocutaneous albinism (OCA)), hyperpigmentation (such as melasma) or mixed hyper-/hypopigmentation (including dyschromatosis symmetrica hereditaria) and may occur with or without alteration of the number of melanocytes [8,17,18]. Congenital defects in melanin synthesis or melanosome function or migration of melanoblasts cause congenital hypopigmented disorders, such as OCA, Chediak Higashi syndrome, or piebaldism. On the other hand, congenital hyperpigmented conditions depend more on locally distributed melanocytes or nevus cells, other types of melanin-producing cells derived from neural crest cells such as linear and whorled nevoid hypermelanosis, and congenital melanocytic nevus. Acquired conditions include vitiligo vulgaris, halo nevus, melasma, and senile pigmented macules, which mainly depend on immune dysfunction, inflammation or aging.

Ectodermal neural crest melanoblasts are the origin of melanocytes, and melanoblasts migrate to different destinations as the embryo develops, including the basal layer of the epidermis, hair follicles, eye uvea, vestibular organ, endolymphatic sac and pia mater [19,20]. Various mediators and transcription factors are involved in maturation and migration of melanoblasts to their final destination, such as Paired box protein Pax-3 (PAX3), sex-determining region Y-box 10 (SOX10), endothelin (EDN) 3, Microphthalmia-associated transcription factor (MITF), and stem cell growth factor receptor (KIT) and its ligand stem cell growth factor (SCF) [20,21]. Once they differentiate into melanocytes in the epidermis, various factors produced by the surrounding keratinocytes influence the function of melanocytes, such as SCF, EDN1, Nerve growth factor (NGF), Hepatocyte growth factor (HGF), granulocyte-macrophage colony stimulating factor (GM-CSF),  $\alpha$ -Melanocyte-

stimulating hormone ( $\alpha$ -MSH), Adrenocorticotrophic hormone (ACTH), prostaglandin F2 (PGF2) and prostaglandin E2 (PGE2).

It is postulated that solar lentigines (SLs), hyper-pigmentary spots on sun-exposed skin of aged persons, could be treated by combined topical treatment with EDN signaling blockers and tyrosinase inhibitors [22], and SCF inhibitors or EDN1-triggered intracellular signaling cascades can downregulate the hyperpigmentation observed in solar lentigo, UVB (ultraviolet ray-B) induced melanosis, and melasma [23]. Melanosomes, subcellular lysosome-like organelles in melanocytes, store melanin [24,25] and synthesize two types of melanin pigment: eumelanin (brown-black or dark insoluble polymer) and pheomelanin (red-yellow soluble polymer). After synthesis, mature melanosomes are carried by molecular motors such as kinesins from the perinuclear region to dendrites of melanocytes, interacting with microtubules. On the other hand, dynein works in maintaining perinuclear distribution of melanosomes in melanocytes [26,27]. In the epidermis, each melanocyte is surrounded by 30 to 40 keratinocytes through dendrites and transfers mature melanosomes into the cytoplasm of keratinocytes [8,19]. The size and the number of melanosomes, as well as amount, type, transfer, and distribution of melanin in keratinocytes, determine phenotypic diversity of pigmentation, whereas melanocyte numbers are relatively constant in different ethnic groups [8,19,20]. Delayed degradation of melanosomes in keratinocytes has been found to increase visible pigmentation in dark-skinned individuals which is due to large, more numerous, and elongated melanosomes [19,20,28]. These differences in melanosomes are observed at the time of birth and are not determined by extrinsic factors such as UVR, which indicates that mutation in specific genes is related to the control of skin pigmentation.

Various molecules are involved in skin pigmentation, divided into three distinct groups: proteins required for melanosome structure, enzymes involved in melanin synthesis, and proteins required for melanosome transport and distribution [29]. There are four stages (I–IV) of melanosomal maturation and three important structural proteins (Figure 1) forming melanosomes, including melanosomal matrix protein (PMEL17/Silv/GP100), melanoma antigen recognized by T cell-1 (MART-1), and glycoprotein nonmetastatic melanoma protein b (GPNMB/DC-HIL/osteostatin). Stages I and II are considered as pre-melanosomes and do not contain melanin. In stage III melanosomes, melanin is deposited on PMEL17 fibrils. Melanin fills the melanosome in the last stage and forms a masked internal structure and dark color. MART-1, found in earlier melanosomal stages, is essential for the maturation of PMEL17 [30]. GPNMB, a melanosome-specific and proteolytically released protein, is abundant in late melanosomes [31] and critical for the formation of melanosomes in an MITF-independent fashion [32]. Scaffold materials are provided by these structural melanosomal proteins for melanin synthesis.



**Figure 1.** Factors that regulate melanosome formation and maturation through four distinct stages (I–IV) during the melanin production by a melanocyte. The melanogenesis takes place in a special organelles called melanosomes. Melanosomal structural proteins (PMEL17, MART-1, and GPNMB) affect melanosome structure. Enzymes (TYR, TYRP1, TYRP2, BLOCs, APs) are involved in melanin synthesis either direct or indirect manner. Proteins (microtubules, F-actin, Kinesin, dynein, MLPH, MYO5A, RAB27A) ensure intracellular transport of melanosomes. Melanocyte-specific transcription factors (specially MITF) regulate the expression and function of many pigment-specific factors. ER: endoplasmic reticulum; PMEL17: melanocytes lineage-specific antigen gp100; MART-1: melanoma antigen recognized by T cell-1; GPNMB: glycoprotein nonmetastatic melanoma protein b; TYR: tyrosinase; TYRP1: tyrosinase-related protein-1; TYRP2: tyrosinase-related protein-2; BLOCs: biogenesis of lysosome-related organelles complexes; Aps: adaptor protein complexes; MLPH: melanophilin; MYO5A: myosin-Va; RAB27A: Ras-related protein Rab27A; MITF: microphthalmia-associated transcription factor.

Melanogenic enzymes need to be delivered in melanosomes to reach the final stage of melanosomal maturation and melanin synthesis (stages III–IV). There are three key enzymatic components of melanosomes playing central roles in melanin synthesis: tyrosinase (TYR), tyrosinase-related protein-1 (TYRP1), and dopachrome tautomerase/tyrosinase-related protein-2 (DCT/TYRP2). TYR is a copper-dependent enzyme catalyzing conversion of L-tyrosine into L-3,4-dihydroxyphenylalanine (L-DOPA), the rate-limiting stage in melanin synthesis. This enzyme is deactivated by copper oxidation but can be activated by electron donors such as L-DOPA, ascorbic acid, superoxide anion, and possibly nitric oxide (NO) [14,17,33]. Protein kinase C- $\beta$  (PKC- $\beta$ ) phosphorylates two serine residues of the cytoplasmic domain which are also important for tyrosinase activation [34]. A most severe form of albinism OCA1a occurs due to the mutation inactivating this enzyme. TYRP1 and TYRP2 also exist in the membrane of melanosomes. Although its specific role is not yet well-established, it is assumed that TYRP1 has a role in activation and stabilization of tyrosinase, melanosome synthesis, increasing eumelanin/pheomelanin ratio and working against oxidative stress due to its peroxidase effect [14,17].

Melanosomes are transported from the perinuclear area to periphery of melanocytes, and melanocytes transfer packaged melanin into adjacent keratinocytes for skin pigmentation (Figure 1) [29]. Early melanosomes, originating in the perinuclear area, gradually become mature then turn to late (pigmented) melanosomes, while they move toward periphery of melanocytes (i.e., dendrites). Microtubules and collaboration of microtubule-associated motor proteins (kinesins and dyneins) are involved in trafficking of mature

melanosomes from the melanocyte perinuclear area to dendrites [26,27]. Kinesins ensure centrifugal movement of melanosomes, and melanosomal cargo is transferred from microtubules to F-actin in dendrites via the complex of RAB27A, melanophilin (MLPH) and myosin-Va (MYO5A). It is demonstrated by mouse Melan-a cells that RAB27A links to synaptotagmin-like 2 (SYTL2) prompting SYTL2 to dock melanosome at the plasma membrane [35,36].

However, the exact transfer mechanism of human melanosome to keratinocytes is still unclear, but four different mechanisms have been proposed: (i) exocytosis, (ii) cytophagocytosis, (iii) fusion of plasma membrane, and (iv) transfer by membrane vesicles [37].

### 3. Melanogenesis Regulation

MITF is a basic helix–loop–helix leucine zipper (bHLH-ZIP) transcription factor belonging to the MIT regulating melanocyte function. MITF controls expression of the melanogenesis enzymes TYR, TYRP1 and TYRP2 [38] and regulates melanocyte functions, including pigmentation, melanocyte differentiation, proliferation, and cell survival [39]. Furthermore, MITF protein plays a vital role in expression regulation of the RAB27A protein, crucial in melanosome transport and the melanosomal matrix protein Pmel17 [17,21,40,41]. It has also been discovered that the MIT (microphthalmia) family is the main transcriptional regulator of many genes coding for proteins involved in pigment synthesis, melanosomal structure, as well as melanosome trafficking [38]. Many other transcription factors are involved in regulating MITF expression, including a paired box protein 3 (PAX3), sex-determining region Y-box 9 and 10 (SOX9, SOX10), lymphoid enhancer-binding factor 1 (LEF-1) and Cyclic adenosine monophosphate (cAMP) responsive-element binding protein (CREB), which is phosphorylated by signals via melanocortin-1 receptor (MC1R) [29]. Recent findings have elucidated MITF expression regulation in mice melanocytes is influenced by salt-inducible kinase (SIK), which phosphorylates CREB-regulated transcription coactivator (CRTC) to prevent its nuclear translocation, resulting in inhibited CREB-induced MITF upregulation [42].

#### *Involvement of Signaling Pathways in Melanogenesis Regulation*

The regulation of melanogenesis is a complex process and more than 150 genes are involved in this regulation [43]. Melanogenesis can be controlled at different levels as many of these genes affect the developmental process crucial to melanoblasts, while others are involved in melanocyte survival and differentiation [44]. At least 25 of these genes are involved in regulating different pigmentation pathways or function of melanosomes. The cAMP/protein kinase A (PKA) signaling pathway is one of the most significant signaling pathways and the most well-known receptor is the MC1R. MC1R belongs to G protein-coupled receptor family and is expressed predominantly in melanocytes. The  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH) binds to MC1R and activates adenylate cyclase (AC), increasing intracellular concentration of secondary messenger, cAMP. The cAMP activates PKA, then PKA phosphorylates CREB, leading to increased expression of MITF, resulting in melanogenesis promotion [17,28,45,46].


In addition to the above, other signaling pathways such as mitogen-activated protein kinase (MAPK), inositol trisphosphate/diacylglycerol (IP3/DAG), wingless-type protein (WNT), and PKC are reported to participate in the melanogenesis process. The IP3/DAG pathway is activated by the  $\alpha$ 1 adrenergic receptor ( $\alpha$ 1AR) and thus increases the intracellular levels of PKC- $\beta$  and activates tyrosinase [47]. The c-KIT, granulocyte-macrophage colony-stimulating factor receptor (GM-CSFR), and HGF receptor-mediated signaling pathways can be activated by SCF, GM-CSF and HGF, respectively, which leads to autophosphorylation and activation of MAPK, thereby phosphorylating MITF and upregulating melanogenesis-related enzyme expression [3,48,49]. The WNT bind to class frizzled (FZD) receptors and the wingless-type protein (WNT)/FZD signaling pathway is involved in several aspects of the mouse melanocyte physiology, from developmental lineage differentiation to subsequent maintenance of melanocytes [50]. Indeed, pigment development

and differentiation from the neural crest cells into melanocytes can be promoted by WNT1 and WNT3A in zebrafish [51]. Several models including mouse Melan-a cells and avian embryo melanocytes have demonstrated WNT1 is responsible for increasing the number of melanoblasts differentiating from melanoblast precursors, whereas WNT3A is involved in promotion of the differentiation of neural crest cells and melanoblasts into melanocytes by maintaining MITF expression [52–54]. In general, the WNT signaling pathway can activate the melanocytes-specific MITF isoform (MITF-M) promoter [55–57] and upregulate MITF expression, thus participating in melanogenesis regulation. Catecholamines can promote melanogenesis through the cAMP/PKA pathway and also mediate melanogenesis through PKC- $\beta$  pathways by  $\alpha$ 1 and  $\beta$ 2 adrenergic receptors [3,47].

#### 4. Skin Inflammation


The skin is the largest organ in the human body and provides the first line of defense against pathogens and other environmental pollutants [58]. These external stimuli as well as intrinsic stimuli, such as neurotransmitters, advanced glycation end products, amyloids, and debris of dead cells, provoke either short-term or long-term skin inflammation to the host [59]. Consequently, the production of reactive oxygen species (ROS), known as the first line of defense against pathogen and external pollutants, is promoted either directly or indirectly, and thus ROS stimulates the inflammation process [60,61]. External and intrinsic stimuli trigger inflammation, resulting in immunologic reactions (e.g., dendritic cell activation followed by T cell activation, and finally leading to inflammatory disorders, such as rheumatoid arthritis and psoriasis), which involve the influx of various cells of the host immune system to the inflammation site and release a large number of inflammatory mediators to enhance inflammation. Inflammation is primarily divided into two types: acute inflammation (winds up less than 48 h and neutrophils are main cell type) and chronic inflammation (rests more than 48 h, lymphocytes, macrophages, plasma cells are major participating cells). Acute inflammation may lead to chronic inflammation. Cytokine expression is induced not only by acute inflammation but also by chronic inflammation and is also considered responsible for skin pigmentation. Proinflammatory cytokines (that favor inflammation) or anti-inflammatory cytokines (that neutralize inflammation) accelerate/decelerate the process of inflammation and are involved in controlling inflammatory reactions either indirectly or directly and stimulate/suppress production of cell adhesion molecules or several other inflammatory molecules in diverse cell types.

Generally, pathogen-associated molecular patterns (PAMPs) are derived from microorganisms and trigger the inflammatory response to infections through activation of germline-encoded pattern-recognition receptors (PRRs) [62,63]. Lipopolysaccharide (LPS) is the most common PAMP found on the outer cell wall of Gram-negative bacteria. On the other hand, damage-associated molecular patterns (DAMPs) are derived from host cells exposed to environmental stimuli, such as trauma, heat, and UVR, or from damaged tissue due to ischemia, hypoxia, or genetic defects. DAMPs do not require pathogenic infection but induce sterile inflammatory responses. PRRs families consist of the Toll-like receptors (TLRs), C-type lectin receptors (CLRs), retinoic acid-inducible gene (RIG)-I-like receptors (RLRs), and NOD-like receptors (NLRs) [64]. Among them, TLRs mostly participate in activation of inflammatory responses as a consequence of PAMPs and DAMPs interaction [65]. TLRs activate an intracellular signaling cascade and transfer signals through myeloid differentiation primary response protein-88 (MyD88) leading to nuclear translocation of transcription factors, such as activator protein-1 (AP-1) and nuclear factor kappa-B (NF- $\kappa$ B) or interferon regulatory factor (IRF) 3 (Figure 2). TLRs are shared by both DAMPs and PAMPs to maintain similarities between infectious and noninfectious inflammatory responses [66,67].



**Figure 2.** DAMPs and PAMPs bind to PRRs (TLRs, NLRs, RLPs) and activate several signaling pathways. **(A)** TLRs activate MyD88 or TAM signaling pathways, which lead to nuclear translocation of NF- $\kappa$ B and AP-1 or IRF3, which plays a crucial role in the regulation of inflammatory response. **(B)** TLRs and some inflammatory cytokines, such as IL-1 and TNF trigger NF- $\kappa$ B pathway resulting activation of RelA/p50 complexes that regulate expression of inflammatory cytokines. IKK subunits phosphorylate I $\kappa$ B and regulate NF- $\kappa$ B pathway activation. **(C)** The binding of IL-6 family members to plasma membrane receptors activates the JAKs. JAKs activate STATs which are dephosphorylated in the nucleus and activate downstream cytokines. **(D)** MAPKs direct cellular response to a variety of stimuli, including stress, mitogens and inflammatory cytokines. MKKKs phosphorylate and activate MKKs. Then, MKKs phosphorylate and activate MAPKs including Erk1/2, JNK, and p38. Activated MAPKs phosphorylate various proteins and take part in regulation of inflammatory responses. DAMPs: damage-associated molecular; PAMPs: pathogen-associated molecular patterns; PRRs: pattern-recognition receptors; TLRs: Toll-like receptors; NLRs: NOD-like receptors; RLRs: retinoic acid-inducible gene-I-like receptors; MyD88: myeloid differentiation primary response protein-88; TAM: Tyro3, Axl, and Mer receptors; NF- $\kappa$ B: nuclear factor kappa-B; AP-1: activator protein-1; IRF3: interferon regulatory factor3; IL-1: interleukin-1; TNF: tumor necrosis factor; IKK: I $\kappa$ B kinase; JAK-STAT: Janus kinase-signal transducer and activator of transcription; IL-6: interleukin-6; MAPKs: mitogen-activated protein kinases; MKKK: MAPK kinase kinase; Erk1/2: extracellular signal-regulated kinase 1/2; MKK: MAPK kinase; JNK: c-Jun N-terminal kinases.

Primary inflammatory stimuli including microbial products and cytokines such as IL-1 $\beta$ , IL-6, and tumor necrosis factor (TNF)- $\alpha$ , interact with receptors including TLRs, IL-1 receptor (IL-1R), IL-6 receptor (IL-6R), and the TNF receptor (TNFR), respectively, and trigger important intracellular signaling pathways, including the MAPK, NF- $\kappa$ B and Janus kinase-signal transducer and activator of transcription (JAK-STAT) pathways [68–70]. Additionally, it is observed in many inflammatory disorders that ROS have the most significant effects in MAPK/AP-1, NF- $\kappa$ B, and JAK-STAT signaling pathways (Figure 3) [71–74].



**Figure 3.** Ultraviolet radiation (UVR), pollutants, and chemicals can activate surface receptors either directly or indirectly by activating reactive oxygen species (ROS). Thus, signaling pathways are activated, resulting in protein damage, cell apoptosis, DNA modification, lipid peroxidation, and production of proinflammatory mediators. UVR: ultraviolet radiation; NF- $\kappa$ B: nuclear factor kappa-B; AP-1: activator protein-1; IL-1R: interleukin-1 receptor; TNFR: tumor necrosis factor receptor; JAK-STAT: Janus kinase-signal transducer and activator of transcription; IL-6R: interleukin-6 receptor; MAPK: mitogen-activated protein kinase; EGFR: epidermal growth factor receptor; PGE2: Prostaglandin E2. Yellow, blue and black arrows indicate direction.

UVR is the most important extrinsic factor in melanogenesis regulation as well as ROS production and is considered to be the main stimulus for induced or acquired pigmentation, known as “tanning” [8,17,19,29]. The main cause of sunburn is UVB irradiation. Skin modulates the immune response to UVB irradiation, leading to neutrophil accumulation in the skin, and mast cells play a crucial role in UVB-induced skin inflammation by inducing a variety of proinflammatory mediators [75–77]. Skin UVB exposure significantly increases release of IL-8 and causes a modest increase in release of IL-1; however, other cytokines, such as IL-2, IL-4, IL-6, IL-10, IL-12, IL-13, TNF- $\alpha$  and IFN- $\gamma$ , remained unchanged. Exposure of skin to both UVA and UVB radiation causes an immediate rise in nicotinamide adenine dinucleotide phosphate (NADPH) oxidase level, resulting in ROS increase which then activates signaling pathways in keratinocytes in the epidermis and fibroblasts in the dermis, leading to the activation of inflammatory genes [78]. UVR can activate signaling pathways linked to surface receptors such as the EGF receptor [79,80], Prostaglandin E2 (PGE2) receptor [81,82], TNF- $\alpha$  receptor [83], and IL-1 receptor [84,85] either by causing receptor clustering [86,87] or by producing ROS, which then promotes phosphorylation and activation of downstream signaling pathway intermediates such as Erk1/2 and p38 [80,88–91].

Inflammatory mediators are produced after activation of intracellular signaling pathways by inflammatory stimuli. These have an important role in melanogenesis.




## 5. Inflammatory Cytokines

Allergens, pathogens, chemical stimuli, physical damage, and UVR can all lead to skin inflammation and inflammatory cytokines are predominantly released from immune cells, including monocytes, macrophages, and lymphocytes, which are considered closely related to skin pigmentation [92,93]. Pro- and anti-inflammatory cytokines facilitate and inhibit inflammation. Cytokines modulate immune response to infection or inflammation and regulate inflammation itself via a complex network of interactions. Th1, Th2 and Th17 cells [94] are the main types of Th cells in inflammation. Th1 cells secrete cytokines such as interferon- $\gamma$  (IFN- $\gamma$ ), TNF, and IL-2 and thus play an important role in cellular immune responses and Th2 cells play a central role in humoral immune responses by secreting IL-4, IL-5, IL-10, and IL-13 as well as other cytokines [95,96]. Th17 cells are a recently recognized subpopulation of T cells and produce IL-17, and IL-22, and IFN- $\gamma$ . Skin samples from atopic dermatitis (AD) patients contain numerous Th17 and Th2 cells, and cytokines produced from these cells (especially Th2 cells), which reduce the epidermis barrier function, elevate trans-epidermal water loss, and increase permeation of environmental irritants and allergens. Antigen-presenting cells (APCs) in the epidermis and Langerhans cells capture allergens with their dendrites extending upwards and process and present them to lymphocytes in the draining lymph nodes, thus stimulate naïve T-cells to be effector memory T cells.

Exposure to sensitized allergens evokes activation and accumulation of effector memory T cells to the exposure site, and these cells are stimulated to produce proinflammatory cytokines into the inflamed area. Atopic dermatitis patients are usually sensitized to food allergens, such as egg, and wheat, or to environmental allergens, such as house dust mites and pollen. These allergens existing everywhere can easily provoke inflammation in barrier-disrupted, easily permeable skin of atopic dermatitis patients. Atopic dirty neck is a well-known pigmentary condition considered to be due to post-inflammatory pigmentation/depigmentation. Th17 secretes IL-17, IL-6, IL-21, and IL-22, which participate in innate and acquired immunity. Psoriasis is another major inflammatory skin disease, where Th17 cells play an inevitable role in its pathogenesis. Treatment with biologics often leaves hypopigmented/hyperpigmented macules in the place of psoriasis plaques, which may be reflected by cytokine profiles in the psoriasis plaques. Treatment with UV (Psoralen-UVA (PUVA) or narrowband UVB) leaves strongly hyperpigmented macules without exception, probably due to the melanogenic function of UV.

## 6. Inflammatory Cytokines Induce Melanogenesis

Several studies have observed the involvement of local inflammatory factors in skin pigmentation regulation, which are presented in Table 1 (Figure 4). It has been revealed that IL-4 directly takes part in inhibiting melanogenesis in normal human epidermal melanocytes (NHEMs) by downregulating MITF, TYRP1, and TYRP2 expression through the JAK2-STAT6 signaling pathway [16]. IL-4 is mainly produced by Th2 cells. In chronic inflammation, basophils, eosinophils, and mast cells can also secrete IL-4 [97,98]. IL-4 plays a crucial role in IgE generation in hypersensitivity, inflammation induction, autoimmunity [99] and is also involved in the maintenance of Th2 lymphocytes and acts as an autocrine growth factor of differentiated Th2 cells [100]. It is hypothesized that skewed balance to Th1 is directly responsible for vitiligo development by changing IFN- $\gamma$ /IL-4, Tbet/Gata3 profiles in vitiligo patients compared to controls [101,102].



**Figure 4.** Involvement of inflammatory factors in melanogenesis. Inflammatory factors including IL-33, IL-18, and GM-CSF promote melanogenesis by stimulating the MAPK and PKA pathways either simultaneously or individually. In addition, PGE2, PGF2 $\alpha$ , and IL-1 $\alpha$  promote melanogenesis by stimulating cAMP-dependent pathway. In contrast, inflammatory factors such as IL-4, IL-13, IL-6, IL-17, IL-1 $\beta$ , TNF and INF- $\gamma$  inhibit melanogenesis by suppressing the PKA and MAPK, or JAK-STAT, or NF- $\kappa$ B and JNK pathways. IL-33: interleukin-33; IL-18: interleukin-18; PGE2: prostaglandin E2; PGF2 $\alpha$ : prostaglandin F2 $\alpha$ ; IL-1 $\alpha$ : interleukin-1 $\alpha$ ; GM-CSF: granulocyte-macrophage colony stimulating factor; IL-4: interleukin-4; IL-13: interleukin-13; IL-6: interleukin-6; IL-17: interleukin-17; IL-1 $\beta$ : interleukin-1 $\beta$ ; TNF: tumor necrosis factor; INF- $\gamma$ : interferons- $\gamma$ ; PKA: protein kinase A; MAPK: mitogen-activated protein kinase; JNK: c-Jun N-terminal kinases; JAK-STAT: Janus kinase-signal transducer and activator of transcription; NF- $\kappa$ B: nuclear factor kappa-B; TYR: tyrosinase; TRRP1: tyrosinase-related protein-1; TRRP2: tyrosinase-related protein-2. Red arrows indicates inhibition; green arrows indicate promotion.

IL-13 is a typical Th2 cell cytokine also secreted by CD4 cells, natural killer T cells, mast cells, basophils, eosinophils, and group 2 innate lymphoid cells (nuocytes) [103] sharing a receptor of similar structure and JAK2-STAT6 signaling pathway with IL-4 [104,105]. Recently, it has been found that activated human epidermal  $\gamma\delta$  T cells produce larger amounts of IL-13 than IL-4 [106] and enhance IL-13 production following treatment with Ginsenoside F1 (GF1) extracted from plant genus *Panax*. This plays a role in the skin-whitening effect of GF1 via effective suppression of tyrosinase and DCT, indicating that IL-13 could be a direct modulator of melanogenesis through the JAK2-STAT6 signaling pathway in skin disorders accompanied by enhanced T-cell responses, such as atopic dermatitis and vitiligo [107].

IL-17 is known to prevent bacterial and fungal infections [108] and can induce the release of large amounts of proinflammatory cytokines from a variety of cells, such as epithelial cells, endothelial cells, and fibroblasts [109], and its effects are vastly increased when cooperating with TNF [110]. IL-17, a proinflammatory cytokine, is produced primarily by Th17 cells, as well as by other immune cells, including neutrophils, natural killer cells, mast cells,  $\alpha\beta$  and  $\gamma\delta$ T cells, and group 3 innate lymphoid cells. Studies have revealed IL-17 can synergize with TNF to inhibit signaling pathway for melanogenesis, thereby inhibiting pigmentation [93]. For example, studies on psoriasis lesions showed that overexpression of IL-17 and TNF increase melanocyte number and cause a simultaneous decrease in pigmentation signaling. Neutralization of TNF and/or IL-17 with mAbs revealed that both can recover growth and pigment production of melanocytes, which may contribute to pigmentation changes associated with psoriasis treatment [93].

**Table 1.** Mechanisms of cytokines in the regulation of melanogenesis.

Cytokines	Main Source	Cells Used for Experiment	Impact on Melanogenesis	Mechanisms of Melanogenesis	Refs.
IL-4	Th-cells	Melanocytes	Inhibition	JAK2-STAT6 signaling pathway is used for downregulating the expression of MITF, TYRP1, TYRP2	[16]
IL-13	Th2	$\gamma\delta$ T-cell	Inhibition	Suppress tyrosinase and DCT through the JAK2-STAT6	[107]
IL-17	Epithelial cells, Endothelial cells, Fibroblasts	Melanocytes, Primary pooled human keratinocytes	Inhibition	TNF and IL-17 simultaneously inhibit melanin formation through PKA and MAPK signaling pathways	[93]
IL-33	Keratinocytes, Fibroblasts	Keratinocytes	Promotion	Activate p38 and PKA pathways and thus promote MITF, TYR, TYRP1, TYRP2 expression	[111,112]
IL-18	Monocytes/Macrophages, Keratinocytes	Melanocytes	Promotion	Increase the activity of tyrosinase and upregulate the expression Of TYRP1 and TYRP2	[113]
IL-1 $\alpha$	Langerhans cells, Melanocytes, keratinocytes	Primary melanocytes and swine skin	Promotion	Melanin deposition is increased when IL-1 $\alpha$ is combined with KGF	[114]
IL-1 $\beta$	Monocytes/Macrophages, Keratinocytes	Melanoma cell lines (LB2259-MEL and CP50-MEL)	Inhibition	Through the NF- $\kappa$ B and JNK pathways, it downregulates MITF-M expression	[115]
IL-6	Macrophages, T-cells, Adipocyte	Melanocytes	Inhibition	Tyrosinase activity is declined	[15]
PGE2 and PGF2 $\alpha$	Fibroblasts, Keratinocytes	Keratinocytes	Promotion	Activate cAMP-dependent pathway and stimulate melanocyte dendrite formation	[116]

Table 1. Cont.

Cytokines	Main Source	Cells Used for Experiment	Impact on Melanogenesis	Mechanisms of Melanogenesis	Refs.
INF- $\gamma$	T-cells, NK cells, NKT cells	B16F10	Inhibition	Block melanosomal maturation and upregulate STAT1 phosphorylation	[117,118]
TNF	Monocytes, Macrophages, Keratinocytes, Dendritic cells, Th1, Th17 and Th22	Melanocytes, Primary pooled human keratinocytes	Promotion	Stimulate PKA and MAPK signaling pathways with combination of IL17, and thus inhibit melanin formation	[93]
GM-CSF	Macrophages, Keratinocytes and Th cells	Melanocytes	Promotion	Stimulate MAPK pathway and promote melanocyte proliferation, melanin and synthesis	[119]

IL-33, interleukin-33; IL-18, interleukin-18; PGE2, prostaglandin E2; PGF2 $\alpha$ , prostaglandin F2 $\alpha$ ; IL-1 $\alpha$ , interleukin-1 $\alpha$ ; GM-CSF, granulocyte-macrophage colony stimulating factor; IL-4, interleukin-4; IL-13, interleukin-13; IL-6, interleukin-6; IL-17, interleukin-17; IL-1 $\beta$ , interleukin-1 $\beta$ ; TNF, tumor necrosis factor; IFN- $\gamma$ , interferons- $\gamma$ ; PKA, protein kinase A; MAPK, mitogen-activated protein kinase; JNK, c-Jun N-terminal kinases; JAK-STAT, Janus kinase-signal transducer and activator of transcription; NF- $\kappa$ B, nuclear factor kappa-B; TYR, tyrosinase; TRRP1, tyrosinase-related protein-1; TRRP2, tyrosinase-related protein-2; MITF, microphthalmia-associated transcription factor; NK cell, natural killer cell; NKT cell, natural killer T cell; Th, T helper cell.

Recent research has shown that IL-33 can improve melanin biosynthesis in NHEM, stimulate phosphorylation of p38 and CREB, and increase TYR, TYRP1 and TYRP2 (DCT) expression through MITF, resulting in augmentation of melanogenesis [112]. Abundant IL-33 mRNA is found in keratinocytes induced by UVB [111] and fibroblasts [120,121], suggesting injury on skin causes IL-33 release, resulting in skin pigmentation. Initially, IL-33 can induce mast cells to produce proinflammatory cytokines and chemokines (including monocyte chemoattractant protein-1 and macrophage inflammatory protein-1 $\alpha$ ) [122–126], leading to activation of macrophages [127–129], CD 4+T cells, basophils, dendritic cells and neutrophils [122,129–132], and likely promoting Th2-skewed skin inflammation, which is another indirect effect on pigmentation by IL-33 [133]. It has also been discovered that IL-33 activates and binds its receptor suppressing tumorigenicity 2 (ST2)L/IL-1RAcP on different types of cells, including mast cells and Th2 cells [134]. Then, IL-33 activates NF- $\kappa$ B, suggesting that it regulates outcome of diseases such as atopic dermatitis, which can trigger pigmentation [134].

IL-18 may also modulate both innate and adaptive immunity and its dysregulation can cause autoimmune or inflammatory disease, but some studies recently suggest its participation in pigmentation regulation. It has been observed that IL-18 promotes melanogenesis and upregulates TYRP1 and TYRP2 expression [113,118] by activating pathways and increasing cascade expressing MITF [113,118]. These results indicate the involvement of IL-18 in the regulation of pigmentation by regulating melanogenesis.

IL-1 is an important proinflammatory cytokine involved in initiating inflammation and immunological reactions [135], which contributes to increased tumor invasiveness, metastasis, and angiogenesis under chronic inflammatory conditions [115]. IL-1 is produced predominantly from monocytes and macrophages [136], although it is also produced by another types of cells, such as keratinocytes and fibroblasts [137]. IL-1 possesses two forms,

IL-1 $\alpha$  and IL-1 $\beta$  [138]. Although they share only 24% identity in protein sequence, IL-1 $\beta$  and IL-1 $\alpha$  fold in an extremely similar manner and bind to the same receptor—the type I and type II IL-1 receptor (IL-1RI, IL-1RII, respectively) [139]. The expression patterns of IL-1 $\alpha$  and IL-1 $\beta$  vary among cell types [140]. IL-1 $\alpha$  is predominantly produced by keratinocytes. Keratinocytes secrete both IL-1 $\alpha$  and IL-1 $\beta$  molecules *in vitro* [137], in addition to the expression of IL-1RI, IL-1RII [141]. The signal transduction is initiated by binding to IL-1RI [142] which can inhibit tyrosinase activity and melanogenesis [15,115] and also stimulates human fibroblasts to produce keratinocyte growth factor (KGF) [143]. IL-1 $\alpha$  released from keratinocytes has been shown to regulate the homeostasis of mammalian skin [144] and sufficient to trigger skin inflammation in humans [141,144] and produce more IL-1 $\alpha$  upon UVB exposure [145]. It is documented that KGF-induced TYR expression in primary melanocytes [114]. The combination of KGF and IL-1 $\alpha$  increases melanin deposition and is responsible for initial stage of human solar lentigines [114]. On the other hand, it was observed that the expression of MITF-M was inhibited after IL-1 $\beta$  treatment to a panel of melanoma cell lines. The inhibitory effects of IL-1 $\beta$  on melanogenesis were removed by inactivation of NF- $\kappa$ B and JNK pathways, suggesting IL-1 $\beta$  could inhibit melanogenesis by downregulating MITF-M through NF- $\kappa$ B and JNK pathways [115].

IL-6 is produced by numerous different cell types including keratinocytes, fibroblasts, and dermal endothelial cells and plays a critical role in regulating acute phase response, hematopoiesis, inflammation, metabolic control, liver regeneration, bone metabolism and cancer progression by promoting cell growth, survival, and differentiation [146]. It has been revealed in recent studies that the IL-6 cytokine elicits a dose-dependent decrease in tyrosinase activity, thus inhibiting melanocyte proliferation and melanogenesis [15].

PGE2 is the most abundant prostaglandin released by keratinocytes in response to UVR and inflammatory conditions such as wound healing and stimulates the formation of dendrites in melanocytes [147,148]. A wide range of physiological processes in the skin, including immune function, carcinogenesis, cutaneous barrier function, cell growth, and differentiation are regulated by PGE2 [149–152]. Biosynthesis of PGE2 is controlled by phospholipase A2 (PLA2), cyclooxygenase (COX), and prostaglandin E synthase (PGES) enzymes. PGE2 $\alpha$  is produced by fibroblasts and keratinocytes and stimulates melanocyte dendrite formation and tyrosinase activation, but not proliferation [151]. PGE2 binds to four distinct G-protein coupled receptors (EP1–4) and EP4 receptor signaling stimulates cAMP production in melanocytes. EP3 receptor-lowered basal cAMP levels suggest relative levels of activity of these receptors' control effects of PGE2 on cAMP in melanocytes that stimulates tyrosinase activity and proliferation [116,151].

IFN- $\gamma$  is also a common secretory cytokine in the skin and the most important endogenous mediator of immunity and inflammation [118]. Th1 lymphocytes, CD 8+ cytotoxic T lymphocytes and NK cells mainly release the proinflammatory cytokine IFN- $\gamma$  [153]. However, IFN- $\gamma$  is also secreted by other cells, including antigen-presenting cells, B cells and NKT cells [154–156]. A clear relationship between increased IFN- $\gamma$  signaling and a decreased number of stage III and IV melanosomes in the lesional skin of leprosy patients have been shown [117]. Specific IRF1 binding sites are in the promoter region of pigmentation-related genes such as TYR, TYRP2 (DCT). It is shown that IFN- $\gamma$  negatively regulates the expression of DCT and other pigmentation-related genes, providing strong evidence that IFN- $\gamma$  had a significant role in downregulating melanosome maturation [117]—i.e., skin hypopigmentation [117]. Two other crucial mediators of skin pigmentation,  $\alpha$ -MSH and TGF- $\beta$ , are involved in the protective tanning response and maintenance of melanocytes in an immature state [157,158] and are released during inflammation or UV exposure. Thus, Natarajan VT et al. proposed an  $\alpha\beta\gamma$ -cytokine regulatory model of melanogenesis [117]. Recent studies in various mouse models of vitiligo have demonstrated that local IFN- $\gamma$  accumulation produced by melanocyte-specific CD 8+ T cells plays an important role in skin depigmented spots [159,160] and also revealed increased IFN- $\gamma$  is essential for vitiligo pathogenesis by inducing apoptosis of melanocytes [161]. Moreover, IFN- $\gamma$  can upregulate STAT1 phosphorylation, and JAK1 inhibitors can suppress inhibiting

effect of IFN- $\gamma$ , thus contributing to melanogenesis. Other studies have also shown that IFN- $\gamma$  inhibits IL-18-induced melanogenesis [118].

TNF functions by binding to two different receptors, TNFR1/p55 and TNFR2/p75 [92], and is secreted mainly by monocytes and macrophages, and also by keratinocytes, dendritic cells, Th1, Th17 and Th22. TNF induces inflammation through activation of vascular endothelial cells and immune cells and plays a crucial role as a regulator of lymphoid tissue development by controlling apoptosis [92]. In several autoimmune diseases, increased levels of TNF have been found at inflammation sites, such as in the lesional skin of psoriasis patients [162]. Overexpression of IL-17 and TNF synergistically modulate cytokine expression, increase melanocyte number while suppressing pigmentation signaling [93]. Therapeutic neutralization of TNF and IL-17 with mAbs resulted in a rapid recovery of pigmentation-related gene expression in psoriasis lesions, indicating IL-17 and TNF can affect pigment production in melanocytes, which may contribute to the hypopigmentation associated with psoriasis. For example, after 24–48h treatment with both IL-17 and TNF of melanocytes, decreased levels of c-KIT, MC1R, MITF, and TYRP2 were observed [93]. Consequently, the levels of tyrosinase and melanin were significantly reduced [93]. It has been suggested the combination of IL-17 and TNF can inhibit melanogenesis through PKA and MAPK signaling pathways [92,93] and blocking TNF can lead to rapid restoration of pigmentation gene expression in psoriatic lesions, indicating that anti-TNF has potential for treating depigmented dermatosis [93]. IL-33 expression is induced by IL-17 and IFN- $\gamma$  in epidermal keratinocytes and induces melanogenesis, which creates a negative feedback loop in epidermal pigmentation [163,164].

GM-CSF may be a vital factor in the melanogenesis of the skin because of its role in promoting melanocyte proliferation and melanin synthesis. GM-CSF is produced by mononuclear macrophages, keratinocytes and Th cells [119]. It has been found GM-CSF synthesis and secretion by HaCaT cells increased cell proliferation in the wound healing process and contributes to wound healing and melanogenesis at the site of injury [165]. In addition, to predict the prognosis of transplantation of cultured autologous melanocytes (TCAM) in vitiligo patients, increased serum levels of GM-CSF could be used as the serum biomarkers [166].

## 7. Conclusions

From the currently available literature, little is known about the relationship between inflammation and pigmentation. We have attempted to show how diversified stimuli induce skin inflammation and how inflammatory pathways and inflammatory cytokines interact with each other, resulting in skin pigmentation. Our review has demonstrated that several inflammatory factors can promote or inhibit the melanogenesis in melanocytes through different mechanisms. It is widely accepted that the involvement of all types of inflammatory cells in the activation and release of inflammatory mediators makes it more complex to understand the regulatory network of inflammation and pigmentation. We believe that our review elucidating the role of inflammatory factors in the regulation of melanogenesis can provide new therapeutic perspectives for treating pigmentary disorders.

**Author Contributions:** M.R.H. wrote the manuscript and designed figures under the supervision of M.K. M.O. organized the main conceptual ideas of this review and T.M.A. accumulated all the sources of the table. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research did not receive any external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author.

**Conflicts of Interest:** The author declares no conflict of interest.

## Abbreviations

ER	Endoplasmic reticulum
PMEL17	Melanocytes lineage-specific antigen gp100
MART-1	Melanoma antigen recognized by T cell-1
GPNMB	Glycoprotein nonmetastatic melanoma protein b
TYR	Tyrosinase
TYRP1	Tyrosinase-related protein-1
TYRP2	Tyrosinase-related protein-2
BLOCs	Biogenesis of lysosome-related organelles complexes
AP-1	Activator protein 1
MLPH	Melanophilin
MYO5A	Myosin-Va
RAB27A	Ras-related protein Rab27A
MITF	Microphthalmia-associated transcription factor
DAMPs	Damage-associated molecular patterns
PAMPs	Pathogen-associated molecular patterns
PRRs	Pattern-recognition receptors
TLRs	Toll-like receptors;
NLRs	NOD-like receptors
RLRs	Retinoic acid-inducible gene-I-like receptors
MyD88	Myeloid differentiation primary response protein-88
TAM	Tyro3, Axl, and Mer receptors
NF- $\kappa$ B	Nuclear factor kappa-B
IRF	Interferon regulatory factor
IL	Interleukin
TNF	Tumor necrosis factor
IKK	I $\kappa$ B kinase
JAK-STAT	Janus kinase-signal transducer and activator of transcription
MAPKs	Mitogen-activated protein kinases
MKK	MAPK kinase
MKKK	MAPK kinase kinase
Erk1/2	Extracellular signal-regulated kinase $\frac{1}{2}$
JNK	c-Jun N-terminal kinases
UVR	Ultraviolet radiation
TNFR	Tumor necrosis factor receptor
EGFR	Epidermal growth factor receptor
PGE2	Prostaglandin E2
PGF2 $\alpha$	Prostaglandin F2 $\alpha$
GM-CSF	Granulocyte-macrophage colony stimulating factor
IFN- $\gamma$	Interferon- $\gamma$
PKA	Protein kinase A
NK cell	Natural killer cell
NKT cell	Natural killer T cell
Th	T helper cell
cAMP	Cyclic adenosine monophosphate
PAX3	Paired box protein Pax-3
SOX10	Sex-determining region Y-box 10
KIT	Stem cell growth factor receptor (CD117)
SCF	Stem cell growth factor
EDN	Endothelin
NGF	Nerve growth factor
HGF	Hepatocyte growth factor
$\alpha$ -MSH	$\alpha$ -Melanocyte-stimulating hormone

ACTH	Adrenocorticotrophic hormone
MLPH	melanophilin
CREB	cAMP responsive-element binding protein
SIK	Salt-inducible kinase
CRTC	CREB-regulated transcription coactivator
MITF-M	Melanocytes-specific MITF isoform

## References

1. Chuong, C.M.; Nickoloff, B.J.; Elias, P.M.; Goldsmith, L.A.; Macher, E.; Maderson, P.A.; Sundberg, J.P.; Tagami, H.; Plonka, P.M.; Thestrup-Pederson, K.; et al. What is the 'true' function of skin? *Exp. Dermatol.* **2002**, *11*, 159–187. [[CrossRef](#)]
2. Brenner, M.; Hearing, V.J. The protective role of melanin against UV damage in human skin. *Photochem. Photobiol.* **2008**, *84*, 539–549. [[CrossRef](#)]
3. Bonaventure, J.; Domingues, M.J.; Larue, L. Cellular and molecular mechanisms controlling the migration of melanocytes and melanoma cells. *Pigment. Cell Melanoma Res.* **2013**, *26*, 316–325. [[CrossRef](#)] [[PubMed](#)]
4. Jan Borovansky, P.A.R. *Melanins and Melanosomes Biosynthesis, Biogenesis, Physiological, and Pathological Functions*; Jan Borovansky, P.A.R., Ed.; Wiley-Blackwell: Weinheim, Germany, 2011; p. 429.
5. Lei, T.C.; Virador, V.; Yasumoto, K.; Vieira, W.D.; Toyofuku, K.; Hearing, V.J. Stimulation of melanoblast pigmentation by 8-methoxypsoralen: the involvement of microphthalmia-associated transcription factor, the protein kinase a signal pathway, and proteasome-mediated degradation. *J. Invest. Dermatol.* **2002**, *119*, 1341–1349. [[CrossRef](#)]
6. Sviderskaya, E.V.; Hill, S.P.; Balachandar, D.; Barsh, G.S.; Bennett, D.C. Agouti signaling protein and other factors modulating differentiation and proliferation of immortal melanoblasts. *Dev. Dyn.* **2001**, *221*, 373–379. [[CrossRef](#)] [[PubMed](#)]
7. Tolleson, W.H. Human melanocyte biology, toxicology, and pathology. *J. Environ. Sci. Health C Environ. Carcinog Ecotoxicol. Rev.* **2005**, *23*, 105–161. [[CrossRef](#)]
8. Lin, J.Y.; Fisher, D.E. Melanocyte biology and skin pigmentation. *Nature* **2007**, *445*, 843–850. [[CrossRef](#)]
9. Martin, S.F. Contact dermatitis: From pathomechanisms to immunotoxicology. *Exp. Dermatol.* **2012**, *21*, 382–389. [[CrossRef](#)] [[PubMed](#)]
10. Miller, L.S.; Cho, J.S. Immunity against *Staphylococcus aureus* cutaneous infections. *Nat. Rev. Immunol.* **2011**, *11*, 505–518. [[CrossRef](#)]
11. Behrends, U.; Peter, R.U.; Hintermeier-Knabe, R.; Eissner, G.; Holler, E.; Bornkamm, G.W.; Caughman, S.W.; Degitz, K. Ionizing radiation induces human intercellular adhesion molecule-1 in vitro. *J. Invest. Dermatol.* **1994**, *103*, 726–730. [[CrossRef](#)]
12. Fuchs, J.; Kern, H. Modulation of UV-light-induced skin inflammation by D-alpha-tocopherol and L-ascorbic acid: A clinical study using solar simulated radiation. *Free Radic. Biol. Med.* **1998**, *25*, 1006–1012. [[CrossRef](#)]
13. Bäsler, K.; Brandner, J.M. Tight junctions in skin inflammation. *Pflugers Arch.* **2017**, *469*, 3–14. [[CrossRef](#)]
14. Slominski, A.; Tobin, D.J.; Shibahara, S.; Wortsman, J. Melanin pigmentation in mammalian skin and its hormonal regulation. *Physiol. Rev.* **2004**, *84*, 1155–1228. [[CrossRef](#)]
15. Swope, V.B.; Abdel-Malek, Z.; Kassem, L.M.; Nordlund, J.J. Interleukins 1 alpha and 6 and tumor necrosis factor-alpha are paracrine inhibitors of human melanocyte proliferation and melanogenesis. *J. Invest. Dermatol.* **1991**, *96*, 180–185. [[CrossRef](#)] [[PubMed](#)]
16. Choi, H.; Choi, H.; Han, J.; Jin, S.H.; Park, J.Y.; Shin, D.W.; Lee, T.R.; Kim, K.; Lee, A.Y.; Noh, M. IL-4 inhibits the melanogenesis of normal human melanocytes through the JAK2-STAT6 signaling pathway. *J. Invest. Dermatol.* **2013**, *133*, 528–536. [[CrossRef](#)]
17. Park, H.Y.; Kosmadaki, M.; Yaar, M.; Gilchrist, B.A. Cellular mechanisms regulating human melanogenesis. *Cell Mol. Life Sci.* **2009**, *66*, 1493–1506. [[CrossRef](#)] [[PubMed](#)]
18. Fistarol, S.K.; Itin, P.H. Disorders of pigmentation. *J. Dtsch. Dermatol. Ges.* **2010**, *8*, 187–201. [[CrossRef](#)] [[PubMed](#)]
19. Tsatmali, M.; Ancans, J.; Thody, A.J. Melanocyte function and its control by melanocortin peptides. *J. Histochem. Cytochem.* **2002**, *50*, 125–133. [[CrossRef](#)]
20. Costin, G.E.; Hearing, V.J. Human skin pigmentation: Melanocytes modulate skin color in response to stress. *FASEB J.* **2007**, *21*, 976–994. [[CrossRef](#)]
21. Dessinioti, C.; Stratigos, A.J.; Rigopoulos, D.; Katsambas, A.D. A review of genetic disorders of hypopigmentation: Lessons learned from the biology of melanocytes. *Exp. Dermatol.* **2009**, *18*, 741–749. [[CrossRef](#)] [[PubMed](#)]
22. Imokawa, G. Melanocyte Activation Mechanisms and Rational Therapeutic Treatments of Solar Lentigos. *Int. J. Mol. Sci.* **2019**, *20*. [[CrossRef](#)] [[PubMed](#)]
23. Imokawa, G.; Ishida, K. Inhibitors of intracellular signaling pathways that lead to stimulated epidermal pigmentation: Perspective of anti-pigmenting agents. *Int. J. Mol. Sci.* **2014**, *15*, 8293–8315. [[CrossRef](#)]
24. Seiberg, M. Keratinocyte-melanocyte interactions during melanosome transfer. *Pigment. Cell Res.* **2001**, *14*, 236–242. [[CrossRef](#)]
25. Ito, S.; Wakamatsu, K. Chemistry of mixed melanogenesis—pivotal roles of dopaquinone. *Photochem. Photobiol.* **2008**, *84*, 582–592. [[CrossRef](#)] [[PubMed](#)]
26. Hara, M.; Yaar, M.; Byers, H.R.; Goukassian, D.; Fine, R.E.; Gonsalves, J.; Gilchrist, B.A. Kinesin participates in melanosomal movement along melanocyte dendrites. *J. Invest. Dermatol.* **2000**, *114*, 438–443. [[CrossRef](#)] [[PubMed](#)]



27. Byers, H.R.; Yaar, M.; Eller, M.S.; Jalbert, N.L.; Gilchrist, B.A. Role of cytoplasmic dynein in melanosome transport in human melanocytes. *J. Investig. Dermatol.* **2000**, *114*, 990–997. [[CrossRef](#)]
28. Schiaffino, M.V. Signaling pathways in melanosome biogenesis and pathology. *Int. J. Biochem. Cell Biol.* **2010**, *42*, 1094–1104. [[CrossRef](#)]
29. Yamaguchi, Y.; Hearing, V.J. Physiological factors that regulate skin pigmentation. *Biofactors* **2009**, *35*, 193–199. [[CrossRef](#)]
30. Hoashi, T.; Watabe, H.; Muller, J.; Yamaguchi, Y.; Vieira, W.D.; Hearing, V.J. MART-1 is required for the function of the melanosomal matrix protein PMEL17/GP100 and the maturation of melanosomes. *J. Biol. Chem.* **2005**, *280*, 14006–14016. [[CrossRef](#)]
31. Hoashi, T.; Sato, S.; Yamaguchi, Y.; Passeron, T.; Tamaki, K.; Hearing, V.J. Glycoprotein nonmetastatic melanoma protein b, a melanocytic cell marker, is a melanosome-specific and proteolytically released protein. *FASEB J.* **2010**, *24*, 1616–1629. [[CrossRef](#)]
32. Zhang, P.; Liu, W.; Zhu, C.; Yuan, X.; Li, D.; Gu, W.; Ma, H.; Xie, X.; Gao, T. Silencing of GPNMB by siRNA inhibits the formation of melanosomes in melanocytes in a MITF-independent fashion. *PLoS ONE* **2012**, *7*, e42955. [[CrossRef](#)]
33. Schallreuter, K.U.; Kothari, S.; Chavan, B.; Spencer, J.D. Regulation of melanogenesis—controversies and new concepts. *Exp. Dermatol.* **2008**, *17*, 395–404. [[CrossRef](#)]
34. Park, H.Y.; Perez, J.M.; Laursen, R.; Hara, M.; Gilchrist, B.A. Protein kinase C-beta activates tyrosinase by phosphorylating serine residues in its cytoplasmic domain. *J. Biol. Chem.* **1999**, *274*, 16470–16478. [[CrossRef](#)]
35. Kondo, T.; Hearing, V.J. Update on the regulation of mammalian melanocyte function and skin pigmentation. *Expert Rev. Dermatol.* **2011**, *6*, 97–108. [[CrossRef](#)]
36. Kuroda, T.S.; Ariga, H.; Fukuda, M. The actin-binding domain of Slac2-a/melanophilin is required for melanosome distribution in melanocytes. *Mol. Cell Biol.* **2003**, *23*, 5245–5255. [[CrossRef](#)] [[PubMed](#)]
37. Van Den Bossche, K.; Naeyaert, J.M.; Lambert, J. The quest for the mechanism of melanin transfer. *Traffic* **2006**, *7*, 769–778. [[CrossRef](#)] [[PubMed](#)]
38. Vachtenheim, J.; Borovanský, J. “Transcription physiology” of pigment formation in melanocytes: Central role of MITF. *Exp. Dermatol.* **2010**, *19*, 617–627. [[CrossRef](#)] [[PubMed](#)]
39. Liu, J.J.; Fisher, D.E. Lighting a path to pigmentation: Mechanisms of MITF induction by UV. *Pigment. Cell Melanoma Res.* **2010**, *23*, 741–745. [[CrossRef](#)] [[PubMed](#)]
40. Park, H.Y. Biology of melanocytes. In *Fitzpatrick's Dermatology in General Medicine*; McGraw-Hill: New York, NY, USA, 2008; pp. 591–608.
41. Steingrímsson, E.; Copeland, N.G.; Jenkins, N.A. Melanocytes and the microphthalmia transcription factor network. *Annu. Rev. Genet.* **2004**, *38*, 365–411. [[CrossRef](#)]
42. Mujahid, N.; Liang, Y.; Murakami, R.; Choi, H.G.; Dobry, A.S.; Wang, J.; Suita, Y.; Weng, Q.Y.; Allouche, J.; Kemeny, L.V.; et al. A UV-Independent Topical Small-Molecule Approach for Melanin Production in Human Skin. *Cell Rep.* **2017**, *19*, 2177–2184. [[CrossRef](#)]
43. Yamaguchi, Y.; Brenner, M.; Hearing, V.J. The regulation of skin pigmentation. *J. Biol. Chem.* **2007**, *282*, 27557–27561. [[CrossRef](#)]
44. Bennett, D.C.; Lamoreux, M.L. The color loci of mice—a genetic century. *Pigment. Cell Res.* **2003**, *16*, 333–344. [[CrossRef](#)]
45. Yuan, X.H.; Jin, Z.H. Paracrine regulation of melanogenesis. *Br. J. Dermatol.* **2018**, *178*, 632–639. [[CrossRef](#)] [[PubMed](#)]
46. Swope, V.B.; Abdel-Malek, Z.A. MC1R: Front and Center in the Bright Side of Dark Eumelanin and DNA Repair. *Int. J. Mol. Sci.* **2018**, *19*. [[CrossRef](#)] [[PubMed](#)]
47. Grando, S.A.; Pittelkow, M.R.; Schallreuter, K.U. Adrenergic and cholinergic control in the biology of epidermis: Physiological and clinical significance. *J. Investig. Dermatol.* **2006**, *126*, 1948–1965. [[CrossRef](#)]
48. Besmer, P.; Murphy, J.E.; George, P.C.; Qiu, F.H.; Bergold, P.J.; Lederman, L.; Snyder, H.W., Jr.; Brodeur, D.; Zuckerman, E.E.; Hardy, W.D. A new acute transforming feline retrovirus and relationship of its oncogene v-kit with the protein kinase gene family. *Nature* **1986**, *320*, 415–421. [[CrossRef](#)]
49. Yarden, Y.; Kuang, W.J.; Yang-Feng, T.; Coussens, L.; Munemitsu, S.; Dull, T.J.; Chen, E.; Schlessinger, J.; Francke, U.; Ullrich, A. Human proto-oncogene c-kit: A new cell surface receptor tyrosine kinase for an unidentified ligand. *Embo J.* **1987**, *6*, 3341–3351. [[CrossRef](#)]
50. Lim, X.; Nusse, R. Wnt signaling in skin development, homeostasis, and disease. *Cold Spring Harb. Perspect. Biol.* **2013**, *5*. [[CrossRef](#)]
51. Dorsky, R.I.; Moon, R.T.; Raible, D.W. Control of neural crest cell fate by the Wnt signalling pathway. *Nature* **1998**, *396*, 370–373. [[CrossRef](#)]
52. Takeda, K.; Yasumoto, K.; Takada, R.; Takada, S.; Watanabe, K.; Udono, T.; Saito, H.; Takahashi, K.; Shibahara, S. Induction of melanocyte-specific microphthalmia-associated transcription factor by Wnt-3a. *J. Biol. Chem.* **2000**, *275*, 14013–14016. [[CrossRef](#)] [[PubMed](#)]
53. Jin, E.J.; Erickson, C.A.; Takada, S.; Burrus, L.W. Wnt and BMP signaling govern lineage segregation of melanocytes in the avian embryo. *Dev. Biol.* **2001**, *233*, 22–37. [[CrossRef](#)]
54. Dunn, K.J.; Brady, M.; Ochsenbauer-Jambor, C.; Snyder, S.; Incao, A.; Pavan, W.J. WNT1 and WNT3a promote expansion of melanocytes through distinct modes of action. *Pigment. Cell Res.* **2005**, *18*, 167–180. [[CrossRef](#)]
55. Dorsky, R.I.; Raible, D.W.; Moon, R.T. Direct regulation of nacre, a zebrafish MITF homolog required for pigment cell formation, by the Wnt pathway. *Genes Dev.* **2000**, *14*, 158–162. [[PubMed](#)]

56. Flaherty, K.T.; Hodi, F.S.; Fisher, D.E. From genes to drugs: Targeted strategies for melanoma. *Nat. Rev. Cancer* **2012**, *12*, 349–361. [[CrossRef](#)]
57. Widlund, H.R.; Horstmann, M.A.; Price, E.R.; Cui, J.; Lessnick, S.L.; Wu, M.; He, X.; Fisher, D.E. Beta-catenin-induced melanoma growth requires the downstream target Microphthalmia-associated transcription factor. *J. Cell Biol.* **2002**, *158*, 1079–1087. [[CrossRef](#)]
58. Benson, H.A.E.; Grice, J.E.; Mohammed, Y.; Namjoshi, S.; Roberts, M.S. Topical and Transdermal Drug Delivery: From Simple Potions to Smart Technologies. *Curr. Drug Deliv.* **2019**, *16*, 444–460. [[CrossRef](#)] [[PubMed](#)]
59. Sander, C.S.; Chang, H.; Hamm, F.; Elsner, P.; Thiele, J.J. Role of oxidative stress and the antioxidant network in cutaneous carcinogenesis. *Int. J. Dermatol.* **2004**, *43*, 326–335. [[CrossRef](#)] [[PubMed](#)]
60. Kohen, R. Skin antioxidants: Their role in aging and in oxidative stress—New approaches for their evaluation. *Biomed Pharm.* **1999**, *53*, 181–192. [[CrossRef](#)]
61. Trouba, K.J.; Hamadeh, H.K.; Amin, R.P.; Germolec, D.R. Oxidative stress and its role in skin disease. *Antioxid. Redox Signal.* **2002**, *4*, 665–673. [[CrossRef](#)] [[PubMed](#)]
62. Gudkov, A.V.; Komarova, E.A. p53 and the Carcinogenicity of Chronic Inflammation. *Cold Spring Harb. Perspect. Med.* **2016**, *6*. [[CrossRef](#)]
63. Brusselle, G.; Bracke, K. Targeting immune pathways for therapy in asthma and chronic obstructive pulmonary disease. *Ann. Am. Thorac. Soc.* **2014**, *11* (Suppl. 5), S322–S328. [[CrossRef](#)]
64. Takeuchi, O.; Akira, S. Pattern recognition receptors and inflammation. *Cell* **2010**, *140*, 805–820. [[CrossRef](#)] [[PubMed](#)]
65. Janeway, C.A., Jr.; Medzhitov, R. Innate immune recognition. *Annu. Rev. Immunol.* **2002**, *20*, 197–216. [[CrossRef](#)] [[PubMed](#)]
66. Adib-Conquy, M.; Cavaillon, J.M. Stress molecules in sepsis and systemic inflammatory response syndrome. *FEBS Lett.* **2007**, *581*, 3723–3733. [[CrossRef](#)]
67. Rubartelli, A.; Lotze, M.T. Inside, outside, upside down: Damage-associated molecular-pattern molecules (DAMPs) and redox. *Trends Immunol.* **2007**, *28*, 429–436. [[CrossRef](#)]
68. Hendrayani, S.F.; Al-Harbi, B.; Al-Ansari, M.M.; Silva, G.; Aboussekhra, A. The inflammatory/cancer-related IL-6/STAT3/NF- $\kappa$ B positive feedback loop includes AUF1 and maintains the active state of breast myofibroblasts. *Oncotarget* **2016**, *7*, 41974–41985. [[CrossRef](#)]
69. Kyriakis, J.M.; Avruch, J. Mammalian mitogen-activated protein kinase signal transduction pathways activated by stress and inflammation. *Physiol. Rev.* **2001**, *81*, 807–869. [[CrossRef](#)]
70. Henríquez-Olguín, C.; Altamirano, F.; Valladares, D.; López, J.R.; Allen, P.D.; Jaimovich, E. Altered ROS production, NF- $\kappa$ B activation and interleukin-6 gene expression induced by electrical stimulation in dystrophic mdx skeletal muscle cells. *Biochim. Biophys. Acta* **2015**, *1852*, 1410–1419. [[CrossRef](#)] [[PubMed](#)]
71. Kim, A.L.; Labasi, J.M.; Zhu, Y.; Tang, X.; McClure, K.; Gabel, C.A.; Athar, M.; Bickers, D.R. Role of p38 MAPK in UVB-induced inflammatory responses in the skin of SKH-1 hairless mice. *J. Investig. Dermatol.* **2005**, *124*, 1318–1325. [[CrossRef](#)]
72. Wang, F.; Lee, E.; Lowes, M.A.; Haider, A.S.; Fuentes-Duculan, J.; Abello, M.V.; Chamian, F.; Cardinale, I.; Krueger, J.G. Prominent production of IL-20 by CD68+/CD11c+ myeloid-derived cells in psoriasis: Gene regulation and cellular effects. *J. Investig. Dermatol.* **2006**, *126*, 1590–1599. [[CrossRef](#)] [[PubMed](#)]
73. Wolk, K.; Kunz, S.; Witte, E.; Friedrich, M.; Asadullah, K.; Sabat, R. IL-22 increases the innate immunity of tissues. *Immunity* **2004**, *21*, 241–254. [[CrossRef](#)]
74. Dhar, A.; Young, M.R.; Colburn, N.H. The role of AP-1, NF- $\kappa$ B and ROS/NOS in skin carcinogenesis: The JB6 model is predictive. *Mol. Cell Biochem.* **2002**, *234–235*, 185–193. [[CrossRef](#)]
75. Endoh, I.; Di Girolamo, N.; Hampartzoumian, T.; Cameron, B.; Geczy, C.L.; Tedla, N. Ultraviolet B irradiation selectively increases the production of interleukin-8 in human cord blood-derived mast cells. *Clin. Exp. Immunol.* **2007**, *148*, 161–167. [[CrossRef](#)]
76. Valencia, A.; Kochevar, I.E. Nox1-based NADPH oxidase is the major source of UVA-induced reactive oxygen species in human keratinocytes. *J. Investig. Dermatol.* **2008**, *128*, 214–222. [[CrossRef](#)] [[PubMed](#)]
77. Beak, S.M.; Lee, Y.S.; Kim, J.A. NADPH oxidase and cyclooxygenase mediate the ultraviolet B-induced generation of reactive oxygen species and activation of nuclear factor- $\kappa$ B in HaCaT human keratinocytes. *Biochimie* **2004**, *86*, 425–429. [[CrossRef](#)]
78. Glady, A.; Tanaka, M.; Moniaga, C.S.; Yasui, M.; Hara-Chikuma, M. Involvement of NADPH oxidase 1 in UVB-induced cell signaling and cytotoxicity in human keratinocytes. *Biochem. Biophys. Rep.* **2018**, *14*, 7–15. [[CrossRef](#)] [[PubMed](#)]
79. Xu, Y.; Shao, Y.; Zhou, J.; Voorhees, J.J.; Fisher, G.J. Ultraviolet irradiation-induces epidermal growth factor receptor (EGFR) nuclear translocation in human keratinocytes. *J. Cell Biochem.* **2009**, *107*, 873–880. [[CrossRef](#)] [[PubMed](#)]
80. Huang, R.P.; Wu, J.X.; Fan, Y.; Adamson, E.D. UV activates growth factor receptors via reactive oxygen intermediates. *J. Cell Biol.* **1996**, *133*, 211–220. [[CrossRef](#)] [[PubMed](#)]
81. Tober, K.L.; Thomas-Ahner, J.M.; Kusewitt, D.F.; Oberyszyn, T.M. Effects of UVB on E prostanoid receptor expression in murine skin. *J. Investig. Dermatol.* **2007**, *127*, 214–221. [[CrossRef](#)]
82. Soontrapa, K.; Honda, T.; Sakata, D.; Yao, C.; Hirata, T.; Hori, S.; Matsuoka, T.; Kita, Y.; Shimizu, T.; Kabashima, K.; et al. Prostaglandin E2-prostaglandin E receptor subtype 4 (EP4) signaling mediates UV irradiation-induced systemic immunosuppression. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 6668–6673. [[CrossRef](#)] [[PubMed](#)]
83. Muthusamy, V.; Piva, T.J. The UV response of the skin: A review of the MAPK, NF $\kappa$ B and TNF $\alpha$  signal transduction pathways. *Arch. Dermatol. Res.* **2010**, *302*, 5–17. [[CrossRef](#)] [[PubMed](#)]

84. Wan, Y.S.; Wang, Z.Q.; Voorhees, J.; Fisher, G. EGF receptor crosstalks with cytokine receptors leading to the activation of c-Jun kinase in response to UV irradiation in human keratinocytes. *Cell Signal*. **2001**, *13*, 139–144. [[CrossRef](#)]
85. Blanton, R.A.; Kupper, T.S.; McDougall, J.K.; Dower, S. Regulation of interleukin 1 and its receptor in human keratinocytes. *Proc. Natl. Acad. Sci. USA* **1989**, *86*, 1273–1277. [[CrossRef](#)]
86. López-Camarillo, C.; Ocampo, E.A.; Casamichana, M.L.; Pérez-Plasencia, C.; Alvarez-Sánchez, E.; Marchat, L.A. Protein kinases and transcription factors activation in response to UV-radiation of skin: Implications for carcinogenesis. *Int. J. Mol. Sci.* **2012**, *13*, 142–172. [[CrossRef](#)]
87. Rosette, C.; Karin, M. Ultraviolet light and osmotic stress: Activation of the JNK cascade through multiple growth factor and cytokine receptors. *Science* **1996**, *274*, 1194–1197. [[CrossRef](#)]
88. Peus, D.; Meves, A.; Vasa, R.A.; Beyerle, A.; O'Brien, T.; Pittelkow, M.R. H<sub>2</sub>O<sub>2</sub> is required for UVB-induced EGF receptor and downstream signaling pathway activation. *Free Radic. Biol. Med.* **1999**, *27*, 1197–1202. [[CrossRef](#)]
89. Madson, J.G.; Hansen, L.A. Multiple mechanisms of ErbB2 action after ultraviolet irradiation of the skin. *Mol. Carcinog.* **2007**, *46*, 624–628. [[CrossRef](#)]
90. Son, Y.; Cheong, Y.K.; Kim, N.H.; Chung, H.T.; Kang, D.G.; Pae, H.O. Mitogen-Activated Protein Kinases and Reactive Oxygen Species: How Can ROS Activate MAPK Pathways? *J. Signal. Transduct.* **2011**, *2011*, 792639. [[CrossRef](#)]
91. Finkel, T. Signal transduction by reactive oxygen species. *J. Cell Biol.* **2011**, *194*, 7–15. [[CrossRef](#)]
92. Grine, L.; DeJager, L.; Libert, C.; Vandenbroucke, R.E. An inflammatory triangle in psoriasis: TNF, type I IFNs and IL-17. *Cytokine Growth Factor Rev.* **2015**, *26*, 25–33. [[CrossRef](#)] [[PubMed](#)]
93. Wang, C.Q.F.; Akalu, Y.T.; Suarez-Farinas, M.; Gonzalez, J.; Mitsui, H.; Lowes, M.A.; Orlow, S.J.; Manga, P.; Krueger, J.G. IL-17 and TNF synergistically modulate cytokine expression while suppressing melanogenesis: Potential relevance to psoriasis. *J. Investig. Dermatol.* **2013**, *133*, 2741–2752. [[CrossRef](#)]
94. Mosmann, T.R.; Sad, S. The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol. Today* **1996**, *17*, 138–146. [[CrossRef](#)]
95. O'Garra, A. Cytokines induce the development of functionally heterogeneous T helper cell subsets. *Immunity* **1998**, *8*, 275–283. [[CrossRef](#)]
96. Reiner, S.L.; Seder, R.A. Dealing from the evolutionary pawnshop: How lymphocytes make decisions. *Immunity* **1999**, *11*, 1–10. [[CrossRef](#)]
97. Barata, L.T.; Ying, S.; Meng, Q.; Barkans, J.; Rajakulasingam, K.; Durham, S.R.; Kay, A.B. IL-4- and IL-5-positive T lymphocytes, eosinophils, and mast cells in allergen-induced late-phase cutaneous reactions in atopic subjects. *J. Allergy Clin. Immunol.* **1998**, *101*, 222–230. [[CrossRef](#)]
98. Min, B.; Prout, M.; Hu-Li, J.; Zhu, J.; Jankovic, D.; Morgan, E.S.; Urban, J.F., Jr.; Dvorak, A.M.; Finkelman, F.D.; LeGros, G.; et al. Basophils produce IL-4 and accumulate in tissues after infection with a Th2-inducing parasite. *J. Exp. Med.* **2004**, *200*, 507–517. [[CrossRef](#)] [[PubMed](#)]
99. Imran, M.; Laddha, N.C.; Dwivedi, M.; Mansuri, M.S.; Singh, J.; Rani, R.; Gokhale, R.S.; Sharma, V.K.; Marfatia, Y.S.; Begum, R. Interleukin-4 genetic variants correlate with its transcript and protein levels in patients with vitiligo. *Br. J. Dermatol.* **2012**, *167*, 314–323. [[CrossRef](#)] [[PubMed](#)]
100. Salgame, P.; Abrams, J.S.; Clayberger, C.; Goldstein, H.; Convit, J.; Modlin, R.L.; Bloom, B.R. Differing lymphokine profiles of functional subsets of human CD4 and CD8 T cell clones. *Science* **1991**, *254*, 279–282. [[CrossRef](#)]
101. Basak, P.Y.; Adiloglu, A.K.; Ceyhan, A.M.; Tas, T.; Akkaya, V.B. The role of helper and regulatory T cells in the pathogenesis of vitiligo. *J. Am. Acad. Dermatol.* **2009**, *60*, 256–260. [[CrossRef](#)]
102. Nouri-Koupaee, A.; Mansouri, P.; Jahانبini, H.; Sanati, M.H.; Jadali, Z. Differential expression of mRNA for T-bet and GATA-3 transcription factors in peripheral blood mononuclear cells of patients with vitiligo. *Clin. Exp. Dermatol.* **2015**, *40*, 735–740. [[CrossRef](#)] [[PubMed](#)]
103. Rael, E.L.; Lockey, R.F. Interleukin-13 signaling and its role in asthma. *World Allergy Organ J.* **2011**, *4*, 54–64. [[CrossRef](#)]
104. Nishimura, Y.; Nitto, T.; Inoue, T.; Node, K. IL-13 attenuates vascular tube formation via JAK2-STAT6 pathway. *Circ. J.* **2008**, *72*, 469–475. [[CrossRef](#)] [[PubMed](#)]
105. Renauld, J.C. New insights into the role of cytokines in asthma. *J. Clin. Pathol.* **2001**, *54*, 577–589. [[CrossRef](#)]
106. Kim, K.; Han, J.; Lee, T.R.; Shin, D.W.; Chang, H.; Cho, A.R.; Choi, S.J.; Jo, S.J.; Kwon, O. Comparative Analysis of Human Epidermal and Peripheral Blood  $\gamma\delta$  T Cell Cytokine Profiles. *Ann. Dermatol.* **2014**, *26*, 308–313. [[CrossRef](#)]
107. Han, J.; Lee, E.; Kim, E.; Yeom, M.H.; Kwon, O.; Yoon, T.H.; Lee, T.R.; Kim, K. Role of epidermal  $\gamma\delta$  T-cell-derived interleukin 13 in the skin-whitening effect of Ginsenoside F1. *Exp. Dermatol.* **2014**, *23*, 860–862. [[CrossRef](#)]
108. Speeckaert, R.; Lambert, J.; Grine, L.; Van Gele, M.; De Schepper, S.; van Geel, N. The many faces of interleukin-17 in inflammatory skin diseases. *Br. J. Dermatol.* **2016**, *175*, 892–901. [[CrossRef](#)]
109. Volpe, E.; Servant, N.; Zollinger, R.; Bogiatzi, S.I.; Hupé, P.; Barillot, E.; Soumelis, V. A critical function for transforming growth factor-beta, interleukin 23 and proinflammatory cytokines in driving and modulating human T(H)-17 responses. *Nat. Immunol.* **2008**, *9*, 650–657. [[CrossRef](#)] [[PubMed](#)]
110. Liang, S.C.; Tan, X.Y.; Luxenberg, D.P.; Karim, R.; Dunussi-Joannopoulos, K.; Collins, M.; Fouser, L.A. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J. Exp. Med.* **2006**, *203*, 2271–2279. [[CrossRef](#)] [[PubMed](#)]

111. Meehansan, J.; Komine, M.; Tsuda, H.; Tominaga, S.; Ohtsuki, M. Ultraviolet B irradiation induces the expression of IL-33 mRNA and protein in normal human epidermal keratinocytes. *J. Dermatol. Sci.* **2012**, *65*, 72–74. [[CrossRef](#)] [[PubMed](#)]
112. Zhou, J.; Song, J.; Ping, F.; Shang, J. Enhancement of the p38 MAPK and PKA signaling pathways is associated with the pro-melanogenic activity of Interleukin 33 in primary melanocytes. *J. Dermatol. Sci.* **2014**, *73*, 110–116. [[CrossRef](#)]
113. Zhou, J.; Shang, J.; Song, J.; Ping, F. Interleukin-18 augments growth ability of primary human melanocytes by PTEN inactivation through the AKT/NF- $\kappa$ B pathway. *Int. J. Biochem. Cell Biol.* **2013**, *45*, 308–316. [[CrossRef](#)] [[PubMed](#)]
114. Chen, N.; Hu, Y.; Li, W.H.; Eisinger, M.; Seiberg, M.; Lin, C.B. The role of keratinocyte growth factor in melanogenesis: A possible mechanism for the initiation of solar lentigines. *Exp. Dermatol.* **2010**, *19*, 865–872. [[CrossRef](#)] [[PubMed](#)]
115. Kholmanskikh, O.; van Baren, N.; Brasseur, F.; Ottaviani, S.; Vanacker, J.; Arts, N.; van der Bruggen, P.; Coulie, P.; De Plaen, E. Interleukins 1 $\alpha$  and 1 $\beta$  secreted by some melanoma cell lines strongly reduce expression of MITF-M and melanocyte differentiation antigens. *Int. J. Cancer* **2010**, *127*, 1625–1636. [[CrossRef](#)] [[PubMed](#)]
116. Scott, G.; Leopardi, S.; Printup, S.; Malhi, N.; Seiberg, M.; Lapoint, R. Proteinase-activated receptor-2 stimulates prostaglandin production in keratinocytes: Analysis of prostaglandin receptors on human melanocytes and effects of PGE2 and PGF2 $\alpha$  on melanocyte dendricity. *J. Investig. Dermatol.* **2004**, *122*, 1214–1224. [[CrossRef](#)] [[PubMed](#)]
117. Natarajan, V.T.; Ganju, P.; Singh, A.; Vijayan, V.; Kirty, K.; Yadav, S.; Puntambekar, S.; Bajaj, S.; Dani, P.P.; Kar, H.K.; et al. IFN- $\gamma$  signaling maintains skin pigmentation homeostasis through regulation of melanosome maturation. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 2301–2306. [[CrossRef](#)] [[PubMed](#)]
118. Zhou, J.; Ling, J.; Wang, Y.; Shang, J.; Ping, F. Cross-talk between interferon-gamma and interleukin-18 in melanogenesis. *J. Photochem. Photobiol. B* **2016**, *163*, 133–143. [[CrossRef](#)]
119. Videira, I.F.; Moura, D.F.; Magina, S. Mechanisms regulating melanogenesis. *An. Bras. Dermatol.* **2013**, *88*, 76–83. [[CrossRef](#)]
120. Arend, W.P.; Palmer, G.; Gabay, C. IL-1, IL-18, and IL-33 families of cytokines. *Immunol. Rev.* **2008**, *223*, 20–38. [[CrossRef](#)]
121. Byrne, S.N.; Beaugie, C.; O’Sullivan, C.; Leighton, S.; Halliday, G.M. The immune-modulating cytokine and endogenous Alarmin interleukin-33 is upregulated in skin exposed to inflammatory UVB radiation. *Am. J. Pathol.* **2011**, *179*, 211–222. [[CrossRef](#)]
122. Ali, S.; Huber, M.; Kollewe, C.; Bischoff, S.C.; Falk, W.; Martin, M.U. IL-1 receptor accessory protein is essential for IL-33-induced activation of T lymphocytes and mast cells. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 18660–18665. [[CrossRef](#)]
123. Allakhverdi, Z.; Smith, D.E.; Comeau, M.R.; Delespesse, G. Cutting edge: The ST2 ligand IL-33 potently activates and drives maturation of human mast cells. *J. Immunol.* **2007**, *179*, 2051–2054. [[CrossRef](#)] [[PubMed](#)]
124. Moulin, D.; Donzé, O.; Talabot-Ayer, D.; Mézin, F.; Palmer, G.; Gabay, C. Interleukin (IL)-33 induces the release of pro-inflammatory mediators by mast cells. *Cytokine* **2007**, *40*, 216–225. [[CrossRef](#)] [[PubMed](#)]
125. Theoharides, T.C.; Zhang, B.; Kempuraj, D.; Tagen, M.; Vasiadi, M.; Angelidou, A.; Alysandratos, K.D.; Kalogeromitros, D.; Asadi, S.; Stavrianeas, N.; et al. IL-33 augments substance P-induced VEGF secretion from human mast cells and is increased in psoriatic skin. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 4448–4453. [[CrossRef](#)] [[PubMed](#)]
126. Pushparaj, P.N.; Tay, H.K.; H’Ng, S.C.; Pitman, N.; Xu, D.; McKenzie, A.; Liew, F.Y.; Melendez, A.J. The cytokine interleukin-33 mediates anaphylactic shock. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 9773–9778. [[CrossRef](#)]
127. Kurowska-Stolarska, M.; Stolarski, B.; Kewin, P.; Murphy, G.; Corrigan, C.J.; Ying, S.; Pitman, N.; Mirchandani, A.; Rana, B.; van Rooijen, N.; et al. IL-33 amplifies the polarization of alternatively activated macrophages that contribute to airway inflammation. *J. Immunol.* **2009**, *183*, 6469–6477. [[CrossRef](#)] [[PubMed](#)]
128. Ohno, T.; Oboki, K.; Kajiwara, N.; Morii, E.; Aozasa, K.; Flavell, R.A.; Okumura, K.; Saito, H.; Nakae, S. Caspase-1, caspase-8, and calpain are dispensable for IL-33 release by macrophages. *J. Immunol.* **2009**, *183*, 7890–7897. [[CrossRef](#)]
129. Schmitz, J.; Owyang, A.; Oldham, E.; Song, Y.; Murphy, E.; McClanahan, T.K.; Zurawski, G.; Moshrefi, M.; Qin, J.; Li, X.; et al. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity* **2005**, *23*, 479–490. [[CrossRef](#)] [[PubMed](#)]
130. Hueber, A.J.; Alves-Filho, J.C.; Asquith, D.L.; Michels, C.; Millar, N.L.; Reilly, J.H.; Graham, G.J.; Liew, F.Y.; Miller, A.M.; McInnes, I.B. IL-33 induces skin inflammation with mast cell and neutrophil activation. *Eur. J. Immunol.* **2011**, *41*, 2229–2237. [[CrossRef](#)]
131. Suzukawa, M.; Iikura, M.; Koketsu, R.; Nagase, H.; Tamura, C.; Komiyama, A.; Nakae, S.; Matsushima, K.; Ohta, K.; Yamamoto, K.; et al. An IL-1 cytokine member, IL-33, induces human basophil activation via its ST2 receptor. *J. Immunol.* **2008**, *181*, 5981–5989. [[CrossRef](#)]
132. Rank, M.A.; Kobayashi, T.; Kozaki, H.; Bartemes, K.R.; Squillace, D.L.; Kita, H. IL-33-activated dendritic cells induce an atypical TH2-type response. *J. Allergy Clin. Immunol.* **2009**, *123*, 1047–1054. [[CrossRef](#)]
133. Hsu, C.L.; Chhiba, K.D.; Krier-Burris, R.; Hosakoppal, S.; Berdnikovs, S.; Miller, M.L.; Bryce, P.J. Allergic inflammation is initiated by IL-33-dependent crosstalk between mast cells and basophils. *PLoS ONE* **2020**, *15*, e0226701. [[CrossRef](#)] [[PubMed](#)]
134. Cevikbas, F.; Steinhoff, M. IL-33: A novel danger signal system in atopic dermatitis. *J. Investig. Dermatol.* **2012**, *132*, 1326–1329. [[CrossRef](#)]
135. Dinarello, C.A. Interleukin-1. *Rev. Infect. Dis.* **1984**, *6*, 51–95. [[CrossRef](#)]
136. Kampschmidt, R.F. The numerous postulated biological manifestations of interleukin-1. *J. Leukoc. Biol.* **1984**, *36*, 341–355. [[CrossRef](#)] [[PubMed](#)]
137. Kupper, T.S. Immune and inflammatory processes in cutaneous tissues. Mechanisms and speculations. *J. Clin. Investig.* **1990**, *86*, 1783–1789. [[CrossRef](#)] [[PubMed](#)]

138. Auron, P.E.; Webb, A.C.; Rosenwasser, L.J.; Mucci, S.F.; Rich, A.; Wolff, S.M.; Dinarello, C.A. Nucleotide sequence of human monocyte interleukin 1 precursor cDNA. *Proc. Natl. Acad. Sci. USA* **1984**, *81*, 7907–7911. [[CrossRef](#)]
139. Sims, J.E.; March, C.J.; Cosman, D.; Widmer, M.B.; MacDonald, H.R.; McMahan, C.J.; Grubin, C.E.; Wignall, J.M.; Jackson, J.L.; Call, S.M.; et al. cDNA expression cloning of the IL-1 receptor, a member of the immunoglobulin superfamily. *Science* **1988**, *241*, 585–589. [[CrossRef](#)]
140. March, C.J.; Mosley, B.; Larsen, A.; Cerretti, D.P.; Braedt, G.; Price, V.; Gillis, S.; Henney, C.S.; Kronheim, S.R.; Grabstein, K.; et al. Cloning, sequence and expression of two distinct human interleukin-1 complementary DNAs. *Nature* **1985**, *315*, 641–647. [[CrossRef](#)]
141. Groves, R.W.; Sherman, L.; Mizutani, H.; Dower, S.K.; Kupper, T.S. Detection of interleukin-1 receptors in human epidermis. Induction of the type II receptor after organ culture and in psoriasis. *Am. J. Pathol.* **1994**, *145*, 1048–1056.
142. Martin, M.U.; Wesche, H. Summary and comparison of the signaling mechanisms of the Toll/interleukin-1 receptor family. *Biochim. Biophys. Acta* **2002**, *1592*, 265–280. [[CrossRef](#)]
143. Tang, A.; Gilchrist, B.A. Regulation of keratinocyte growth factor gene expression in human skin fibroblasts. *J. Dermatol. Sci.* **1996**, *11*, 41–50. [[CrossRef](#)]
144. Groves, R.W.; Ross, E.; Barker, J.N.; Camp, R.D.; MacDonald, D.M. Effect of in vivo interleukin-1 on adhesion molecule expression in normal human skin. *J. Investig. Dermatol.* **1992**, *98*, 384–387. [[CrossRef](#)] [[PubMed](#)]
145. Kondo, S.; Sauder, D.N.; Kono, T.; Galley, K.A.; McKenzie, R.C. Differential modulation of interleukin-1 alpha (IL-1 alpha) and interleukin-1 beta (IL-1 beta) in human epidermal keratinocytes by UVB. *Exp. Dermatol.* **1994**, *3*, 29–39. [[CrossRef](#)]
146. Hirano, T.; Ishihara, K.; Hibi, M. Roles of STAT3 in mediating the cell growth, differentiation and survival signals relayed through the IL-6 family of cytokine receptors. *Oncogene* **2000**, *19*, 2548–2556. [[CrossRef](#)] [[PubMed](#)]
147. Pentland, A.P.; Mahoney, M.; Jacobs, S.C.; Holtzman, M.J. Enhanced prostaglandin synthesis after ultraviolet injury is mediated by endogenous histamine stimulation. A mechanism for irradiation erythema. *J. Clin. Investig.* **1990**, *86*, 566–574. [[CrossRef](#)]
148. Pentland, A.P.; Mahoney, M.G. Keratinocyte prostaglandin synthesis is enhanced by IL-1. *J. Investig. Dermatol.* **1990**, *94*, 43–46. [[CrossRef](#)]
149. Fosslie, E. Molecular pathology of cyclooxygenase-2 in neoplasia. *Ann. Clin. Lab. Sci.* **2000**, *30*, 3–21.
150. Pentland, A.P.; Needleman, P. Modulation of keratinocyte proliferation in vitro by endogenous prostaglandin synthesis. *J. Clin. Investig.* **1986**, *77*, 246–251. [[CrossRef](#)]
151. Scott, G.; Jacobs, S.; Leopardi, S.; Anthony, F.A.; Learn, D.; Malaviya, R.; Pentland, A. Effects of PGF2alpha on human melanocytes and regulation of the FP receptor by ultraviolet radiation. *Exp. Cell Res.* **2005**, *304*, 407–416. [[CrossRef](#)]
152. Holtzman, M.J.; Zhang, V.; Hussain, H.; Roswit, W.T.; Wilson, J.D. Prostaglandin H synthase and lipoxygenase gene families in the epithelial cell barrier. *Ann. N. Y. Acad. Sci.* **1994**, *744*, 58–77. [[CrossRef](#)]
153. Bach, E.A.; Aguet, M.; Schreiber, R.D. The IFN gamma receptor: A paradigm for cytokine receptor signaling. *Annu. Rev. Immunol.* **1997**, *15*, 563–591. [[CrossRef](#)]
154. Carnaud, C.; Lee, D.; Donnars, O.; Park, S.H.; Beavis, A.; Koezuka, Y.; Bendelac, A. Cutting edge: Cross-talk between cells of the innate immune system: NKT cells rapidly activate NK cells. *J. Immunol.* **1999**, *163*, 4647–4650.
155. Frucht, D.M.; Fukao, T.; Bogdan, C.; Schindler, H.; O’Shea, J.J.; Koyasu, S. IFN-gamma production by antigen-presenting cells: Mechanisms emerge. *Trends Immunol.* **2001**, *22*, 556–560. [[CrossRef](#)]
156. Flaishon, L.; Hershkoviz, R.; Lantner, F.; Lider, O.; Alon, R.; Levo, Y.; Flavell, R.A.; Shachar, I. Autocrine secretion of interferon gamma negatively regulates homing of immature B cells. *J. Exp. Med.* **2000**, *192*, 1381–1388. [[CrossRef](#)] [[PubMed](#)]
157. Nishimura, E.K.; Suzuki, M.; Igras, V.; Du, J.; Lonning, S.; Miyachi, Y.; Roes, J.; Beermann, F.; Fisher, D.E. Key roles for transforming growth factor beta in melanocyte stem cell maintenance. *Cell Stem Cell* **2010**, *6*, 130–140. [[CrossRef](#)] [[PubMed](#)]
158. Abdel-Malek, Z.; Scott, M.C.; Suzuki, I.; Tada, A.; Im, S.; Lamoreux, L.; Ito, S.; Barsh, G.; Hearing, V.J. The melanocortin-1 receptor is a key regulator of human cutaneous pigmentation. *Pigment. Cell Res.* **2000**, *13* (Suppl. 8) (Suppl. 8), 156–162. [[CrossRef](#)]
159. Harris, J.E.; Harris, T.H.; Weninger, W.; Wherry, E.J.; Hunter, C.A.; Turka, L.A. A mouse model of vitiligo with focused epidermal depigmentation requires IFN- $\gamma$  for autoreactive CD8<sup>+</sup> T-cell accumulation in the skin. *J. Investig. Dermatol.* **2012**, *132*, 1869–1876. [[CrossRef](#)]
160. Gregg, R.K.; Nichols, L.; Chen, Y.; Lu, B.; Engelhard, V.H. Mechanisms of spatial and temporal development of autoimmune vitiligo in tyrosinase-specific TCR transgenic mice. *J. Immunol.* **2010**, *184*, 1909–1917. [[CrossRef](#)] [[PubMed](#)]
161. Yang, L.; Wei, Y.; Sun, Y.; Shi, W.; Yang, J.; Zhu, L.; Li, M. Interferon-gamma Inhibits Melanogenesis and Induces Apoptosis in Melanocytes: A Pivotal Role of CD8<sup>+</sup> Cytotoxic T Lymphocytes in Vitiligo. *Acta Derm. Venereol.* **2015**, *95*, 664–670. [[CrossRef](#)] [[PubMed](#)]
162. Kristensen, M.; Chu, C.Q.; Eedy, D.J.; Feldmann, M.; Brennan, F.M.; Breathnach, S.M. Localization of tumour necrosis factor-alpha (TNF-alpha) and its receptors in normal and psoriatic skin: Epidermal cells express the 55-kD but not the 75-kD TNF receptor. *Clin. Exp. Immunol.* **1993**, *94*, 354–362. [[CrossRef](#)] [[PubMed](#)]
163. Meehansan, J.; Tsuda, H.; Komine, M.; Tominaga, S.; Ohtsuki, M. Regulation of IL-33 expression by IFN- $\gamma$  and tumor necrosis factor- $\alpha$  in normal human epidermal keratinocytes. *J. Investig. Dermatol.* **2012**, *132*, 2593–2600. [[CrossRef](#)] [[PubMed](#)]
164. Meehansan, J.; Komine, M.; Tsuda, H.; Karakawa, M.; Tominaga, S.; Ohtsuki, M. Expression of IL-33 in the epidermis: The mechanism of induction by IL-17. *J. Dermatol. Sci.* **2013**, *71*, 107–114. [[CrossRef](#)] [[PubMed](#)]

- 
165. Park, K.C.; Kim, D.S.; Kim, H.J.; Seo, K.I.; Kim, K.H.; Chung, J.H.; Eun, H.C.; Jung, H.C. Growth related secretion and production of GM-CSF by epithelial cell line. *J. Dermatol. Sci.* **2001**, *25*, 53–58. [[CrossRef](#)]
  166. Wu, X.G.; Hong, W.S.; Xu, A. GM-CSF: A possible prognostic serum biomarker of vitiligo patients' considered for transplantation treatment with cultured autologous melanocytes: A pilot study. *J. Eur. Acad. Dermatol. Venereol.* **2016**, *30*, 1409–1411. [[CrossRef](#)] [[PubMed](#)]