

SHORT REPORT

Open Access

Evaluation of coding-independent functions of the transcribed bovine aromatase pseudogene *CYP19P1*

Marina Chwalisz and Rainer Fürbass *

Abstract

Background: *CYP19A1* encodes the aromatase which catalyzes the final reaction of estrogen biosynthesis. The bovine genome also contains a non-coding copy of *CYP19A1*, the transcribed pseudogene *CYP19P1*. Whereas *CYP19A1* is transcribed in all estrogen-producing tissues, mainly in the placenta and gonads, the *CYP19P1* transcript so far was detected in the placenta. Strikingly, one sequence segment of both transcripts exhibits an exceptional high identity of 98%, which implies selective pressure and suggests some kind of function. Only recently, indeed, coding-independent functions of several transcribed pseudogenes were reported. Therefore, we analyzed *CYP19P1* and *CYP19A1* transcripts with the aim to detect clues for gene-pseudogene interference.

Findings: The *CYP19P1* transcript was first examined *in silico* for the presence of microRNA coding sequences and microRNA targets. Further, to identify tissues where *CYP19P1* and *CYP19A1* transcripts are co-expressed, as a pre-requisite for transcript interference, expression profiling was performed in a variety of bovine tissues. Our *in silico* analyses did neither reveal potential microRNA coding sequences, nor microRNA targets. Co-expression of the *CYP19* loci was demonstrated in placental cotyledons and granulosa cells of dominant follicles. However, in granulosa cells of dominant follicles the concentration of *CYP19P1* mRNA was very low compared to *CYP19A1* mRNA.

Conclusions: *CYP19P1* and *CYP19A1* transcripts might interfere in placental cotyledons. However, in granulosa cells of dominant follicles relevant interference between gene and pseudogene transcripts is unlikely to occur because of the very low *CYP19P1/CYP19A1* transcript ratio.

Keywords: *CYP19A1*, Placenta, Granulosa, Gene-pseudogene interference

Findings

Background

CYP19A1 encodes the aromatase which catalyzes the conversion of androgens to estrogens. During an earlier screening of a bovine placental cDNA library for *CYP19A1* clones, we also isolated cDNA clones of a homologous pseudogene, *CYP19P1* [1,2]. Gene and pseudogene transcripts exhibit major differences due to the loss of several exons in the *CYP19P1* transcript. Furthermore, numerous mutations in the pseudogene sequence produced multiple translational stop codons in all reading frames and thereby abrogated its protein-coding function. Strikingly, however, mutations are unevenly distributed. A sequence segment

of 177 bp corresponding to exon 5 of *CYP19A1* is highly conserved, showing 98% identity, compared to 89% sequence identity in general [2]. Both *CYP19* loci are located on the same strand of chromosome 10, being separated by 20 kb of genomic DNA [3,4].

Pseudogenes for long were considered as defunct copies of functional genes. However, recent evidence suggests that some pseudogenes might exert coding-independent functions [5,6]. Interestingly, gene-pseudogene interference was demonstrated by selective knock-down of the *ABCC6P1* pseudogene, which led to a decreased transcription of its *ABCC6* parent gene [7]. This recent evidence prompted us to take up again our analysis of the *CYP19P1* pseudogene with the aim to detect clues of gene-pseudogene interference. To this end, transcripts were searched *in silico* for microRNA-coding sequences and microRNA targets. Further, expression profiles of

* Correspondence: fuerbass@fhn-dummerstorf.de
Leibniz Institute for Farm Animal Biology (FBN), Wilhelm-Stahl-Allee 2,
Dummerstorf 18196, Germany

CYP19A1 and *CYP19P1* were analyzed in a variety of bovine tissues.

Materials and methods

Samples from placentas, ovarian granulosa cells, fetal ovaries, endometria, adrenal glands and livers were collected from slaughtered cows in a local abattoir. Tissue samples were stored in RNAlater (Qiagen, Hilden, Germany) at -20°C. Granulosa cells were frozen in liquid nitrogen and stored at -80°C. Placental cotyledons and caruncles were separated manually. Despite careful separation, caruncle samples might contain traces of cotyledonary cells. Ovarian dominant and pre-ovulatory follicles collected before and after the LH-surge, respectively, were identified as described in [8]. Follicles were punctured with 18G needles and granulosa cells were aspirated. Total RNA was prepared using the NucleoSpin RNA II Kit (Macherey-Nagel, Düren, Germany), according

to the supplier's protocol. This procedure included on-column DNaseI digestion to remove traces of DNA. RNA was quantified in a NanoDrop 1000 spectrophotometer (PeQLab, Erlangen, Germany). RNA integrity was confirmed by denaturing agarose gel electrophoresis.

The abundance of *CYP19P1* and *CYP19A1* transcripts was measured by quantitative reverse transcription PCR (qPCR). For normalization purposes, the *RPLP0* transcript encoding a ribosomal large subunit protein was also measured. RNA (100 ng) was reverse transcribed in a 25 µl reaction volume using a mixture of random hexameric and oligo dT primers (Roche, Mannheim, Germany) and M-MLV reverse transcriptase (Promega, Mannheim, Germany). cDNA was purified with the High Pure PCR Product Purification Kit (Roche). For subsequent real time PCR, cDNA was amplified in a 12 µl reaction volume with the SensiFast SYBR No-ROX Kit (Bioline, Luckenwalde, Germany) using following primer pairs:




Figure 1 Expression of *CYP19P1* and *CYP19A1* in bovine tissues. **A)** Quantitative reverse transcription PCR analysis of *CYP19P1* and *CYP19A1* transcripts. Concentration values were normalized using *RPLP0* as an internal control. The abundance of the *RPLP0* transcript in the bovine tissues is indicated in the inserted diagram. The results are shown as means \pm SEM of $n = 3$ independent experiments. Analyzed tissues were placental cotyledons (Cot) and caruncles (Car), fetal ovaries (fOv) and granulosa cells from dominant and pre-ovulatory follicles (dGC and pGC, respectively). *CYP19A1* and *CYP19P1* transcripts were not detected in pGC, adrenal glands, endometria and livers. **B)** Agarose gels confirming the generation of only the expected products during the qPCR analysis shown in the panel A. The products of qPCR reactions including reverse transcription are indicated below the gels by white boxes labeled + RT. Control PCR reactions without prior reverse transcription (indicated by black boxes labeled -RT) did not yield products. Lanes labeled with +, - and M contain positive PCR controls (PCR products from cloned cDNAs), negative PCR controls (PCR reactions without templates) and molecular weight markers, respectively.


CYP19P1_for, 5'-TCATTACAACGCATCCCCAGGTTGA-3'/**CYP19P1_rev**, 5'-CTAGGTCCATGACGGGCTGGTATCA-3', **CYP19A1_for**, 5'-GGATCGGCAGTGCCTGCAATTACTA-3'/**CYP19A1_rev**, 5'-ATGCCGATGAACTGCAACCCAAGTT-3' and **RPLP0_for**, 5'-TGGTTACCCAACCGTCGCATCTGTA-3'/**RPLP0_rev**, 5'-CACAAAGGCAGATGGATCAGCCAAG-3' (Sigma-Aldrich, Taufkirchen, Germany). The primer pairs were designed to flank intronic sequences of genomic DNA. The expected products from cDNA were 156, 174 and 140 bp in length, respectively. The amplification and quantification of resulting PCR products were performed in a Light-Cycler 480 instrument (Roche) under following cycling conditions: Pre-incubation at 95°C for 5 min, followed by 40 cycles of denaturation at 95°C for 20 s, annealing at 60°C for 15 s, and extension at 72°C for

15 s, and single point fluorescence acquisition at 75°C for 10 s to avoid quantifying primer artifacts. The generation of only the expected products was confirmed by melting curve analysis and agarose gel electrophoresis, which did not reveal PCR products from genomic DNA (Figure 1B). External standard curves were generated by co-amplification of various dilutions of cloned PCR products (5×10^{-12} to 5×10^{-16} g DNA/reaction) with the corresponding primer pairs.

The statistical analyses were performed with the Sigma Plot 12.0 Analysis System (Jandel Scientific, San Rafael, CA, USA).

Results and discussion

The remarkably high conservation of *CYP19P1* in the exon 5-homologous sequence segment (henceforth referred to



as P1-exon 5) implies selective pressure. Hence, it might well be that the pseudogene exerts a coding-independent biological activity via the P1-exon 5. Because pseudogene-derived small interfering RNAs were shown to regulate gene expression in mouse oocytes [9], we examined if also the *CYP19P1* transcript encodes microRNAs. We performed *in silico* analyses of the P1-exon 5 using the free software RNAfold [10] to detect putative stable pre-microRNA hairpins. However, no such structures were predicted by the program. The cellular abundance of the tumor suppressor gene *PTEN* transcript was found to be regulated by the transcript of the highly homologous *PTENP1* pseudogene via competition for microRNA binding [11]. To evaluate if the *CYP19* gene pair could also interfere this way, we searched the P1-exon 5 for microRNA target sites using the MIRANDA software [12]. However, probable targets of known microRNAs were not found. Other pseudogenes are transcribed in an anti-sense orientation and lead to silencing of their parent genes by translational interference [5]. However, this mode of action can not apply to *CYP19P1* and *CYP19A1* which are both encoded by the same strand of chromosome 10 [4] and hence are transcribed in the same orientation.

For possible interference, *CYP19P1* and *CYP19A1* transcripts need to be co-expressed. So far, expression of *CYP19P1* has not been studied in bovine tissues other than placenta. To evaluate co-expression as a prerequisite for transcript interference, we performed qPCR to measure the abundance of *CYP19P1* and *CYP19A1* transcripts in a variety of bovine tissues. These included granulosa cells from dominant follicles and placental cotyledons, which were known to express *CYP19A1* [13]. Further, granulosa cells from pre-ovulatory follicles, fetal ovaries, placental caruncles, endometrium, adrenal gland and liver were also analyzed, to find out whether the tissue-specificity of *CYP19A1* expression is retained in *CYP19P1*. The results of the qPCR experiments are presented in Figure 1A. As expected, cotyledons and granulosa cells from dominant follicles expressed considerable amounts of the *CYP19A1* transcript. In contrast, the *CYP19A1* transcript concentration was low or undetectable in the remaining tissue samples. Likewise, the *CYP19P1* transcript was found in cotyledons and granulosa cells from dominant follicles, however, at a very low concentration. The ratio *CYP19P1*/*CYP19A1* of mean transcript concentrations was 0.16 and 0.016, respectively. Considering these results *CYP19P1* and *CYP19A1* transcripts might interfere in placental cotyledons, although the underlying mechanism remains unclear. On the other hand it is unlikely that the *CYP19P1* transcript plays a major role in the regulation of *CYP19A1* expression in granulosa cells of dominant follicles. Interestingly, however, in granulosa cells from pre-ovulatory follicles *CYP19P1* expression is shut down, just as known from *CYP19A1*, as a consequence of the pre-ovulatory

LH-surge [14,8]. The apparent congruent tissue-specificity of *CYP19* gene and pseudogene could be explained by promoter conservation. A genomic DNA sequence covering the entire *CYP19A1* and *CYP19P1* loci is present in the database [GenBank:NW_003104282]. We defined the genomic sequence immediately upstream of the transcribed *CYP19P1* sequence as potential pseudogene promoter and compared it with promoters of *CYP19A1* by BLAST analysis. Thereby we found, that 483 bp of the proximal *CYP19A1* promoter 2 ([15]; [GenBank:Z69242]) were duplicated during pseudogene genesis (Figure 2). We have verified the sequence by sequencing cloned PCR products (data not shown). Gene and pseudogene promoters exhibit 83% sequence identity. Although binding motifs of GATA and SF1 transcription factors are retained, the putative pseudogene promoter has lost the functional TATA box, which might explain the low promoter activity and the use of an alternative transcription start site.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MC implementation, analysis of data and preparation of the manuscript. RF initiation of the study, analysis of data, preparation of the manuscript. Both authors read and approved the final manuscript.

Acknowledgements

We thank Dr. R. M. Brunner and F. Hadlich for providing assistance with *in silico* analyses, Maren Anders and Veronica Schreiter for excellent technical assistance. This work was supported by a grant from the Deutsche Forschungsgemeinschaft (DFG).

Received: 30 September 2013 Accepted: 13 June 2014

Published: 20 June 2014

References

1. Vanselow J, Fürbass R: Novel aromatase transcripts from bovine placenta contain repeated sequence motifs. *Gene* 1995, **154**:281–286.
2. Fürbass R, Vanselow J: An aromatase pseudogene is transcribed in the bovine placenta. *Gene* 1995, **154**:287–292.
3. Goldammer T, Guérin G, Brunner RM, Vanselow J, Fürbass R, Schwerin M: Chromosomal mapping of the bovine aromatase gene (*CYP19*) and an aromatase pseudogene to chromosome 10 and syntenic group U5. *Mam Genome* 1994, **5**:822–823.
4. Brunner RM, Goldammer T, Fürbass R, Vanselow J, Schwerin M: Genomic organization of the bovine aromatase encoding gene and a homologous pseudogene as revealed by DNA fiber FISH. *Cytogenet Cell Genet* 1998, **82**:37–40.
5. Pink R, Wicks K, Caley D, Punch E, Jacobs L, Carter D: Pseudogenes: pseudo-functional or key regulators in health and disease? *RNA* 2011, **17**:792–798.
6. Pei B, Sisu C, Frankish A, Howald C, Habegger L, Mu X, Harte R, Balasubramanian S, Tanzer A, Diekhans M, Reymond A, Hubbard T, Harrow J, Gerstein M: The GENCODE pseudogene resource. *Genome Biol* 2012, **13**:R51.
7. Pihler AP, Hellum M, Wenzel J, Kaminski E, Haug K, Kierulf P, Kaminski W: The human ABC transporter pseudogene family: evidence for transcription and gene-pseudogene interference. *BMC Genomics* 2008, **9**:165.
8. Nimz M, Spitschak M, Schneider F, Fürbass R, Vanselow J: Down-regulation of genes encoding steroidogenic enzymes and hormone receptors in late preovulatory follicles of the cow coincides with an accumulation of intrafollicular steroids. *Domest Anim Endocrinol* 2009, **37**:45–54.
9. Tam O, Aravin A, Stein P, Girard A, Murchison E, Cheloufi S, Hodges E, Anger M, Sachidanandam R, Schultz R, Hannon G: Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 2008, **453**:534–539.

10. Gruber A, Lorenz R, Bernhard S, Neuböck R, Hofacker I: **The Vienna RNA websuite.** *Nucleic Acids Res* 2008, **36**:W70–W74.
11. Poliseno L, Salmena L, Zhang J, Carver B, Haveman W, Pandolfi P: **A coding-independent function of gene and pseudogene mRNAs regulates tumor biology.** *Nature* 2010, **465**:1033–1040.
12. Enright A, John B, Gaul U, Tuschl T, Sander C, Marks D: **MicroRNA targets in *Drosophila*.** *Genome Biol* 2003, **5**:R1.
13. Vanselow J, Fürbass R, Rehbock F, Klautschek G, Schwerin M: **Cattle and sheep use different promoters to direct the expression of aromatase cytochrome P450 encoding gene, *Cyp 19*, during pregnancy.** *Domest Anim Endocrinol* 2004, **27**:99–114.
14. Voss AK, Fortune JE: **Levels of messenger ribonucleic acid for cytochrome P450 17alpha-hydroxylase and P450 aromatase in preovulatory bovine follicles decrease after the luteinizing hormone surge.** *Endocrinology* 1993, **132**:2239–2245.
15. Fürbass R, Kalbe C, Vanselow J: **Tissue-specific expression of the bovine aromatase encoding gene uses multiple transcriptional start sites and alternative first exons.** *Endocrinology* 1997, **138**:2813–2819.

doi:10.1186/1756-0500-7-378

Cite this article as: Chwalisz and Fürbass: Evaluation of coding-independent functions of the transcribed bovine aromatase pseudogene *CYP19P1*. *BMC Research Notes* 2014 **7**:378.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

