


# Potential Value of Haptic Feedback in Minimally Invasive Surgery for Deep Endometriosis

Surgical Innovation  
2020, Vol. 27(6) 623–632  
© The Author(s) 2020



Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: 10.1177/1553350620944267  
[journals.sagepub.com/home/sri](https://journals.sagepub.com/home/sri)



Stijn L. Vlek, MD<sup>1</sup> , Rens Burm, MSc<sup>2,3</sup>, Tim M. Govers, PhD<sup>2</sup>, Michel P. H. Vleugels, MD, PhD<sup>4</sup>, Jurriaan B. Tuynman, MD, PhD<sup>1</sup>, and Velja Mijatovic, MD, PhD<sup>5</sup>

## Abstract

**Introduction.** Laparoscopic treatment of deep endometriosis (DE) is associated with intra- and post-operative morbidity. New technological developments, such as haptic feedback in laparoscopic instruments, could reduce the rate of complications. The aim of this study was to assess the room for improvement and potential cost-effectiveness of haptic feedback instruments in laparoscopic surgery. **Methods.** To assess the potential value of haptic feedback, a decision analytical model was constructed. Complications that could be related to the absence of haptic feedback were included in the model. Costs of complications were based on the additional length of hospital stay, operating time, outpatient visits, reinterventions, and/or conversions to laparotomy. The target population consists of women who are treated for DE in the Netherlands. A headroom analysis was performed to estimate the maximum value of haptic feedback in case it would be able to prevent all selected intra- and post-operative complications. **Results.** A total of 9.7 intraoperative and 47.0 post-operative complications are expected in the cohort of 636 patients annually treated for DE in the Netherlands. Together, these complications cause an additional length of hospital stay of 432.1 days, 10.2 additional outpatient visits, 73.9 reinterventions, and 4.2 conversions. Most consequences are related to post-operative complications. The total additional annual costs due to complications were €436 623, amounting to €687 additional costs per patient. **Discussion.** This study demonstrated that the potential value for improvement in DE laparoscopic surgery by using haptic feedback instruments is considerable, mostly caused by the potential prevention of major post-operative complications.

## Keywords

gynaecological laparoscopy, colorectal surgery, the business of surgery

## Introduction

Endometriosis is a benign disorder defined as the presence of endometrial-like tissue outside the uterus which induces an inflammatory response. This response is accompanied by adhesions, fibrosis, angiogenesis, neuronal infiltration, and anatomical distortion, resulting in pelvic pain, pelvic organ dysfunction, and infertility. It affects 5-10% of women of reproductive age.<sup>1</sup> In women with endometriosis, the prevalence of involvement of the gastrointestinal tract is 8-12%. The prevalence of involvement of the urinary tract is lesser and amounts 1-5.5%.<sup>2-4</sup> Surgical treatment can be used as first-line therapy in patients with (imminent) organ dysfunction or initiated after failed medical therapies including nonsteroidal anti-inflammatory drugs and hormonal treatments. The majority of these surgical procedures are performed laparoscopically in the Netherlands; the remainder is performed by laparotomy or robot-assisted surgery.<sup>5</sup>

Surgical treatment of deep endometriosis (DE) involving the bowel or urinary tract can be challenging due

to the extensive inflammation and fibrosis destroying normal anatomy. The normal embryological planes present in normal tissue are no longer visible in severe endometriosis. Therefore, endometriosis surgery is associated with a risk of major complications.<sup>6,7</sup> For treatment of DE of the urinary tract, intraoperative accidental injuries of the ureters and bleeding are reported. Major post-operative complications include anastomotic leakage,

<sup>1</sup>Endometriosis Center, Department of Surgery, Amsterdam University Medical Center, Amsterdam, the Netherlands

<sup>2</sup>MedValue, Nijmegen, the Netherlands

<sup>3</sup>NewCompliance, The Hague, the Netherlands

<sup>4</sup>HCB Asistel Moraira, Spain

<sup>5</sup>Endometriosis Center, Department of Reproductive Medicine, Amsterdam University Medical Center, Amsterdam, the Netherlands

## Corresponding Author:

Stijn L. Vlek, Endometriosis Center, Department of Surgery, Amsterdam University Medical Center, Location VUmc, De Boelelaan 1117, Amsterdam, the Netherlands.  
Email: [s.vlek@amsterdamumc.nl](mailto:s.vlek@amsterdamumc.nl)

ureteral fistulae, and vesicovaginal fistulae. Conversions or re-interventions are prevalent in 3-6.7% and 3.9% of cases, respectively.<sup>8</sup> Treatment of DE of the bowel is associated with iatrogenic damage, bleeding, bowel perforation, and intraoperative anastomotic leakage. Post-operative complications include anastomotic leakage, late bowel perforation, intra-abdominal infection, and rectovaginal fistulae.<sup>9</sup> Intraoperative complications lead to significantly increased post-operative morbidity.

Laparoscopic surgery has 3 general limitations: reduced hand-eye coordination, reduced depth perception, and reduced haptic feedback.<sup>10</sup> Reduced haptic feedback leads to grasping and tearing of tissue with excessive force.<sup>11</sup> Which in turn causes tissue damage and more complications. Henceforth, force reflecting operative instruments (FROIs) have been developed to increase haptic feedback during laparoscopic surgery.

Haptic feedback is the human physiological process of touch. Haptic perception incorporates tactile and kinaesthetic perception. Tactile perception is based on pressure receptors in our skin of our fingers and hand, which detect pressure, vibration, and texture. Our kinaesthetic perception is based on receptors in our muscles, tendons, and joints. They detect position, movement, and force applied.<sup>12</sup>

This very sensitive input and output of nerve signals are a continuously ongoing motion loop with hardly noticeable delay. Although this cycle is automatic and independently controlled, the surgeon can adjust this cycle by increasing or decreasing on purpose the position and power of the applied muscles. Overall, the effect on tissue manipulation and movement and the position of instruments is based on visual control of the kinaesthetic part of the process of feeling the haptic process.

This physiological process of feeling touch has been copied in haptic feedback graspers for use in laparoscopic surgery. The nerve endings have been replaced by the tip of the glass fibres at the end of the graspers. The small nerve sensors have been replaced by fibre Bragg gratings of 2 to 4 mm. The shift of light frequencies, on the level of nanometres, caused by stretching or compressing these gratings, is transported through the long glass fibre to the control unit. In the unit, the special gator measures the information and translates this into a signal to a control board. The output of an electrical signal is modulated to control the actuator in the hand piece of the surgeon and creates the resistance felt on the grasper tip to the gripper on the fingers of the surgeon.<sup>11,13,14</sup>

So, the two-way physiological system in the human is copied in this FROI instrument to create real haptic feedback without any delay, as the frequency is 6000 per second. The modulation of the feeling on purpose can be done by changing the gain or pre-settings.

The introduction of a new technology in surgery is related with complications and poor outcome.<sup>15</sup> The

introduction of new technology in surgery may also be influenced by industry's financial incentives, physician's drive to remain competitive, and the lure of new technology.<sup>16</sup> SAGES has created guidelines for the introduction process of new technology and techniques.<sup>17</sup> They have recommended that health technology assessment (HTA) is the currently best accepted method for the analysis of optimisation of health outcomes and related costs and should be the initial step for introduction of a new technology.<sup>18</sup> In this article, we have performed a HTA by the use of a decision analytic model comparing the current standard technology, conventional laparoscopic surgery, to the new technology haptic feedback in laparoscopic surgery for DE of the bowel and urinary tract. The aim of this study was to estimate the potential value of the use of haptic surgery during laparoscopic DE surgery involving the bowel and/or urinary tract by using a decision analytic model.

## Methods

### Model

To assess the potential value of haptic feedback during laparoscopy for DE, a decision analytical model was constructed. The model was used to synthesise various sources of evidence in order to estimate expected differences in costs and effects for the 2 strategies under comparison: the current standard of care, in which minimally invasive surgery without haptic feedback is performed, and the strategy with haptic feedback added to laparoscopic instruments. Therefore, complications were included in the model that could be related to the absence of haptic feedback.

### Population

No approval by the institutional medical ethics review board was required. The target population of the model consists of patients with DE in the Netherlands, who received a surgical procedure for removal of a primary bowel or urinary tract DE lesion. The starting population in the model is an annual number of 636 patients registered at the Dutch Endometriosis Foundation who have undergone surgery to remove DE lesions in the Netherlands.<sup>5</sup> Large variation exists regarding the exact distribution between primary lesions (primary bowel lesions or primary urinary tract lesions). Therefore, we made an assumption of this distribution based on the prevalence of urinary tract DE (19.5%) found in a study with 221 patients with DE.<sup>19</sup> We assumed that the remainder of patients (80.5%) had a primary bowel DE lesion.

### Selection of Complications Through an Expert Panel Using a Delphi Procedure

There was no clinical evidence on which intraoperative and post-operative complications might be prevented


by the use of haptic feedback in laparoscopic surgery. Therefore, an expert panel (consisting of gynaecologists and surgeons) made a selection of intraoperative and post-operative complications (due to undetected intraoperative injury) which can potentially be reduced by the advantages of having haptic feedback in laparoscopic surgery (Supplementary Material S1). Consensus was achieved according to the Delphi method.<sup>20</sup> Their selection was supported by several preclinical studies and veterinary studies.<sup>11-14,21</sup> The following intraoperative complications were included in the model: bowel injury, haemorrhage, and ureteral injury. Furthermore, the following post-operative

complications were included in the model: late bowel perforation, vesicovaginal fistula, ureteral fistula, rectovaginal fistula, and anastomotic leakage.

### Model Structure

A decision tree was developed to simulate the chain of events in the target population within both strategies. For both strategies, the structure of the model was similar.

A complete overview of the decision tree is shown in Figure 1. First of all, a division was made between laparoscopic procedures performed for primary bowel DE



**Figure 1.** Decision tree comparing laparoscopic instruments with and without haptic feedback graspers.

lesions and for primary urinary tract lesions. Secondly, a division was made between either having intraoperative or post-operative complications. Thirdly, the intraoperative and post-operative complications were divided in the specific complications and subsequent clinical consequences per complication.

### Model Input

We searched the peer-reviewed scientific medical literature to inform model parameters in PubMed. Complication rates for primary bowel DE were often shown separately per surgical technique: shaving, discoid excision, or segmental resection. The use of these techniques varies per centre and country. For the Netherlands, the distribution was assumed to be 10% shaving, 10% discoid excision, and 80% segmental resection. Mean weighed complication rates for primary bowel DE were subsequently calculated based on this distribution.<sup>6,22</sup> Bowel DE complication rates in current standard of care were based on a recent review by Donnez and Roman<sup>22</sup> as well as on the results of a consecutive cohort of 23 laparoscopically treated cases with bowel DE in Amsterdam University Medical Centre location VUmc. Urinary tract DE complication rates were based on a systematic review by Cavaco-Gomes et al.<sup>8</sup> An overview of all transition probabilities is shown in [Table 1](#).

### Costs

A Dutch healthcare perspective was used in the model, and healthcare-related costs were included. Treatments to solve intraoperative or post-operative complications in the model were associated with a combination of additional length of hospital stay, additional operating time, additional outpatient visits as well as a number of reinterventions and/or conversions to laparotomy. These healthcare resources were subsequently valued by cost prices per unit to derive additional costs due to complications. All costs were inflated to 2017 prices using the Dutch consumer price index. An overview of all costs per type of complication is shown in [Table 2](#). Colostomy and subsequent colostomy reversal procedures were performed for multiple complications.

### Assumptions

Some assumptions had to be made due to the limited amount of available data or for the simplicity of the model. It was assumed in the model that no complications occur during or following a reintervention. Also, patients were assumed to have either an intraoperative or a post-operative complication and no elevated probability of a post-operative complication after an intraoperative complication.

Based on expert opinion and the overall strategy in the Netherlands, it was assumed that in case of conversion for

intraoperative bowel injury, most patients receive a temporary colostomy. The reasoning behind this assumption is that conversion to laparotomy is usually performed when serious intraoperative injury occurs, for which colostomy would also be indicated. For late bowel perforations, it was assumed that 70% of patients undergo a colostomy procedure and the remaining 30% undergo a laparoscopic resuturing procedure.<sup>40,41</sup>

### Analyses

Since haptic feedback in laparoscopic graspers is not yet widely implemented and used, there is no clinical evidence on the effect it may have on intraoperative and post-operative complications. Therefore, a headroom analysis was conducted which shows the value of haptic feedback in case it would be able to prevent all selected intraoperative and post-operative complications occurring in primary bowel and urinary tract DE, that is the maximum potential value. The costs of the laparoscopic grasper with haptic feedback are currently unknown and therefore not included in the model.

Since the real performance of haptic feedback is likely to be lower than 100% (as assumed in the headroom analysis), we performed scenario analyses to get insight into the potential value of haptic feedback with different levels of effectiveness on reducing complications. In sensitivity analyses, we varied the value of several input variables to assess their influence on the outcomes.

### Results

A total of 9.7 intraoperative and 47.0 post-operative complications are expected in the cohort of 636 patients annually treated for DE in the Netherlands under the current standard of care. The distribution of various types of complications is listed in [Table 3](#). Of these complications, 83.0% occurred post-operatively and 85.7% of complications were related to primary bowel DE.

Using the model, the additional length of stay, additional number of outpatient visits, number of reinterventions, and number of conversions due to the selected intraoperative and post-operative complications were calculated. These results are presented in [Table 4](#). Next, we calculated the costs associated with the clinical consequences related to the surgical complications. Intraoperative and post-operative complications caused an additional length of stay of 432.1 days for the yearly cohort of patients. Furthermore, these complications initiated 10.2 additional outpatient visits, 73.9 additional reinterventions, and 4.2 additional conversions. The total additional annual costs due to complications were €436.623, amounting to €687 additional costs per patient. It was seen that most additional days of hospital stay, outpatient visits, surgical reinterventions, and ultimately

**Table 1.** Transition Probabilities Used in the Model.

Parameter	Value (%)	Source Used to Estimate Parameter
Surgical procedures for primary urinary tract DE	19.5	Gabriel et al <sup>19</sup>
Surgical procedure for primary bowel DE	80.5	Based on 1% of urinary tract DE
Intraoperative complications		
Bowel injury		
Incidence in primary bowel DE	.23	Donnez and Roman <sup>22</sup> and VUmc database
Incidence in primary urinary tract DE	n/a	
Conversion necessary	40.0	Kambakamba et al <sup>9</sup>
Colostomy in case of conversion	100	Expert opinion
No conversion necessary	60.0	Kambakamba et al <sup>9</sup>
Haemorrhage		
Incidence in primary bowel DE	.61	Donnez and Roman <sup>22</sup> and VUmc database
Incidence in primary urinary tract DE	.44	Cavaco-Gomes et al <sup>8</sup>
Conversion necessary	47.6	Kambakamba et al <sup>9</sup>
No conversion necessary	52.4	Kambakamba et al <sup>9</sup>
Ureteral injury		
Incidence in primary bowel DE	.11	Donnez and Roman <sup>22</sup> and VUmc database
Incidence in primary urinary tract DE	3.46	Cavaco-Gomes et al <sup>8</sup>
Conversion necessary	40.0	Kambakamba et al <sup>9</sup>
No conversion necessary	60.0	Kambakamba et al <sup>9</sup>
Post-operative complications		
Late bowel perforation		
Incidence in primary bowel DE	1.28	Donnez and Roman <sup>22</sup> and VUmc database
Incidence in primary urinary tract DE	.15	Cavaco-Gomes et al <sup>8</sup>
Primary laparoscopic bowel repair	30	Expert opinion
Colostomy	70	Expert opinion
Vesicovaginal fistula		
Incidence in primary bowel DE	.23	Balla et al <sup>23</sup>
Incidence in primary urinary tract DE	.15	Cavaco-Gomes et al <sup>8</sup>
Surgical management (cystorrhaphy)	33.3	Minelli et al <sup>24</sup>
Conservative management (catheterisation)	66.7	Minelli et al <sup>24</sup>
Ureteral fistula		
Incidence in primary bowel DE	.28	Donnez and Roman <sup>23</sup> and VUmc database
Incidence in primary urinary tract DE	2.05	Cavaco-Gomes et al <sup>8</sup>
Surgical management (reimplantation/anastomosis)	33.3	Donnez et al <sup>25</sup>
Conservative management (JJ stent)	66.7	Donnez et al <sup>25</sup>
Rectovaginal fistula		
Incidence in primary bowel DE	3.75	Donnez and Roman <sup>23</sup> and VUmc database
Incidence in primary urinary tract DE	n/a	
Surgical management (transabdominal procedure)	89.0	Kondo et al <sup>26</sup> and UpToDate <sup>27</sup>
Colostoma in case of surgical management	71.4	Minelli et al <sup>24</sup>
Conservative management (antibiotics and nutrition)	11.0	Kondo et al <sup>26</sup> and Donnez et al <sup>25</sup>
Anastomotic leakage		
Incidence in primary bowel DE	3.0	Donnez and Roman <sup>23</sup> and VUmc database
Incidence in primary urinary tract DE	.3	Cavaco-Gomes et al <sup>8</sup>
Surgical management (colostomy)	100	Seracchioli et al <sup>2</sup> and Belghiti et al <sup>28</sup>

Abbreviation: DE = deep endometriosis.

costs were associated with post-operative complications. Post-operative late bowel perforations, rectovaginal fistulas, and anastomotic leakages were responsible for 93% of the total costs.

The results listed in Table 4 represent the room for improvement, that is the consequences and subsequent costs

prevented in case haptic feedback was to reduce all complications. Additional scenario analysis was performed to show the effectiveness rate of haptic feedback per percentage point reduction of possible complications. If 1% reduction of complications is anticipated, €4366 cost reduction is achieved. Since it is a linear model, with a 2% reduction of

**Table 2.** Overview of all Costs per Type of Complication.

Parameter	Value	Source Used to Estimate Parameter
Unit costs		
OR time	€14.12 per minute	Local hospital data, Costing guideline (2015) <sup>29</sup> , and CAO UMC 2015-2017 (2015) <sup>30</sup>
Hospital stay	€486.95 per day	Costing guideline (2015) <sup>29</sup>
Outpatient visit	€93.09 per visit	Costing guideline (2015) <sup>29</sup>
Blood transfusion	€255.78	Local hospital data
Sterilisation of an open surgical instrument set in case of conversion	€148.35	Local hospital data
Colostoma disposables	€18.36 per day	Vennix et al <sup>31</sup>
Additional healthcare resource use due to intraoperative complications		
Additional length of stay in case of a conversion	3.1 days	Yang et al <sup>32</sup>
Additional OR time in case of a conversion	56.4 minute	Yang et al <sup>32</sup>
Additional instrument use in case of conversion	Open set	Expert opinion
Additional length of stay in case of intraoperative repair	2.0 days	Expert opinion
Additional OR time in case of intraoperative repair of haemorrhage or bowel injury	20.0 minute	Expert opinion
Additional OR time in case of intraoperative repair of ureteral injury	60.0 minute	Expert opinion
Additional healthcare resource use due to post-operative complications		
Primary laparoscopic repair of late bowel perforation		
OR duration	104.0 minute	Bleier et al <sup>33</sup>
Length of stay	5.1 days	Bleier et al <sup>33</sup>
Colostomy procedure		
OR duration	113.0 minute	Ivatury et al <sup>34</sup>
Length of stay	7.0 days	Ivatury et al <sup>34</sup>
Colostoma duration	91.0 days	Darai et al <sup>35</sup> and Donnez et al <sup>25</sup>
Colostomy reversal procedure		
OR duration	188.0 minute	Arkenbosch et al <sup>36</sup>
Length of stay	5.4 days	Arkenbosch et al <sup>36</sup>
Rectovaginal fistula repair and transabdominal procedure		
OR duration	145.0 minute	Hagen et al <sup>37</sup>
Length of stay	4.0 days	Hagen et al <sup>37</sup>
Ureteral fistula repair, reimplantation/anastomosis		
OR duration	141.0 minute	Wang et al <sup>38</sup>
Length of stay	5.6 days	Wang et al <sup>38</sup>
Outpatient visits to place and remove JJ stent	2.0 visits	Expert opinion
Ureteral fistula and conservative management		
Outpatient visits to place and remove JJ stent	2.0 visits	Expert opinion
Vesicovaginal fistula repair and laparoscopic cystorrhaphy		
OR duration	144.8 minute	Miklos et al <sup>39</sup>
Length of stay	1.2 days	Miklos et al <sup>39</sup>
Outpatient visits	1.0 visit	Miklos et al <sup>39</sup>
Vesicovaginal fistula and conservative management for catheter insertion and removal		
Outpatient visits	2.0 visits	Miklos et al <sup>39</sup> and Minelli et al <sup>24</sup>

Abbreviation: OR = operation room.

complications, €8732 is saved, with 3% reduction, €13 098 is saved, etc. These results are listed in [Table 5](#).

In sensitivity analyses, we varied the value of several input variables to assess their influence on the outcomes.

We varied the distribution of primary bowel DE and primary urinary tract DE. Instead of the distribution of 19.5/80.5% used in the base case analysis, we used a 5/95% distribution based on expert opinion. In addition, we

**Table 3.** Base Case Analysis Results, Number of Complications Occurring Annually in Primary Bowel and Urinary Tract DE in the Yearly Cohort of 636 Patients.

	Primary Bowel DE	Primary Urinary Tract DE	Total
Intraoperative complications	4.8	4.8	9.7
Bowel injury	1.2	—	1.2
Haemorrhage	3.1	0.5	3.7
Ureteral injury	0.5	4.3	4.8
Post-operative complications	43.7	3.3	47.0
Late bowel perforation	6.5	0.2	6.7
Vesicovaginal fistula	1.2	0.2	1.3
Ureteral fistula	1.4	2.5	4.0
Rectovaginal fistula	19.2	—	19.2
Anastomotic leakage	15.4	0.4	15.7
Total	48.5	8.1	56.6

Abbreviation: DE = deep endometriosis.

**Table 4.** Base Case Analysis Results, Annual Consequences of Intraoperative and Post-operative Complications for the Yearly Cohort of 636 Patients.

	Length of Stay	Outpatient Visit	Reintervention	Conversion	Cost
Intraoperative subtotal	26.3	0	0.5	4.2	€22 876
Bowel DE	14.5	0	0.5	2.2	€12 889
Urinary tract DE	11.7	0	0	2.0	€9988
Post-operative subtotal	405.8	10.2	73.9	0	€413 747
Bowel DE	394.6	4.8	72.0	0	€403 059
Urinary tract DE	11.2	5.4	1.9	0	€10 688
Total	432.1	10.2	73.9	4.2	€436 623

Abbreviation: DE = deep endometriosis.

**Table 5.** Results of Scenario Analyses with Different Effectiveness Rates of Haptic Feedback on the Possible Reduction of Complications in an Annual Cohort of 636 Patients.

Effectiveness (%)	Length of Stay	Outpatient Visit	Reintervention	Conversion	Cost
100	432.1	10.2	73.9	4.2	€436 623
50	216.1	5.1	37	2.1	€218 312
10	43.2	1	7.4	0.2	€43 662
5	21.6	0.5	3.7	0.1	€21 831
2	8.6	0.2	1.5	0	€8732
1	4.3	0.1	0.7	0	€4366

**Table 6.** Results of Sensitivity Analyses Showing the Total Annual Costs Related to Complications According to the Distributions in Primary Bowel and Urinary Tract Deep Endometriosis Lesion Sites.

Urinary Tract/Bowel Distribution (%)	Total Annual Cost
19.5/80.5	€436.623
5.0/95.0	€496.171
52.6/47/4	€300.689

used a 52.6/47.4% distribution, which was the upper bound with regard to primary urinary tract involvement of the prevalence range that was mentioned in the literature.<sup>42</sup> Results are displayed in [Table 6](#).

## Discussion

This study aimed to assess the potential value in reduction of surgical complications and costs due to the use of haptic surgery during DE surgery. This was done through the use

of decision analytic modelling by comparing the current standard of care, in which laparoscopic endometriosis surgery without haptic feedback is performed, and the strategy with haptic feedback added to laparoscopic instruments. Annually, a total of 57 complications could be prevented in 636 surgical procedures for DE in the Netherlands. The majority of these complications occur post-operatively. By preventing all complications included in the model, €436 623 could potentially be saved annually, which equals to €687 per patient. The largest part of these costs arises from post-operative complications, especially from late bowel perforations, rectovaginal fistulas, and anastomotic leakages. This maximum potential value (headroom) is expected to rise further when societal costs (eg costs caused by loss of work related productivity while having a colostomy) would be included in addition to healthcare costs, which were considered in this assessment. Also this study focussed on Clavien–Dindo grade III complications, while several Clavien–Dindo grade I/II complications, such as voiding dysfunction, also result in increased medical costs and social impairment.<sup>43</sup> In addition, complication rates in smaller centres with less experience in treating DE are likely to be higher than the rates reported by expert centres for DE in the literature, which were used as input for the model.

There is considerable variation in the literature on the reporting of complications after bowel or urinary tract surgery, which greatly influences the outcomes of the model, as is shown by the sensitivity analyses. In addition, the literature on surgical treatment of DE is of mediocre to poor quality overall, consisting mostly of case series and retrospective cohort studies, which is believed to result in overestimating the beneficial effect of surgery and under-reporting the total number of complications. Therefore, the room for improvement might be larger in practice.

Since the effectiveness of haptic feedback on reducing intraoperative and post-operative complications in complex endometriosis surgery is currently unknown, we first of all conducted a headroom analysis which did not include costs of research and development. Results of the headroom analysis show the room for improvement in this disease area, if haptic feedback would be able to prevent all selected complications. However, a 100% reduction of complications is unrealistic. We therefore also showed the potential value in case haptic feedback would be able to reduce less complications, although no literature describes the effectiveness of haptic feedback on reducing complications yet.

Some study limitations merit consideration. Since there was no clinical literature available on which types of intraoperative and post-operative complications haptic feedback might have an effect, an expert panel (consisting of leading gynaecologists and surgeons) made a selection of both intraoperative and post-

operative complications (due to undetected intraoperative injury) on which they expect a reduction by using haptic feedback. Their selection was based on the available literature which has proven the assumed benefits of haptic feedback in preclinical and veterinary studies.<sup>11,12,21</sup> The assumed effects which were taken in account are reduced applied gripping forces and better recognition of tissue properties including arterial pulsations resulting in a reduction of a selection of surgical complications.

According to SAGES guidelines, the first step for introducing a new technology or technique should be a HTA to assess potential improvement in health care and potential reduction of associated costs.<sup>17</sup> The next step is preclinical research, for which excellent results are reported in *in vitro*<sup>11</sup> and *in vivo* models<sup>21</sup> regarding the added value of haptic feedback in laparoscopic instruments. Finally, this technology should be introduced in clinical practice; however, physicians should be trained first and patients should be informed pre-operatively.<sup>17</sup>

The weakness related to the use of HTA is in general the dependence on the assumptions that have to be made to build the models. These assumptions are made on literature, interviews for clinical experience, and in this study, interview according to a Delphi procedure. However, in this study, we had to use the distribution of incidence of complications reported in the literature which showed a wide variation in outcomes. Moreover, as has been mentioned earlier, the exact impact of haptic feedback in the clinical practice is also unknown and has to be estimated based on scarce preclinical studies and personal experience. However, this study demonstrates that the potential value for improvement in DE by using haptic feedback instruments is considerable, mostly caused by the potential prevention of major post-operative complications. Sensitivity and scenario analyses showed that the number of complications which could be prevented and the potential cost savings are still considerable when key model parameters are varied.

This study demonstrated that the potential value for improvement in DE laparoscopic surgery by using haptic feedback instruments is considerable, mostly caused by the potential prevention of major post-operative complications.

### Author Contributions

Study concept and design: Stijn L. Vlek, Rens Burm, Tim M. Govers, Michel P. H. Vleugels, Jurriaan B. Tuynman, and Velja Mijatovic

Acquisition of data: Rens Burm, Tim M. Govers, and Michel P. H. Vleugels

Analysis and interpretation: Stijn L. Vlek, Rens Burm, Tim M. Govers, and Michel P. H. Vleugels

Study supervision: Michel P. H. Vleugels, Jurriaan B. Tuynman, and Velja Mijatovic



## Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

## ORCID iD

Stijn L. Vlek  <https://orcid.org/0000-0001-8614-0626>

## Supplemental Material

Supplemental material for this article is available online.

## References

- Giudice LC. Clinical practice. Endometriosis. *The New England Journal of Medicine*. 2010;362:2389-2398. doi:10.1056/NEJMcp1000274
- Seracchioli R, Poggioli G, Pierangeli F, et al. Surgical outcome and long-term follow up after laparoscopic resectosigmoid resection in women with deep infiltrating endometriosis. *BJOG*. 2007;114:889-895. doi:10.1111/j.1471-0528.2007.01363.x
- Wills HJ, Reid GD, Cooper MJ, et al. Fertility and pain outcomes following laparoscopic segmental bowel resection for colorectal endometriosis: A review. *Aust N Z J Obstet Gynaecol*. 2008;48:292-295. doi:10.1111/j.1479-828X.2008.00871.x
- Scala C, Barra F, Ferrero S, et al. Ureteral endometriosis: A systematic review of epidemiology, pathogenesis, diagnosis, treatment, risk of malignant transformation and fertility. *Human Reproduction Update*. 2018;24:710-730. doi:10.1093/humupd/dmy027
- Nederland ES. *Endometriose: zorg in ziekenhuizen 2016-2018*. 2018.
- Meuleman C, Tomassetti C, D'Hoore A, et al. Surgical treatment of deeply infiltrating endometriosis with colorectal involvement. *Human Reproduction Update*. 2011;17:311-326. doi:10.1093/humupd/dmq057
- Chudzinski A, Collinet P, Flamand V, et al. Ureterovesical reimplantation for ureteral deep infiltrating endometriosis: A retrospective study. *Journal of Gynecology Obstetrics and Human Reproduction*. 2017;46:229-233. doi:10.1016/j.jogoh.2017.01.001
- Cavaco-Gomes J, Martinho M, Gilabert-Aguilar J, et al. Laparoscopic management of ureteral endometriosis: A systematic review. *European Journal of Obstetrics, Gynecology, and Reproductive Biology*. 2017;210:94-101. doi:10.1016/j.ejogrb.2016.12.011
- Kambakamba P, Dindo D, Nocito A, et al. Intraoperative adverse events during laparoscopic colorectal resection—better laparoscopic treatment but unchanged incidence. Lessons learnt from a Swiss multi-institutional analysis of 3,928 patients. *Langenbeck's Archives of Surgery/Deutsche Gesellschaft für Chirurgie*. 2014;399:297-305. doi:10.1007/s00423-013-1156-4
- Westerbring-van der Putten EP, Goossens RH, Jakimowicz JJ, Dankelman J. Haptics in minimally invasive surgery - A review. *Minim Invasive Ther Allied Tech*. 2008;17(1):3-16.
- Alleblas CCJ, Vleugels MPH, Coppus S, et al. The effects of laparoscopic graspers with enhanced haptic feedback on applied forces: A randomized comparison with conventional graspers. *Surgical Endoscopy*. 2017;31:5411-5417. doi:10.1007/s00464-017-5623-9
- Wottawa CR, Genovese B, Nowroozi BN, et al. Evaluating tactile feedback in robotic surgery for potential clinical application using an animal model. *Surgical Endoscopy*. 2016;30:3198-3209. doi:10.1007/s00464-015-4602-2
- Vleugels MPH. Force reflecting instruments (FROI): The future becomes reality? *Journal of Minimally Invasive Gynecology*. 2007;14:S105. doi:10.1016/j.jmig.2007.08.164
- Vleugels M, Nieboer B. Real time haptic feedback in endoscopy: The proof of concept. *J Minim Invasive Gynecol*. 2015;22:S210-S211. doi:10.1016/j.jmig.2015.08.752
- Peltola M, Malmivaara A, Paavola M. Introducing a knee endoprosthesis model increases risk of early revision surgery. *Clinical Orthopaedics and Related Research*. 2012;470:1711-1717. doi:10.1007/s11999-011-2171-9
- Sachdeva AK, Russell TR. Safe introduction of new procedures and emerging technologies in surgery: Education, credentialing, and privileging. *The Surgical Clinics of North America*. 2007;87:853-866. doi:10.1016/j.suc.2007.06.006
- Stefanidis D, Fanelli RD, Price R, et al. SAGES guidelines for the introduction of new technology and techniques. *Surgical Endoscopy*. 2014;28:2257-2271. doi:10.1007/s00464-014-3587-6
- McGregor M, Brophy JM. End-user involvement in health technology assessment (HTA) development: A way to increase impact. *International Journal of Technology Assessment in Health Care*. 2005;21:263-267.
- Gabriel B, Nassif J, Trompoukis P, et al. Prevalence and management of urinary tract endometriosis: A clinical case series. *Urology*. 2011;78:1269-1274. doi:10.1016/j.urology.2011.07.1403
- Hsu C-C, Sandford BA. The Delphi technique: Making sense of consensus. *Practical Assessment, Research & Evaluation*. 2007;12:1-8.
- Alleblas CCJ, Vleugels MPH, Stommel MWJ, et al. Performance of a haptic feedback grasper in laparoscopic surgery: A randomized pilot comparison with conventional graspers in a porcine model. *Surgical Innovation*. 2019;26(5):573-580. doi:10.1177/1553350619848551
- Donnez O, Roman H. Choosing the right surgical technique for deep endometriosis: Shaving, disc excision, or bowel resection? *Fertil Steril*. 2017;108:931-942. doi:10.1016/j.fertnstert.2017.09.006
- Balla A, Quaresima S, Subiela JD, et al. Outcomes after resectosigmoid resection for endometriosis: A systematic literature review. *International Journal of Colorectal Disease*. 2018;33:835-847. doi:10.1007/s00384-018-3082-y
- Minelli L, Fanfani F, Fagotti A, et al. Laparoscopic colorectal resection for bowel endometriosis: Feasibility, complications, and clinical outcome. *Arch Surg*. 2009;144:234-239. doi:10.1001/archsurg.2008.555

25. Donnez J, Jadoul P, Colette S, et al. Deep rectovaginal endometriotic nodules: Perioperative complications from a series of 3,298 patients operated on by the shaving technique. *Gynecological Surgery*. 2013;10:31-40. doi:10.1007/s10397-012-0759-z
26. Kondo W, Bourdel N, Tamburro S, et al. Complications after surgery for deeply infiltrating pelvic endometriosis. *Bjog*. 2011;118:292-298. doi:10.1111/j.1471-0528.2010.02774.x
27. Marc R, Toglia M. *Rectovaginal and anovaginal fistulas*. [https://www.uptodate.com/contents/rectovaginal-and-anovaginal-fistulas?search=rectovaginal%20fistula&source=search\\_result&selectedTitle=1~37&usage\\_type=default&display\\_rank=1#H16](https://www.uptodate.com/contents/rectovaginal-and-anovaginal-fistulas?search=rectovaginal%20fistula&source=search_result&selectedTitle=1~37&usage_type=default&display_rank=1#H16)
28. Belghiti J, Ballester M, Zilberman S, et al. Role of protective defunctioning stoma in colorectal resection for endometriosis. *J Minim Invasive Gynecol*. 2014;21:472-479. doi:10.1016/j.jmig.2013.12.094
29. Roijen LH-v, Linden Nvd, Bouwmans C, et al. *Kostenhandleiding: Methodologie van kostenonderzoek en referentieprijzen voor economische evaluaties in de gezondheidszorg*Zorginstituut Nederland. 2015.
30. *Salarisschalen Cao umc 2015-2017, Bijlage A. 14-10-2015*. NFU; 2015.
31. Vennix S, van Dieren S, Opmeer BC, et al. Cost analysis of laparoscopic lavage compared with sigmoid resection for perforated diverticulitis in the Ladies trial. *The British Journal of Surgery*. 2017;104:62-68. doi:10.1002/bjs.10329
32. Yang C, Wexner SD, Safar B, et al. Conversion in laparoscopic surgery: Does intraoperative complication influence outcome? *Surgical Endoscopy*. 2009;23:2454-2458. doi:10.1007/s00464-009-0414-6
33. Bleier JI, Moon V, Feingold D, et al. Initial repair of iatrogenic colon perforation using laparoscopic methods. *Surgical Endoscopy*. 2008;22:646-649. doi:10.1007/s00464-007-9429-z
34. Ivatury SJ, Bostock Rosenzweig IC, Holubar SD. Short-term outcomes After open and laparoscopic colostomy creation. *Diseases of the Colon and Rectum*. 2016;59:543-550. doi:10.1097/dcr.0000000000000581
35. Darai E, Bazot M, Rouzier R, et al. Outcome of laparoscopic colorectal resection for endometriosis. *Current Opinion in Obstetrics & Gynecology*. 2007;19:308-313. doi:10.1097/GCO.0b013e328216f6bc
36. Arkenbosch J, Miyagaki H, Kumara HM, et al. Efficacy of laparoscopic-assisted approach for reversal of Hartmann's procedure: Results from the American college of surgeons national surgical quality improvement program (ACS-NSQIP) database. *Surgical Endoscopy*. 2015;29:2109-2114. doi:10.1007/s00464-014-3926-7
37. Van der Hagen SJ, Soeters PB, Baeten CG, et al. Laparoscopic fistula excision and omentoplasty for high rectovaginal fistulas: A prospective study of 40 patients. *International Journal of Colorectal Disease*. 2011;26:1463-1467. doi:10.1007/s00384-011-1259-8
38. Wang Z, Chen Z, He Y, et al. Laparoscopic ureteroureterostomy with an intraoperative retrograde ureteroscopy-assisted technique for distal ureteral injury secondary to gynecological surgery: A retrospective comparison with laparoscopic ureteroneocystostomy. *Scandinavian Journal of Urology*. 2017;51:329-334. doi:10.1080/21681805.2017.1304989
39. Miklos JR, Moore RD. Laparoscopic extravesical vesicovaginal fistula repair: Our technique and 15-year experience. *International Urogynecology Journal*. 2015;26:441-446. doi:10.1007/s00192-014-2458-y
40. Daams F, Sliker JC, Tedja A, et al. Treatment of colorectal anastomotic leakage: Results of a questionnaire amongst members of the Dutch society of gastrointestinal surgery. *Dig Surg*. 2012;29:516-521. doi:10.1159/000346348
41. Watson AJ, Krukowski ZH, Munro A. Salvage of large bowel anastomotic leaks. *The British Journal of Surgery*. 1999;86:499-500. doi:10.1046/j.1365-2168.1999.01096.x
42. Knabben L, Imboden S, Fellmann B, et al. Urinary tract endometriosis in patients with deep infiltrating endometriosis: Prevalence, symptoms, management, and proposal for a new clinical classification. *Fertil Steril*. 2015;103:147-152. doi:10.1016/j.fertnstert.2014.09.028
43. Poupon C, Owen C, Arfi A, et al. Nomogram predicting the likelihood of complications after surgery for deep endometriosis without bowel involvement. *European Journal of Obstetrics & Gynecology and Reproductive Biology: X*. 2019;3. doi:10.1016/j.eurox.2019.100028