

## Review

# Evidence-based approaches for establishing the 2015 Dietary Reference Intakes for Koreans

Sangah Shin<sup>1\*</sup>, Subeen Kim<sup>2\*</sup> and Hyojee Joung<sup>1,3§</sup>

<sup>1</sup>Department of Food and Nutrition, Chung-Ang University, 4726, Seodong-daero, Daedeok-myeon, Anseong, Gyeonggi 17546, Korea

<sup>2</sup>Department of Public Health, Graduate School of Public Health, Seoul National University, Seoul 08826, Korea

<sup>3</sup>Institute of Health and Environment, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea

**BACKGROUND/OBJECTIVES:** The Dietary Reference Intakes for Koreans (KDRIs), a set of reference intake values, have served as a basis for guiding a balanced diet that promotes health and prevents disease in the general Korean population. In the process of developing DRIs, a systematic review has played an important role in helping the DRI committees make evidence-based and transparent decisions for updating the next DRIs. Thus, the 2015 KDRI steering committee applied the systematic review framework to the revision process of the KDRIs. The purpose of this article is to summarize the revision process for the 2015 KDRIs by focusing on the systematic review framework.

**MATERIALS/METHODS:** The methods used to develop the systematic review framework for 2015 KDRIs followed the Agency for Healthcare Research and Quality and the Tufts Evidence-based Practice Center. The framework for systematic review of the 2015 KDRIs comprised of the 3 following steps: (1) development of an analytic framework and refinement of key questions and search terms; (2) literature search and data extraction; and, (3) appraisal of the literature and summarizing the results.

**RESULTS:** A total of 203,237 studies were retrieved through the above procedure, with 2,324 of these studies included in the analysis. General information, main results, comments of reviewers, and results of quality assessment were extracted and organized by study design. The average points of quality appraisals were 3.0 (range, 0-5) points for intervention, 6.1 (0-9) points for cohort, 6.0 (3-9) points for nested case-control, 5.4 (1-8) points for case-control, 14.6 (0-22) points for cross-sectional studies, and 7.0 (0-11) points for reviews.

**CONCLUSIONS:** Systematic review helped to establish the 2015 KDRIs as a useful tool for evidence-based approach. Collaborative efforts to improve the framework for systematic review should be continued for future KDRIs.

Nutrition Research and Practice 2018;12(6):459-468; <https://doi.org/10.4162/nrp.2018.12.6.459>; pISSN 1976-1457 eISSN 2005-6168

**Keywords:** Dietary reference intakes for Koreans, systematic review, evidence-based

## INTRODUCTION

Dietary reference intakes (DRIs) are reference values that guide the planning and assessment of nutrient intake in healthy populations [1]. The progenitor of DRI was the Recommended Dietary Allowances (RDAs) concept for energy, protein, and 8 vitamins and minerals, which was established in 1941 by the US National Research Council at the National Defense Advisory Commission [2]. RDAs were reference values that focused on preventing nutritional deficiencies [3]. According to the Institute of Medicine, RDA is "the average daily dietary nutrient intake level sufficient to meet the nutrient requirement of nearly all (97-98%) healthy individuals in a particular life stage and gender group" [4]. However, as various chronic diseases that were

derived from excess or imbalance intake became major issues in public health in the mid-1990s, the need for a comprehensive approach for nutrient intake recommendations became apparent, which resulted in the new paradigm called DRIs [5].

Most countries or regions recommend nutrient intake values that are specific to their populations because there are different dietary cultures and environments, as well as health/disease problems, among different populations. Traditionally, these standards were set at a level that would cover the dietary requirements of practically all healthy persons in a given population. As new knowledge of human nutrient requirements has grown, these standards require reassessment and revision when appropriate. Therefore, most countries examine the status of scientific evidence and update their nutrient intake values

We would like to thank the chairs and members of the 2015 KDRIs review committee, and the 6 working subcommittees (age and physique, infants, energy & macronutrients, vitamins, minerals, and applications), for their tremendous efforts for establishing the 2015 KDRI. This project was funded by the Korean Ministry of Health and Welfare (Contract No. 2013: 12138153800; 2014: 25143119100; 2015: 25153092900).

<sup>§</sup> Corresponding Author: Hyojee Joung, Tel. 82-2-880-2716, Fax. 82-2-883-2832, Email. [hjjoung@snu.ac.kr](mailto:hjjoung@snu.ac.kr)

Received: July 25, 2018, Revised: August 21, 2018, Accepted: November 5, 2018

\* These authors contributed equally to this work.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

periodically (about every 5-10 years) [6].

The initial Korean RDA (KRDA) for 10 nutrients, including energy, was published in 1962 by the Food and Agriculture Organization (FAO) Korean Office and has been periodically revised (the 7th revision was published in 2000) [7]. However, the need for new nutrient reference values has arisen as dietary and lifestyle changes have resulted in an increased prevalence of chronic diseases in the mid-20th century in Korea. Thus, in 2002, the Korean Nutrition Society (KNS) organized a committee to revise the KRDA and subsequently decided to expand the recommendation to the Dietary Reference Intakes for Koreans (KDRI), in order to address the diverse issues related to nutrition. In 2005, the KNS officially established the KDRI, which consisted of Estimated Average Requirement (EAR), Recommended Nutrient Intake (RNI), Adequate Intake (AI), Tolerable Upper Intake Level (UL) for nutrients, and Estimated Energy Requirement (EER) for energy and Acceptable Macronutrient Distribution Range (AMDR) of macronutrients [8]. The KDRI were revised in 2010 by the KNS.

Since the National Nutrition Management Act was announced in 2010, the task for establishing and revising the KDRI was conducted by the Korean Ministry of Health and Welfare (MOHW). Therefore, the MOHW recommended the task of revising the KDRI to the KNS, and for the first time published the KDRI at the national level in 2015 [9]. The 2015 KDRI are publicly available on the MOHW website (<http://www.mohw.go.kr>). In particular, the committee for the establishment of the 2015 KDRI put emphasis on an evidence-based multidisciplinary approach and the promotion of KDRI utilization.

The purpose of this article is to summarize the revision process for the 2015 KDRI by focusing on the systematic review framework of scientific evidence.

## MATERIALS AND METHODS

### *The development process of the 2015 KDRI*

DRI should be developed from scientific evidence that includes their role in eliminating nutritional deficiencies and reducing the risk of chronic diseases [10]. Where adequate information is available, each nutrient has a set of DRI. A nutrient has either an EAR and an RNI, or an AI. EAR and RNI are set when there is adequate information regarding the mean and standard deviation (SD) for the requirements of a given nutrient [9]. RNI is set to 2 SDs above the mean requirement, or a coefficient of variation is used instead of SD when insufficient data is available, such that 97.5% of the population's required intake would be below the recommended intake level [9]. When sufficient scientific evidence to derive the EAR and RNI cannot be obtained, the AI is used, which represents the adequate nutrient level for health [9]. UL is defined as "the highest average daily nutrient intake level that is likely to pose no risk of adverse health effects to almost all individuals in the general population" by the IOM [4]. UL is calculated based on the evidence of No Observed Adverse Effect Level (NOAEL), Lowest Observed Adverse Effect Level (LOAEL), and Uncertainty Factor (UF) [9]. The UL is determined only when there is evidence from the literature regarding adverse or toxic effects, such as NOAEL and LOAEL, from excessive intake of a given

nutrient [9]. AMDR indicates appropriate ranges of intakes for energy sources among macronutrients, such as carbohydrates, proteins, and lipids, for the prevention of chronic diseases and nutritional imbalance [7,9]. EER means the estimated value of total energy intake for individuals, which is calculated by the predictive equation taking into account age, height, weight, physical activity level, and life cycle [7,9]. Table 1 shows the established components of the 2015 KDRI by nutrients for individuals aged 1 year and older.

In Korea, an evidence-based research methodology has been applied to KDRI for developing the appropriate reference values for the Korean population from the initial 2005 version of the KDRI [8], with the first revision published in 2010 [11]. Evidence from new and relevant scientific reports has accumulated in recent decades, in addition to the systematic review framework for the United States and Canadian DRI (e.g., Ca and Vitamin D) published in 2011 [12]. Therefore, the committee for developing the 2015 KDRI applied the systematic review framework for establishing the 2015 KDRI.

### *The committee for developing the 2015 KDRI*

The KNS, at the request of the Korean MOHW, assembled a steering committee to establish the KDRI, based upon a rigorous and comprehensive review of the available scientific data. The KNS established a steering committee and a review board. The steering committee organized 6 subcommittees with a broad range of expertise (Fig. 1). The 6 subcommittees included: (1) age and physical standards; (2) standards for infants; (3) energy and macronutrients; (4) vitamins; (5) minerals; and, (6) applications of the KDRI. The review board was organized by experts, who participated in establishing or revising previous KDRI and reviewed both the procedures and results of the 2015 KDRI development.

### *Age subgroups and physical standards for the 2015 KDRI*

Age and gender subgroup categories, as well as the physical standards for each subgroup, were revised for the 2015 KDRI. To categorize age for the 2015 KDRI, the committee reviewed the age group categories of the 2010 KDRI and the available national statistics, including Korean statistical information and educational statistics. In addition, the committee considered the age group categories used for the DRI of other countries for comparison. As a result, age was categorized into 13 subgroups; 0-5 months, 6-11 months, 1-2 years, 3-5 years, 6-8 years, 9-11 years, 12-14 years, 15-18 years, 19-29 years, 30-49 years, 50-64 years, 65-74 years, and 75 years and older.

Physical standards were derived from the data of Korea National Health and Nutrition Examination Survey 2008-2012, the Agency for Technology and Standards survey data (5th, 2003-2004; 6th, 2010), and the 2007 Korean growth chart, according to the age and gender subgroups of the KDRI.

### *Nutrients included in the 2015 KDRI*

One of the important roles of the committee is to screen the target nutrients to be included in the KDRI. Target nutrients are selected by the KDRI committee through an examination of the available scientific evidence and have extended its scope from 34 types of nutrients and energy in the 2005 KDRI, to




Fig. 1. Organization chart of the 2015 KDRI. <sup>1)</sup>Numbers in parentheses indicate the number of nutrients. Source: The Korean Nutrition Society (KNS), 2014-2015 [27,28].

Table 1. 2015 KDRI components for 1 year and older

Nutrients	Components of Dietary Reference Intakes			
	EAR	RNI	AI	UL
Energy and macronutrients				
Energy	O <sup>1)</sup>			
Carbohydrate <sup>2)</sup>				
Sugars <sup>3)</sup>				
Lipids <sup>2)</sup>				
Protein <sup>2)</sup>	O	O		
Amino acids	O	O		
Total fiber			O	
Water			O	
Vitamins				
Fat-soluble vitamins				
Vitamin A	O	O		O
Vitamin D			O	O
Vitamin E			O	O
Vitamin K			O	
Water-soluble vitamins				
Vitamin C	O	O		O
Thiamin	O	O		
Riboflavin	O	O		
Niacin	O	O		O
Vitamin B <sub>6</sub>	O	O		O
Folate	O	O		O
Vitamin B <sub>12</sub>	O	O		
Pantothenic acid			O	
Biotin			O	

Table 1. continued

Nutrients	Components of Dietary Reference Intakes			
	EAR	RNI	AI	UL
<b>Minerals</b>				
<b>Macrominerals</b>				
Calcium	○	○		○
Phosphorus	○	○		○
Sodium			○	○ <sup>4)</sup>
Chloride			○	
Potassium			○	
Magnesium	○	○		○
<b>Microminerals</b>				
Iron	○	○		○
Zinc	○	○		○
Copper	○	○		○
Fluoride			○	○
Manganese			○	○
Iodine	○	○		○
Selenium	○	○		○
Molybdenum	○	○		○
Chromium			○	

EAR, estimated average requirement; RNI, recommended nutrient intake; UL, tolerable upper intake level; AI, adequate intake

<sup>1)</sup> Estimated energy requirement for energy

<sup>2)</sup> Acceptable macronutrient distribution range for energy sources among macronutrients

<sup>3)</sup> Recommendations for sugars

<sup>4)</sup> Intake goal

Source: Ministry of Health and Welfare (MOHW) & KNS, 2015 [9].

36 types in the 2015 KDRI [9]. Total sugars in 2010 and chromium in 2015 were newly added to their respective KDRI. As a result, 2015 KDRI included energy and 7 macronutrients (carbohydrate, sugars, lipids, protein, amino acids, total fiber, and water), 13 vitamins (vitamin A, D, E, K, C, thiamin, riboflavin, niacin, vitamin B<sub>6</sub>, folate, vitamin B<sub>12</sub>, pantothenic acid, and biotin), and 15 minerals (calcium, phosphorus, sodium, chloride, potassium, magnesium, iron, zinc, copper, fluoride, manganese, iodine, selenium, molybdenum, and chromium) in the KDRI (Table 1).

*The systematic review process for the development of the 2015 KDRI*

The KDRI are quantitative nutrient reference values that reflect both adequate and safe upper levels of intake, and comprise the evidentiary base and reference standards that underpin government food and nutrition programs, policies, and regulation. This includes food fortification, nutrition labeling, food expenditure, and guidelines for food and nutrition assistance programs. KDRI also serve as a basis for developing and revising national dietary and food guidelines. In addition, many other stakeholders use the KDRI for health delivery, as well as education, research, and other program initiatives. Therefore, KDRI should be based on scientific evidence of the association between nutrients and health outcomes, including basic requirements and the prevention of adverse effects. Particularly, the application of scientific evidence provides a foundation for establishing KDRI (Fig. 2) and may improve the validity and reliability of KDRI by making the best use of the existing scientific literature [9].

The 2015 KDRI were developed by using systematic evidence

review and analysis, based on the systematic review framework used by the Agency for Healthcare Research and Quality (AHRQ) and the Tufts Evidence-based Practice Center (EPC) for updating the 2011 DRI for calcium and vitamin D [12,13]. The systematic review framework for the 2015 KDRI was adapted by the steering committee using the 3 following steps: (1) development of an analytic framework and refinement of key questions and search terms; (2) literature search and data extraction; and, (3)




Fig. 2. Revision process of the 2015 KDRI. KNHANES, Korean National Health & Nutrition Examination Survey; EAR, estimated average requirement; RNI, recommended nutrient intake; UL, tolerable upper intake level; AI, adequate intake. Source: MOHW & KNS, 2015 [9].

appraisal of the literature and summarizing the results. After development, the steering committee provided training on the systematic review framework to each subcommittee member at least once to assist them in fully understanding the framework and to be able to conduct the systematic review. Each subcommittee performed a literature search, evaluated the literature selected, and summarized the main findings following the developed framework. Subsequently, the main findings of the systematic review were used for the 2015 revision of the KDRI. The contents of the systematic review framework for the 2015 KDRI are described in detail in the following paragraphs.

#### *Development of the analytic framework and refinement of key questions and search terms*

An analytic framework is a visual map that shows a complex, connected structure of exposures, health outcomes (or clinical outcomes), biological functions of a given nutrient, modifying factors, and the population of interest, based on the existing scientific knowledge [14]. The analytic framework helps researchers consolidate and translate study results and plays a guiding role in the integration of information from a variety of sources [14]. In addition, it is useful in determining key questions in the next step of the process by summing-up the relationship between exposure and health outcomes in the analytic framework [14]. The analytic framework for DRIs includes 2 aspects of nutrient intake on health outcomes: beneficial effects and adverse effects [12,13]. The analytic framework for beneficial effects is used for establishing EAR values, and the analytic framework for adverse effects is used to determine the UL values. For the development of the analytic framework, it is necessary to identify which biomarkers are appropriate to explain the underlying association between exposure and health outcomes through the existing scientific evidence. The analytic framework consists of 4 components: (1) exposures and/or sources (e.g., nutrient intake levels); (2) indicators of exposure reflecting nutritional status (e.g., blood concentration of the nutrient); (3) surrogate outcomes or intermediate outcomes that may predict or substitute for health outcomes (e.g., blood pressure, or blood cholesterol levels); and, (4) health outcomes or clinical outcomes (e.g., cardiovascular disease) [9,12]. Given that the relationship of these components varies for each nutrient, different analytic frameworks are needed for each nutrient of interest [15]. As an example for the 2015 KDRI revision, the analytic framework components relevant to establishing the EAR and UL for folate are identified as follows.

Regarding EAR, the exposures (and/or sources) of folate were food, fortified foods, and supplements. The indicators of exposures were serum folate, red blood cell (RBC) folate, and plasma homocysteine. The surrogate or intermediate outcomes were megaloblastic anemia, gastrointestinal dysfunction, and depression. The health or clinical outcomes were pregnancy and childbirth, neural tube defects in the fetus, cardiovascular disease, development of some cancers, and delay of growth. Regarding UL, the exposures (and/or sources) were fortified foods and supplements. The indicators of exposures were serum folate and RBC folate. The surrogate or intermediate outcomes were not identified. The health outcomes or clinical outcomes were the neurological disorder, masking vitamin B<sub>12</sub> deficiency,

and lowering zinc status.

Based on the analytic frameworks, the next step is to formulate key questions regarding the relationship of the analytic framework components and then determine the key search terms [15]. Key questions define the objective and range that need to be covered prior to the literature search to achieve the objective and also limit the scope of a study at an early stage of research [15]. While key questions should be clearly articulated and specified [14,15], these questions must be comprehensive in nature. Therefore, a collaborative process is required for formulating key questions, which embraces stakeholders, targeted users, and both authorities and committee members [14]. Also, the Population, Intervention or exposure, Control or Comparator, and Outcomes (PICO) strategy [16] is commonly used to enhance transparency and reliability in the process of developing key questions [15]. The committee for the 2015 KDRI carefully determined the key questions that should be answered for setting reference values by considering the relationship between these indicators. For example, several key questions to set the reference values for folate were developed such as: "What effects do dietary folate and folic acid (folate supplements) have on serum folate, RBC folate, and plasma homocysteine level"; and "What effects does folate intake have on surrogate (intermediate) outcomes like megaloblastic anemia and health (clinical) outcomes like neural tube defects in the fetus".

The committee for the 2015 KDRI categorized the analytic framework components into 2 parts: the exposures (or sources) of nutrients, and the outcomes of interest. Using folate as an example, the key search terms for exposures were folate and folic acid, and the key search terms for health outcomes were megaloblastic anemia, neural tube defects, hyperhomocysteinemia, cardiovascular disease, cancer or neoplasms, preterm or low birth weight, and breast milk or lactation. Then, terminology refinements, such as dealing with similar terms, were needed to determine both the final key search terms and the search strategies.

#### *Literature search and data extraction*

The literature search should be conducted using a well-structured search strategy with predefined criteria. The 2015 KDRI committee searched articles using refined key search terms and search strategies through domestic search engines [KISS (Koreanstudies Information Service System), RISS (Research Information Sharing Service), NDSL (National Digital Science Library), DBpia], as well as international search engines (PubMed, Cochrane Library, EMBASE, Science-Direct, Scopus). Eligibility criteria of the target population, language, journal, publication year, and study design are described in detail below. All articles reviewed for the 2015 KDRI only included articles that guaranteed quality using a peer-review system. The target population of study was healthy individuals, excluding infants under 6 months old. For infants, the subcommittee for infants independently conducted the literature review. There are English and Korean-language restrictions, but an exception was made in case of need for studies published in either Japanese or Chinese.

The availability of new and relevant scientific research was

**Table 2.** Extracted data according to each study design

Extracted data	Study design					
	Review <sup>1)</sup>	Intervention <sup>2)</sup>	Cohort <sup>3)</sup>	Nested case-control <sup>3)</sup>	Case-control <sup>3)</sup>	Cross-sectional <sup>4)</sup>
<b>General information</b>						
Authors	○	○	○	○	○	○
Article Title	○	○	○	○	○	○
Journal	○	○	○	○	○	○
Publication year	○	○	○	○	○	○
Volume, page	○	○	○	○	○	○
<b>Main results</b>						
Population	○	○	○	○	○	○
Background data		○				
Comparison	○	○	○	○	○	
Adjusted variables			○		○	○
Confounding variables				○		
Compliance		○				
Measure of exposure		○	○	○	○	○
Amount of intake (intake level)						○
Life stage		○	○	○	○	
Outcomes		○	○	○	○	○
Follow-up period		○	○	○	○	
Criteria of group classification			○	○	○	
Intervention (dose per one day)		○				
Category			○	○		○
Dose-rank			○	○	○	○
Number of Analyzed subjects		○				
Unit		○				
Number of case		○	○	○	○	○
Number of non-case			○	○	○	○
Number of total		○				
Mean, SD		○	○	○	○	○
Relative risk		○	○	○	○	○
Standard error		○				
95%CI (lower and upper limit)		○	○	○	○	○
Difference in net (95%CI)		○				
P-value		○	○	○	○	○
Summary of main findings	○					
Comments	○	○	○	○	○	○
Results of quality assessment <sup>5)</sup>	○	○	○	○	○	○

<sup>1,5)</sup> For review articles, AMSTAR [23] was used to assess the quality of studies.

<sup>2,5)</sup> For intervention studies, Jadad scale [20] was used to assess the quality of studies.

<sup>3,5)</sup> For cohort, nested case-control, and case-control studies, Newcastle-Ottawa [21] was used to assess the quality of studies.

<sup>4,5)</sup> For cross-sectional studies, STROBE [22] was used to assess the quality of studies.

Source: KNS, 2014 [27].

the primary consideration for establishing the 2015 KDRI. Thus, the availability of relevant research published after 2008 was first examined to cover the period after the 2010 KDRI. Then, the researches that were examined for the 2010 KDRI were re-reviewed for establishing the 2015 KDRI. Considering the strength and quality of scientific evidence according to each type of study [17,18], the committee gave a high priority to the results from systematic reviews and meta-analyses and also examined intervention studies, cohort studies, and nested case-control studies. In addition, case-control studies and cross-sectional studies were included in the systematic review if the research was conducted with a Korean or Asian population.

Retrieved articles without duplicates were first screened for eligibility through abstract review, and then the full-text of the articles were reviewed for inclusion in the systematic review. General information of the selected studies needed to be extracted and documented clearly, which includes information on the authors, article title, the journal of publication, and study design (Table 2), for assessing study quality and extracting study findings in the next step of the overall systematic review process.

#### *Appraisal of the literature and summarizing the results*

The final step in the systematic review framework for the 2015

KDRIs is to evaluate the quality of the selected studies and to summarize the main results of each study, as well as to provide reviewer comments on the study findings. The committee for the 2015 KDRIs considered a broad range of quality assessment tools that were verified internationally, according to each type of study design, for the appraisal of study quality [19]. The quality assessment tools used in the recent DRIs for calcium and vitamin D in the United States and Canada, the tools suggested by the Academy of Nutrition and Dietetics of the United States, and other tools widely used in the similar research field were examined thoroughly. Finally, 4 quality assessment tools were chosen for the systematic review for the 2015 KDRIs, as follows: (1) the Jadad scale [20] for intervention studies; (2) the Newcastle-Ottawa scale [21] for cohort, nested

case-control, and case-control studies; (3) the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) statement [22] for cross-sectional studies; and, (4) A Mea Surement Tool to Assess systematic Reviews (AMSTAR) [23] for review articles. Along with the quality appraisal, study findings were concisely summarized in a specific table to serve as key evidence for the KDRIs. Data regarding the study findings for the 2015 KDRIs included study population, measures of exposure, intervention, outcomes, relative risk, and so on (Table 2).

## RESULTS

### Results of systematic review for 2015 KDRIs

Out of the total 203,237 retrieved articles for the 2015 KDRIs

**Table 3.** Number of studies reviewed for the 2015 KDRIs

Nutrients		Retrieved studies	Primary selection	Secondary selection	
				Original articles	Review articles
Energy and macronutrients	Energy	6,135	68	57 (6/1/0/6/44) <sup>1)</sup>	9
	Carbohydrate	2,055	95	74 (64/3/1/0/6)	12
	Sugars	2,134	1,317	163 (27/34/0/8/94)	0
	Lipids	16,817	408	229 (84/60/14/17/54)	51
	Protein	32,619	337	92 (23/35/1/3/30)	33
	Amino acids	5,836	36	18 (13/1/1/0/3)	0
	Total fiber	54,324	415	358 (288/42/2/0/26)	13
	Water	2,076	85	62 (10/4/0/10/38)	12
Vitamins	Vitamin A	8,644	398	40 (33/4/0/0/3)	16
	Vitamin D	2,538	448	141 (29/66/26/4/16)	18
	Vitamin E	52	52	30 (21/1/4/1/3)	15
	Vitamin K	2,967	85	39 (24/6/2/1/6)	11
	Vitamin C	499	62	46 (13/16/2/5/10)	9
	Thiamin	2,990	25	22 (2/9/2/1/8)	1
	Riboflavin	3,367	188	82 (27/13/4/4/34)	8
	Niacin	295	15	9 (4/1/0/1/3)	6
	Vitamin B <sub>6</sub>	4,252	33	25 (1/6/2/4/12)	2
	Folate	1,539	76	23 (3/8/1/9/2)	4
	Vitamin B <sub>12</sub>	17,524	41	32 (4/7/1/5/15)	4
	Pantothenic acid	21	21	11 (3/1/0/0/7)	2
	Biotin	290	27	23 (12/1/0/3/7)	0
Minerals	Calcium	235	143	97 (20/16/1/5/55)	3
	Phosphorus	1,087	119	58 (10/0/0/5/43)	22
	Sodium/chloride	1,208	91	35 (4/5/0/0/26)	13
	Potassium	2,826	52	27 (10/1/0/2/14)	3
	Magnesium	11,600	92	59 (10/25/0/4/20)	17
	Iron	401	70	66 (21/6/1/5/33)	2
	Zinc	12,102	216	12 (8/0/0/1/3)	1
	Copper	3,700	56	25 (3/0/0/1/21)	8
	Fluoride	1,726	74	15 (8/1/0/0/6)	6
	Manganese	575	23	11 (0/5/0/1/5)	2
	Iodine	387	11	9 (1/0/0/1/7)	2
	Selenium	298	94	21 (10/3/2/0/6)	4
	Molybdenum <sup>2)</sup>	-	-	-	-
	Chromium	118	4	3 (3/0/0/0/0)	1
<b>Total number</b>		<b>203,237</b>	<b>5,277</b>	<b>2,014</b>	<b>310</b>

<sup>1)</sup> Numbers in parentheses indicate the number of intervention studies, cohort studies, nested case-control studies, case-control studies, and cross-sectional studies in order.

<sup>2)</sup> Review of molybdenum study was not available due to the lack of study published after 2008.

Source: KNS, 2015 [28].

**Table 4.** Results of the quality assessment of studies used for the 2015 KDRI

Nutrients		Study design					Review <sup>4)</sup>
		Intervention <sup>1)</sup>	Cohort <sup>2)</sup>	Nested case-control <sup>2)</sup>	Case-control <sup>2)</sup>	Cross-sectional <sup>3)</sup>	
Energy and macronutrients	Energy, carbohydrate, sugars, lipids, protein, amino acids, total fiber, and water	2.4 (0-5) <sup>5)</sup>	5.8 (0-9)	5.4 (4-8)	5.4 (1-7)	15.1 (0-22)	6.8 (1-11)
Vitamins	Vitamin A, D, E, K, C, thiamin, riboflavin, niacin, vitamin B <sub>6</sub> , folate, vitamin B <sub>12</sub> , pantothenic acid, and biotin	3.4 (0-5)	6.5 (0-9)	6.5 (3-9)	5.4 (2-7)	15.0 (0-22)	7.5 (0-11)
Minerals	Calcium, phosphorus, sodium/chloride, potassium, magnesium, iron, zinc, copper, fluoride, manganese, iodine, selenium, molybdenum <sup>6)</sup> , and chromium	2.9 (0-5)	5.7 (1-8)	7.2 (4-8)	5.4 (3-8)	13.4 (2-20)	6.6 (0-11)
Average (min-max) of total points		3.0 (0-5) of 5	6.1 (0-9) of 9	6.0 (3-9) of 9	5.4 (1-8) of 9	14.6 (0-22) of 22	7.0 (0-11) of 11

<sup>1)</sup> For intervention studies, Jadad scale [20] was used to assess the quality of studies.

<sup>2)</sup> For cohort, nested case-control, and case-control studies, Newcastle-Ottawa [21] was used to assess the quality of studies.

<sup>3)</sup> For cross-sectional studies, STROBE [22] was used to assess the quality of studies.

<sup>4)</sup> For review articles, AMSTAR [23] was used to assess the quality of studies.

<sup>5)</sup> Data presents average points and range (minimum to maximum points).

<sup>6)</sup> Molybdenum study was not included in the assessment due to the lack of study published after 2008.

Source: KNS, 2015 [28].

revision, 2,324 articles were included in the final systematic review (1,183 articles for energy and macronutrients, 619 articles for vitamins, and 522 articles for minerals) (Table 3). General information regarding the selected 2,324 articles such as authors, title, journal, and study design were extracted and documented in spreadsheets (Table 2). The information of excluded articles was also documented along with the reason for exclusion.

The committee made quality appraisals of the selected articles according to the manual of each quality assessment tool. The appraisal results were reported as scores with reviewer comments on the study evaluations, and the scores were converted to the score percentages out of the total scores as 100%. Table 4 shows the results of the quality appraisal according to each type of study design. On average, the score percentages of most studies were higher than a 60% total score percentage. For intervention studies, 799 studies were scored 3.0 (range, 0-5) points out of 5 total points on average. Cohort (n = 381), nested case-control (n = 67), and case-control studies (n = 107) were scored as 6.1 (range, 0-9), 6.0 (range, 3-9), and 5.4 (range, 1-8) points out of 9 total points, respectively. Cross-sectional studies (n = 660) received 14.6 (range, 0-22) points out of 22 total points and reviews (n = 310) were scored 7.0 (range, 0-11) points out of 11 total points (Table 4).

After the appraisal, data pertaining to the study findings were extracted and summarized in a table, along with the general information and appraisal results of the study. The results table concisely presents the study characteristics and findings, so that it assists the committee in translating the study outcomes to address the predetermined key questions [14]. Although the data components related to study findings are slightly different according to each type of study design, the results table for the 2015 KDRI contained study populations, comparisons, measures of exposure, life stage, interventions, outcomes, relative risk, 95% confidence intervals, reviewer's comments, and the results of the quality appraisal (Table 2). The results tables functioned as the key evidence for setting the reference values of the 2015 KDRI.

## DISCUSSION

This study has comprehensively detailed the systematic review process for the development of the 2015 KDRI. The 2015 KDRI provide not only a basic guide for a balanced diet to promote health, but it is also an important standard for the evaluation of nutrient intake levels for Koreans [9]. To establish scientifically sound and reliable KDRI, the decision making should proceed in a manner that is evidence-based and transparent. Unbiased systematic review plays an essential role in the decision-making process of the KDRI committee and facilitates the revision process of DRIs as additional findings are accumulated in the future [19]. For these reasons, the systematic review framework was developed and applied to the revision process for the 2015 KDRI.

The incorporation of the systematic review framework into the revision process for the 2015 KDRI was a complex task. However, considerable efforts on improving the framework should be continually developed to respond to global harmonization and the establishment of more elaborate KDRI. There are several suggestions that may be considered to improve the systematic review framework for future KDRI. First, an active discussion will be needed to include a wide range of chronic disease indicators into the analytic framework. The KDRI committee has made an effort to develop reference values for the prevention of chronic diseases and nutrient deficiency. To strengthen this effort, potential indicators related to chronic disease and nutrient exposure should be first identified. The potential indicators that can be incorporated into the analytic framework are the verified indicators that can be identified on the causal pathways between nutrient exposure and chronic disease [1]. However, it is difficult to identify these indicators for most chronic diseases [1]. Consequently, it requires an active discussion on how to select the appropriate chronic disease indicators, including proper surrogate indicators involved in chronic disease development (e.g., low-density lipoprotein-cholesterol for coronary heart disease) [24], and how to incorporate them into the analytic framework [12,25]. Second,



given the important role of selecting key questions that set the objectives and scope of the systematic review, it is desirable to encourage a broad participation of stakeholders, targeted users, authorities, and systematic review methodologists, as well as committee members, when developing the key questions [19]. Once the multi-sectoral team is organized, careful consideration is necessary to determine which topics should be addressed in setting KDRI as the key questions [18]. Lastly, nutrition-specific quality assessment tools that reflect the characteristics of nutrition-based research could be introduced during the study quality appraisal stage [19]. The causal relationship between nutrient exposure and health outcome is more complicated than the relationship between pharmacologic treatment and health outcome [19]. In addition, there are unique kinds of covariates, confounders, and sources of error to consider in nutrition research [26]. Therefore, for a critical appraisal of each study regarding the association between nutrient exposure and health outcomes, especially more complex chronic diseases, the nutrition-specific assessment tools could be utilized. For example, a new nutrition-specific assessment tool can be developed by adding evaluation items reflecting the characteristics of nutrition research to the existing evaluation tools [24,26].

In the process of revising DRIs, it is clear that the systematic review plays a key role, but the results of this process alone is not a substitute for the decision-making process of the review committee [12]. In other words, the systematic review is not the end of the committee's decision-making process, but is an integral part of the process for transparent and objective decision making [14]. Therefore, the systematic review framework for KDRI should be constantly improved and developed as a useful tool to help decision making of the committee.

At present, many experts and researchers are still making a commitment to future KDRI. The endeavors include not only improvements of the systematic review framework, but also the consideration of new approaches to establish reference values related to various chronic diseases, expansion of nutrients in KDRI, response to population aging and global harmonization, and so on [30]. The effort for promoting the use and application of KDRI should be made particularly through a close cooperation with professionals in government and fields. On the basis of the cooperation, it might be able to develop the dietary guidelines for Koreans, which is coordinated with KDRI. Although the 2015 KDRI were developed using evidence-based approaches, limited research evidence was available on the relationship between diet and health status in the Korean population. Given that the dietary recommendations and guidelines are applied to the dietary habit of the target population, further research about the Korean diet should be conducted for developing the next KDRI.

## CONFLICT OF INTEREST

The authors declare no potential conflicts of interests.

## ORCID

Sangah Shin: <https://orcid.org/0000-0003-0094-1014>

Subeen Kim: <https://orcid.org/0000-0003-4056-6925>

Hyojee Joung: <https://orcid.org/0000-0003-1182-7786>

## REFERENCES

- Murphy SP, Yates AA, Atkinson SA, Barr SI, Dwyer J. History of nutrition: the long road leading to the Dietary Reference Intakes for the United States and Canada. *Adv Nutr* 2016;7:157-68.
- National Research Council. Recommended Dietary Allowances. Washington, D.C.: The National Academies Press; 1941.
- National Research Council. Recommended Dietary Allowances: 10th Edition. Washington, D.C.: The National Academies Press; 1989. p.21.
- Institute of Medicine. Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate. Washington, D.C.: The National Academies Press; 2004. p.3.
- Institute of Medicine. The Development of DRIs 1994-2004: Lessons Learned and New Challenges: Workshop Summary. Washington, D.C.: The National Academies Press; 2008. p.2.
- King JC, Vorster HH, Tome DG. Nutrient intake values (NIVs): a recommended terminology and framework for the derivation of values. *Food Nutr Bull* 2007;28:S16-26.
- Paik HY. Dietary Reference Intakes for Koreans (KDRI). *Asia Pac J Clin Nutr* 2008;17:416-9.
- The Korean Nutrition Society. Dietary Reference Intakes for Koreans 2005. Seoul: The Korean Nutrition Society; 2005.
- Ministry of Health and Welfare; The Korean Nutrition Society. Dietary Reference Intakes for Koreans 2015. Seoul: The Korean Nutrition Society; 2015.
- Atkinson SA. Defining the process of Dietary Reference Intakes: framework for the United States and Canada. *Am J Clin Nutr* 2011;94:655S-657S.
- The Korean Nutrition Society. Dietary Reference Intakes for Koreans 2010. Seoul: The Korean Nutrition Society; 2010.
- Institute of Medicine. Dietary Reference Intakes for Calcium and Vitamin D. Washington, D.C.: The National Academies Press; 2011. p.523-1065.
- Chung M, Balk EM, Brendel M, Ip S, Lau J, Lee J, Lichtenstein A, Patel K, Raman G, Tatsioni A, Terasawa T, Trikalinos TA. Vitamin D and calcium: a systematic review of health outcomes. *Evid Rep Technol Assess (Full Rep)* 2009;(183):1-420.
- Lichtenstein AH, Yetley EA, Lau J. Application of systematic review methodology to the field of nutrition. *J Nutr* 2008;138:2297-306.
- Russell R, Chung M, Balk EM, Atkinson S, Giovannucci EL, Ip S, Lichtenstein AH, Mayne ST, Raman G, Ross AC, Trikalinos TA, West KP Jr, Lau J. Opportunities and challenges in conducting systematic reviews to support the development of nutrient reference values: vitamin A as an example. *Am J Clin Nutr* 2009;89:728-33.
- da Costa Santos CM, de Mattos Pimenta CA, Nobre MR. The PICO strategy for the research question construction and evidence search. *Rev Lat Am Enfermagem* 2007;15:508-11.
- Croswell JM, Kramer BS. Clinical trial design and evidence-based outcomes in the study of liver diseases. *J Hepatol* 2009;50:817-26.
- Joung HJ, Kim BH, Song YJ, Lee JE. Nutrition in Public Health. Paju: Yangseowon; 2013. p.184.
- Chung M, Balk EM, Ip S, Lee J, Terasawa T, Raman G, Trikalinos T, Lichtenstein AH, Lau J. Systematic review to support the development of nutrient reference intake values: challenges and solutions. *Am J Clin Nutr* 2010;92:273-6.

20. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, McQuay HJ. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17:1-12.
21. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses [Internet]. Ottawa: The Ottawa Hospital Research Institute; 2009 [cited 2018 June 16]. Available from: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp).
22. STROBE group. Strengthening the reporting of observational studies in epidemiology [Internet]. Bern: Institute of Social and Preventive Medicine at the University of Bern; 2009 [cited 2018 June 16]. Available from: <https://www.strobe-statement.org/index.php?id=available-checklists>.
23. Shea BJ, Grimshaw JM, Wells GA, Boers M, Andersson N, Hamel C, Porter AC, Tugwell P, Moher D, Bouter LM. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7:10.
24. Yetley EA, MacFarlane AJ, Greene-Finestone LS, Garza C, Ard JD, Atkinson SA, Bier DM, Carriquiry AL, Harlan WR, Hattis D, King JC, Krewski D, O'Connor DL, Prentice RL, Rodricks JV, Wells GA. Options for basing Dietary Reference Intakes (DRIs) on chronic disease endpoints: report from a joint US-/Canadian-sponsored working group. *Am J Clin Nutr* 2017;105:249S-85S.
25. Taylor CL. Framework for DRI development t: components "Known" and Components "To Be Explored" [Internet]. Washington, D.C.: The National Academies Press; 2008 [cited 2018 July 4]. Available from: [https://www.nal.usda.gov/sites/default/files/fnic\\_uploads/Framework\\_k\\_DRI\\_Development.pdf](https://www.nal.usda.gov/sites/default/files/fnic_uploads/Framework_k_DRI_Development.pdf).
26. National Academies of Sciences, Engineering, and Medicine. Global Harmonization of Methodological Approaches to Nutrient Intake Recommendations: Proceedings of a Workshop. Washington, D.C.: The National Academies Press; 2018. p.57-79.
27. The Korean Nutrition Society. 2014 Report on Support Project for Establishing 2015 Dietary Reference Intakes for Koreans. Seoul: The Korean Nutrition Society; 2014.
28. The Korean Nutrition Society. 2015 Report on Support Project for Establishing 2015 Dietary Reference Intakes for Koreans. Seoul: The Korean Nutrition Society; 2015.
29. The Korean Nutrition Society. 2013 Report on Support Project for Establishing 2015 Dietary Reference Intakes for Koreans. Seoul: The Korean Nutrition Society; 2013.
30. The Korean Nutrition Society. Report on the Plan for Ensuring Efficiency in Revision and Application of Dietary Reference Intakes for Koreans. Seoul: The Korean Nutrition Society; 2017.