



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Evaluation of the role of *Mycobacterium tuberculosis* pili (MTP) as an adhesin, invasin, and cytokine inducer of epithelial cells

Saiyur Ramsugit^a, Balakrishna Pillay^b, Manormoney Pillay^{a,*}

^a Medical Microbiology and Infection Control, University of KwaZulu-Natal, Durban, South Africa

^b School of Life Sciences, University of KwaZulu-Natal, Durban, South Africa

ARTICLE INFO

Article history:

Received 17 August 2015

Accepted 26 November 2015

Available online 31 December 2015

Keywords:

Mycobacterium tuberculosis

Curli pili

Adhesin

Invasin

ABSTRACT

This study was undertaken in order to assess the involvement of *Mycobacterium tuberculosis* pili (MTP) as an adhesin, invasin, and cytokine inducer in the *M. tuberculosis*-epithelial cell interaction. A MTP-deficient strain of *M. tuberculosis* demonstrated a significant reduction of 69.39% ($p = 0.047$) and 56.20% ($p = 0.033$) in its ability to adhere to and invade A549 pulmonary epithelial cells, respectively, in comparison with the wild-type strain. Complementation of the MTP-deficient mutant restored its adhesion and invasion capacity back to the wild-type levels. Overall, it was found that similar concentrations of IL-1 β , IL-4, IL-6, IL-8, G-CSF, IFN- γ , MCP-1, and TNF- α were induced in A549 cells infected with the MTP-proficient and MTP-deficient strains. However, at 48 h post-infection, the MTP-deficient mutant induced significantly lower levels of TNF- α than the wild-type strain ($p = 0.033$). Furthermore, at 72 h post-infection, the mutant induced significantly higher levels of IL-8 than the wild-type ($p = 0.005$). We conclude that MTP is an adhesin/invasin of epithelial cells and, while playing a role in *M. tuberculosis* entry, they do not appear to largely influence the epithelial cell cytokine response.

© 2016 Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Mycobacterium tuberculosis, the notorious causative agent of tuberculosis (TB), remains one of the main causes of human mortality by an infectious disease. The lack of effective treatment, diagnostic, and preventative strategies continue

to impede TB control, which is further compounded by the increasing prevalence of multi-, extensively, and totally drug-resistant strains of the pathogen and the HIV/AIDS co-epidemic. A better understanding of the virulence factors that are associated with the pathogenesis of this organism is crucial to improve strategies to control and reduce global TB burdens.

* Corresponding author at: Medical Microbiology and Infection Control, School of Laboratory Medicine and Medical Sciences, College of Health Sciences, University of KwaZulu-Natal, 1st Floor Doris Duke Medical Research Institute, Private Bag 7, Congella, 4013 Durban, South Africa.

E-mail address: [\(M. Pillay\).](mailto:Pillayc@ukzn.ac.za)

<http://dx.doi.org/10.1016/j.bjid.2015.11.002>

1413-8670/© 2016 Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Adhesins are molecules or structures that are present extracellularly on the bacterial cell surface. They play an important role in the initial host-pathogen interaction by mediating adherence, a precursor to host cell entry. A major *M. tuberculosis* adhesin of epithelial cells is the heparin-binding hemagglutinin adhesin (HBHA), which has also been implicated in extrapulmonary dissemination of the pathogen.¹ Bacterial species are known to produce multiple adhesins to effectively infect host cells. Therefore, other adhesins may contribute to *M. tuberculosis* adhesion to epithelial cells.²

Pili are a well-characterized family of bacterial adhesins. Two pili types have been described for *M. tuberculosis*, namely curli and type IV pili.³ The curli or coiled pilus morphotype, first identified in *Escherichia coli* and *Salmonella* spp., are known to be mediators in biofilm formation, adherence to and colonization of the host, and induction of host inflammation.⁴ The curli-like pili of *M. tuberculosis*, MTP, identified by Alteri et al.,⁵ resemble the pili fibers of *E. coli* and *Salmonella enterica* and are encoded by the *mtp* (*Rv3312A*) gene. These researchers also reported on the ability of MTP to bind to the extracellular matrix protein laminin *in vitro*, which suggested that MTP may function as an adherence factor.⁵

A central focus of our research group currently is elucidating the function of MTP. We have previously identified the role of MTP in *in vitro* biofilm formation⁶ and in the adherence to and invasion of macrophages.⁷ We have also shown that the *mtp* gene sequence is unique to *M. tuberculosis* complex strains and highly conserved amongst clinical isolates and, thus, MTP may be a suitable biomarker for a TB diagnostic test.⁸

In this study, we compared adhesion and invasion of a wild-type, Δmtp mutant, and *mtp*-complemented *M. tuberculosis* strains to the A549 pulmonary epithelial cell line. We also compared cytokine production by A549 cells infected with these strains.

Materials and methods

Bacterial strains and growth conditions

M. tuberculosis clinical isolate V9124⁹ and the previously constructed MTP-deficient Δmtp mutant and MTP-overexpressing *mtp*-complemented strains⁶ were used in this study. Cultures were grown at 37 °C in Middlebrook 7H9 broth (Difco), supplemented with 10% (v/v) oleic acid-albumin-dextrose-catalase (OADC; Becton Dickinson), 0.5% (v/v) glycerol (Sigma), and 0.05% (v/v) Tween 80 (Sigma). Hygromycin (75 µg mL⁻¹; Roche Applied Sciences) or kanamycin (30 µg mL⁻¹; Sigma) were included for the mutant and complemented strain cultures, respectively.

Epithelial cell preparation

The A549 human type II alveolar epithelial cell line (ATCC CCL-185) was maintained in Eagle's Minimum Essential Medium with Earle's Balanced Salt Solution and 2 mM L-glutamine (Lonza), supplemented with 10% (v/v) fetal bovine serum (Biowest). Following trypsinization, 2 × 10⁵ cells were seeded into 24-well plates (NEST Biotechnology) and incubated for 24 h at 37 °C in 5% CO₂.

Infection of epithelial cells

Logarithmic phase mycobacterial cultures were pelleted by centrifugation (2000 × g for 10 min; Heraeus Multifuge 3S-R Centrifuge, Thermo Scientific) and resuspended in fresh cell culture media. The epithelial cells were infected with the bacterial strains at a multiplicity of infection (MOI) of 1:1 for the adhesion and invasion assays. A MOI of 10:1 was used for the cytokine assay to ensure sufficient cytokine production.¹⁰ Plates were incubated at 37 °C and 5% CO₂ at the desired time intervals. The inocula were also serially diluted and plated in triplicate onto Middlebrook 7H11 (Difco) agar plates, containing 10% (v/v) OADC and 0.5% (v/v) glycerol, to enumerate viable bacilli used for infection. Trypan blue exclusion was used to determine the number of viable epithelial cells at the time of infection, to confirm the MOI.

Adhesion assay

The adhesion assay was performed as previously described.⁷ At the end of the 1 h incubation period, media was removed and the epithelial cells washed thrice with 1 mL phosphate-buffered saline (PBS; Oxoid). Adherent epithelial cells were lysed with 1 mL of 0.1% (v/v) Triton X-100 (Sigma) for 20 min at 37 °C and 5% CO₂. The number of adherent bacteria was determined by plating ten-fold serial dilutions of the lysate, in triplicate, onto 7H11 agar plates. Colonies were counted after 3 weeks of incubation at 37 °C.

Invasion assay

The invasion assay was performed as previously described.⁷ After 2 h incubation, monolayers were treated with amikacin (300 µg mL⁻¹; Sigma) for 1 h at 37 °C in a 5% CO₂ atmosphere. The media was removed and plated onto 7H11 plates to confirm the death of non-invaded microbes. The amikacin was removed by washing thrice with PBS. Adherent epithelial cells were lysed and the number of viable invaded organisms was determined as in the adhesion assay.

Cytokine analysis

Following 4 h of infection, spent media was removed, the wells washed with PBS, and fresh tissue culture media added. At 24, 48, and 72 h post-infection, cell culture supernatants were removed and filtered through a 0.2 µm filter (PALL Life Sciences). Bovine serum albumin (BSA; Sigma) was added at a final concentration of 0.5% (w/v) and the samples were stored at -80 °C. The cytokine levels in supernatants were measured using a Bio-Plex Pro Human Cytokine Multi-Plex Panel (Bio-Rad) in a Bio-Plex 200 System (Bio-Rad), according to the manufacturer's instructions.

Statistical analysis

Adhesion/invasion assays were performed at least three independent times in duplicate. Cytokine experiments were performed at two independent times and assayed in duplicate. The significance of differences between the strains was determined by one-way analysis of variance (ANOVA). All data




Fig. 1 – The role of *M. tuberculosis* pili (MTP) in the adhesion (A) to and invasion (B) of epithelial cells. A549 epithelial cells were infected with a wild-type, MTP-deficient mutant, and MTP-overexpressing complemented strains for 1 h (adhesion assay) or 2 h (invasion assay). Amikacin-treated and -untreated infected epithelial cells were washed, lysed, and plated onto agar plates to enumerate viable invaded and adherent bacilli, respectively. Results are expressed as the mean percentage adhesion to or invasion of the mutant and complemented strains, relative to that of the wild-type. Error bars are $\pm 1 \times$ SEM.

were analyzed using SPSS 22.0 statistical software. $p < 0.05$ was considered significant.

Results and discussion

The entry of tubercle bacilli into the alveolar space of the lungs is followed by the invasion or internalization of the organism by alveolar macrophages.¹¹ This interaction has important implications for the pathogen's survival and replication, evasion of the host immune response, and in its dissemination. Although less widely studied, *M. tuberculosis* also has the ability to invade and replicate in epithelial cells.¹²⁻¹⁴ Mehta et al.¹³ showed that whilst *M. tuberculosis* invades macrophages more efficiently than epithelial cells, the intracellular replication within the latter cell type was significantly higher. Epithelial cells may, therefore, provide a less harsh intracellular environment for the pathogen than macrophages. Furthermore, they are present in greater numbers than macrophages within alveoli¹⁵ and, thus, may represent the first host cell type that the bacilli encounter in the lung.

Using immunofluorescence microscopy with anti-MTP antibodies, Alteri¹⁶ detected expression of MTP in a previous study with A549 epithelial cells. This finding suggested a possible role of MTP in the *M. tuberculosis*-epithelial cell interaction. To ascertain whether MTP enable *M. tuberculosis* entry into epithelial cells, we compared adhesion and invasion capabilities of a MTP-deficient and a MTP-overexpressing *M. tuberculosis* strain in A549 cells. MTP-deficiency significantly diminished adhesion to and invasion of A549 cells compared with the parental strain, by 69.39% ($p = 0.047$) and 56.20% ($p = 0.033$), respectively (Fig. 1). Complementation restored adhesion ($p = 0.027$) and invasion ($p = 0.027$) to the wild-type levels (Fig. 1). Increased MTP expression by the complemented strain did not, however, significantly alter adhesion ($p > 0.999$) to and invasion ($p > 0.999$) of epithelial cells compared with the wild-type (Fig. 1).

Our findings are similar to those of other studies, which have shown that curliated bacteria are better able to adhere to and enter epithelial cells than non-curliated bacteria. Curliated *E. coli* strains were reported to be better at adhering to uroepithelial cells¹⁷ and human laryngeal epithelial (Hep-2) cells.¹⁸ They also displayed increased invasion of human cervical epithelial (HeLa) cells¹⁹ and Hep-2 cells.²⁰ Similarly, curliated *Salmonella typhimurium* promoted attachment to cultured mouse small intestinal epithelial cells.²¹

The current results differ from our previous study, which found that the adherence of *M. tuberculosis* to THP-1 macrophages was dependent on the density of pilation.⁷ The complemented strain expressed a 51-fold higher *mtp* expression than the wild-type, driven by the constitutively expressed *hsp60* promoter of the pMV261 vector used for complementation.⁶ This MTP-overexpression significantly increased adhesion to THP-1 macrophages, but not A549 epithelial cells, compared with the wild-type. This observation may possibly indicate different mechanisms or host receptor interaction of MTP with these two host cell types or, alternatively, varying expression of the same host cell receptor between macrophages and epithelial cells.

The only definitive adhesin/invasin previously described for *M. tuberculosis* is the HBHA. Pethe et al.¹ reported a 60% and 80% reduction in the adhesion to and invasion of A549 pneumocytes, respectively, by a HBHA-deficient strain compared with the wild-type. Loss of HBHA did not influence adhesion to and invasion of macrophages.¹ Due to the significant reduction in adherence to and invasion of epithelial cells, HBHA was regarded as the principal adhesin of epithelial cells. However, our study confirms that *M. tuberculosis* possesses more than a single adhesin, with a redundant function, which may account for the success of the pathogen. Therapeutic strategies targeting multiple adhesins may, therefore, prove most beneficial to incapacitating the pathogen.

In addition to serving as an adhesin and invasin of epithelial cells, we previously reported a 42% and 69% decrease in

adhesion to and invasion of THP-1 macrophages, respectively, by the MTP-deficient strain.⁷ The difference in adhesion and invasion rates suggest that MTP may play a more significant role as an adhesin in epithelial cells than in macrophages, whilst the opposite may be true for its function as an invasin.

Pethe et al.¹ suggested that following entry and replication within macrophages, *M. tuberculosis* enters non-phagocytic cells, which facilitate extrapulmonary dissemination. The function of MTP in both phagocytic and non-phagocytic host cell interactions implies that MTP is a major adhesin and participates in multiple stages of *M. tuberculosis* pathogenesis. During host colonization, bacteria are also known to develop into structured microbial communities or biofilms, which are associated with drug tolerance.²² In addition to serving as an adhesin of host cells, MTP mediate the formation of *in vitro* *M. tuberculosis* biofilms.⁶

Given the collective data on both adhesins, it seems that MTP may be a marginally better adhesin than HBHA and that the latter may, in turn, be more effective as an invasin compared with the former. This can only be confirmed by testing these deletion mutants under the same experimental conditions. However, both these adhesins/invasins appear to play major roles in the virulence of this pathogen and suggest that in combination, they may represent powerful targets for adjunctive immuno- or chemotherapy or potential vaccine development.

The outcome of *M. tuberculosis* infection is dependent on both the virulence of the pathogen and the host cellular defence mechanisms. *M. tuberculosis*-infected epithelial cells are known to secrete cytokines and chemokines, including interleukin-8 (IL-8) and monocyte chemotactic protein-1 (MCP-1), which contribute to the host inflammatory response.¹⁰ Pili do not only function in the host-pathogen interaction by enabling adhesion to and invasion of host cells, but the interaction of pili with host cell receptors is essential to the induction of inflammation via the production of inflammatory markers, such as cytokines.²³ Several studies have demonstrated that pathogen-induced inflammation on epithelial cells can be attributed to pathogen-associated molecular patterns, such as pili or flagella, which stimulate pro-inflammatory signals.²⁴⁻²⁷ In order to determine whether MTP are activators of the epithelial cell cytokine response, we compared the *in vitro* cytokine profiles of A549 cells infected with the wild-type, Δ mtp mutant, and complemented strains, using a multiplex assay.

The A549 cells were found to produce IL-1 β , IL-4, IL-6, IL-8, granulocyte colony-stimulating factor (G-CSF), interferon-gamma (IFN- γ), MCP-1, and TNF- α in this study, although at low concentrations for some analytes. Titers for IL-1 β , IL-4, IL-6, G-CSF, and IFN- γ were not significantly different for the epithelial cells infected with the wild-type, mutant, and complemented strains at all three time points tested (Fig. 2a, b, c, e and f). Similarly, there were no significant differences in IL-8, MCP-1, and TNF- α levels elicited by the MTP-deficient and MTP-proficient strains at most of the time intervals, with a few exceptions (Fig. 2d, g and h). The only indication that MTP induce the cytokine response was the observation that at 48 h post-infection, the mutant strain displayed a significant

reduction in TNF- α production compared with the wild-type ($p=0.033$) (Fig. 2h). However, at 24 and 72 h post-infection, the mutant strain showed, on average, higher levels of TNF- α induction than both the wild-type and complemented strains (Fig. 2h). The mutant strain also induced the lowest levels of IL-4, IL-8, and IFN- γ at 48 h post-infection compared with the MTP-proficient strains, but this did not reach statistical significance (Fig. 2b, d and f).

These findings suggest that, apart from TNF- α (and to a lesser extent IL-4, IL-8, and IFN- γ) at 48 h post-infection, MTP-mediated entry into epithelial cells does not trigger responses of the cytokines tested. These findings contrast other studies that have associated curli pili as inducers of the immune system. Tükel et al.²⁸ showed that *S. enterica* serotype Typhimurium curli are recognized by Toll-like receptor 2, resulting in the activation of IL-8. Bian et al.²⁹ demonstrated that curliated *E. coli* result in the activation of IL-6, IL-8, and TNF- α . Macrophages and human embryonic kidney cells, and not epithelial cells, were the host cell models in these studies. However, others have demonstrated an association between bacterial invasion and cytokine production by epithelial cells.³⁰⁻³² Interestingly, the mutant strain induced significantly higher levels of IL-8 than the wild-type at 72 h post-infection ($p=0.005$) (Fig. 2d). In addition, significantly more MCP-1 ($p=0.048$) and TNF- α ($p=0.004$) were secreted by mutant-infected epithelial cells at 72 and 24 h post-infection, respectively, compared with the complemented strain (Fig. 2g and h). Whilst not reaching statistical significance, the mutant strain showed, on average, elevated levels of all cytokines detected in this study, except IL-4 and IFN- γ , at 24 h post-infection than the wild-type and complemented strains (Fig. 2). Similarly, at 72 h post-infection, only the G-CSF concentrations were not the highest for the mutant strain samples (Fig. 2). In addition, at 48 h post-infection, the mutant strain induced the highest concentrations of IL-1 β , G-CSF, and MCP-1 (Fig. 2a, e and g). It is, therefore, tempting to speculate that MTP-mediated invasion of epithelial cells may confer an advantage to the pathogen by dampening the epithelial cell cytokine response, which is a key factor in infection control. This would represent an additional strategy of *M. tuberculosis* to ensure survival by escaping the host immune system. Alternately, the decreased cytokine levels may be attributed to increased cytotoxicity of wild-type and complemented strain-infected epithelial cells, due to the higher bacterial loads than for the mutant-infected epithelial cells. It has recently been reported that a positive correlation exists between *M. tuberculosis* adhesion to and invasion of epithelial cells and its cytotoxic effect.³³ Another possible reason for this phenomenon could be that MTP-deficiency triggers the expression of other adhesins, which induce cytokine production. Future cytotoxicity studies and cytokine assays on epithelial cells treated with purified MTP protein would help to verify these postulates.

To conclude, we have shown that MTP function in the adherence and entry of *M. tuberculosis* into epithelial cells. However, MTP did not largely influence the overall cytokine responses of *M. tuberculosis*-infected epithelial cells. Future *in vivo* studies will address the importance of these *in vitro* findings.




Fig. 2 – The role of *M. tuberculosis* pili (MTP) in epithelial cell cytokine production. A549 epithelial cells were exposed to the wild-type, MTP-deficient mutant, and MTP-overexpressing complemented strains for 4 h. Extracellular bacteria were removed by washing, fresh media added, and cell culture supernatants at 24, 48, and 72 h post-infection were assayed for levels of the cytokines IL-1 β (A), IL-4 (B), IL-6 (C), IL-8 (D), G-CSF (E), IFN- γ (F), MCP-1 (G), and TNF- α (H), using a Bio-Plex assay (Bio-Rad). Error bars are $\pm 1 \times$ SEM.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

The authors would like to thank the National Research Foundation of South Africa and the Canon Collins Educational and Legal Assistance Trust for financial support. We also thank Dr. Lenine Liebenberg (CAPRISA) for assistance with the cytokine data analysis.

REFERENCES

- Pethe K, Alonso S, Biet F, et al. The heparin-binding haemagglutinin of *M. tuberculosis* is required for extrapulmonary dissemination. *Nature*. 2001;412:190–4.
- Govender VS, Ramsugit S, Pillay M. Mycobacterium tuberculosis adhesins: potential biomarkers as anti-tuberculosis therapeutic and diagnostic targets. *Microbiology*. 2014;160:1821–31.
- Ramsugit S, Pillay M. Pili of *Mycobacterium tuberculosis*: current knowledge and future prospects. *Arch Microbiol*. 2015;197:737–44.
- Barnhart MM, Chapman MR. Curli biogenesis and function. *Annu Rev Microbiol*. 2006;60:131–47.
- Alerti CJ, Xicohténcatl-Cortes J, Hess S, Caballero-Olín G, Girón JA, Friedman RL. *Mycobacterium tuberculosis* produces pili during human infection. *Proc Natl Acad Sci U S A*. 2007;104:5145–50.
- Ramsugit S, Guma S, Pillay B, et al. Pili contribute to biofilm formation in vitro in *Mycobacterium tuberculosis*. *Antonie Van Leeuwenhoek*. 2013;104:725–35.
- Ramsugit S, Pillay M. *Mycobacterium tuberculosis* pili promote adhesion to and invasion of THP-1 macrophages. *Jpn J Infect Dis*. 2014;67:476–8.
- Naidoo N, Ramsugit S, Pillay M. *Mycobacterium tuberculosis* pili (MTP), a putative biomarker for a tuberculosis diagnostic test. *Tuberculosis (Edinb)*. 2014;94:338–45.

9. Pillay M, Sturm AW. Nosocomial transmission of the F15/LAM4/KZN genotype of *Mycobacterium tuberculosis* in patients on tuberculosis treatment. *Int J Tuberc Lung Dis.* 2010;14:223–30.
10. Lin Y, Zhang M, Barnes PF. Chemokine production by a human alveolar epithelial cell line in response to *Mycobacterium tuberculosis*. *Infect Immun.* 1998;66:1121–6.
11. Smith I. *Mycobacterium tuberculosis* pathogenesis and molecular determinants of virulence. *Clin Microbiol Rev.* 2003;16:463–96.
12. Bermudez LE, Goodman J. *Mycobacterium tuberculosis* invades and replicates within type II alveolar cells. *Infect Immun.* 1996;64:1400–6.
13. Mehta PK, King CH, White EH, Murtagh JJ Jr, Quinn FD. Comparison of in vitro models for the study of *Mycobacterium tuberculosis* invasion and intracellular replication. *Infect Immun.* 1996;64:2673–9.
14. Ashiru OT, Pillay M, Sturm AW. Adhesion to and invasion of pulmonary epithelial cells by the F15/LAM4/KZN and Beijing strains of *Mycobacterium tuberculosis*. *J Med Microbiol.* 2010;59:528–33.
15. Crandall ED, Kim KJ. Alveolar epithelial barrier properties. In: Crystal RG, West JB, editors. *The lung: scientific foundations*. New York: Raven Press; 1991.
16. Alteri CJ (Ph.D. thesis) Novel pili of *Mycobacterium tuberculosis*. University of Arizona; 2005.
17. Kikuchi T, Mizunoe Y, Takade A, Naito S, Yoshida S. Curli fibers are required for development of biofilm architecture in *Escherichia coli* K-12 and enhance bacterial adherence to human uroepithelial cells. *Microbiol Immunol.* 2005;49:875–84.
18. Kim SH, Kim YH. *Escherichia coli* 0157:H7 adherence to HEp-2 cells is implicated with curli expression and outer membrane integrity. *J Vet Sci.* 2004;5:119–24.
19. Gophna U, Barlev M, Seijffers R, Oelschlager TA, Hacker J, Ron EZ. Curli fibers mediate internalization of *Escherichia coli* by eukaryotic cells. *Infect Immun.* 2001;69:2659–65.
20. Uhlich GA, Keen JE, Elder RO. Variations in the *csgD* promoter of *Escherichia coli* 0157:H7 associated with increased virulence in mice and increased invasion of HEp-2 cells. *Infect Immun.* 2002;70:395–9.
21. Sukupolvi S, Lorenz RG, Gordon JI, et al. Expression of thin aggregative fimbriae promotes interaction of *Salmonella* *typhimurium* SR-11 with mouse small intestinal epithelial cells. *Infect Immun.* 1997;65:5320–5.
22. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. *Science.* 1999;284:1318–22.
23. Sauer FG, Mulvey MA, Schilling JD, Martinez JJ, Hultgren SJ. Bacterial pili: molecular mechanisms of pathogenesis. *Curr Opin Microbiol.* 2000;3:65–72.
24. Hedges S, Svensson M, Svartborg C. Interleukin-6 response of epithelial cell lines to bacterial stimulation in vitro. *Infect Immun.* 1992;60:1295–301.
25. Berin MC, Darfeuille-Michaud A, Egan LJ, Miyamoto Y, Kagnoff MF. Role of EHEC 0157:H7 virulence factors in the activation of intestinal epithelial cell NF-κappaB and MAP kinase pathways and the upregulated expression of interleukin 8. *Cell Microbiol.* 2002;4:635–48.
26. Xicohencatl-Cortés J, Lyons S, Chaparro AP, et al. Identification of proinflammatory flagellin proteins in supernatants of *Vibrio cholerae* O1 by proteomics analysis. *Mol Cell Proteomics.* 2006;5:2374–83.
27. Ledesma MA, Ochoa SA, Cruz A, et al. The hemorrhagic coli pilus (HCP) of *Escherichia coli* 0157:H7 is an inducer of proinflammatory cytokine secretion in intestinal epithelial cells. *PLoS ONE.* 2010;5:e12127.
28. Tükel C, Raffatellu M, Humphries AD, et al. CsgA is a pathogen-associated molecular pattern of *Salmonella enterica* serotype Typhimurium that is recognized by Toll-like receptor 2. *Mol Microbiol.* 2005;58:289–304.
29. Bian Z, Brauner A, Li Y, Normark S. Expression of and cytokine activation by *Escherichia coli* curli fibers in human sepsis. *J Infect Dis.* 2000;181:602–12.
30. Eckmann L, Kagnoff MF, Fierer J. Intestinal epithelial cells as watchdogs for the natural immune system. *Trends Microbiol.* 1995;3:118–20.
31. Hedges SR, Agace WW, Svartborg C. Epithelial cytokine responses and mucosal cytokine networks. *Trends Microbiol.* 1995;3:266–70.
32. Wilson M, Seymour R, Henderson B. Bacterial perturbation of cytokine networks. *Infect Immun.* 1998;66:2401–9.
33. Ashiru OT, Sturm AW. Cytotoxicity induction in A549 alveolar epithelial cells by *Mycobacterium tuberculosis* isolates cultured in the presence and absence of oxygen. *J Basic Appl Sci.* 2015;11:118–24.