## Retrovirology



Poster presentation

**Open Access** 

# In silico analysis of mobile elements-derived sequences in schizophrenia-related genes

Emanuele F Osimo\*<sup>1</sup>, Marquis Vawter<sup>2</sup>, Steven G Potkin<sup>2</sup>, Fabio Macciardi<sup>1,2</sup> and Simona Gaudi<sup>3</sup>

Address: <sup>1</sup>Department of Science and Biomedical Technology, University of Milan, Milan, Italy, <sup>2</sup>Dept of Psychiatry and Human Behavior, UCI, Irvine, CA, USA and <sup>3</sup>Istituto Superiore di Sanità, Roma, Italy

\* Corresponding author

from Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts Montpellier, France. 21-23 September 2009

Published: 24 September 2009

Retrovirology 2009, 6(Suppl 2):P68 doi:10.1186/1742-4690-6-S2-P68

This abstract is available from: http://www.retrovirology.com/content/6/S2/P68

© 2009 Osimo et al: licensee BioMed Central Ltd.

### **Background**

Regulation of Transposable Elements (TEs) expression has already been associated with complex human diseases, in particular cancer [1]. Also, complex diseases cannot be explained only by genetic factors and are likely to be the result of gene-environment interactions with the contribution of TEs. The detection of retroviral transcripts in the brains of schizophrenics suggests that activation or upregulation of distinct human endogenous retroviruses (HERVs) may play a role in the etiopatho-genesis of neuropsychiatric diseases [2], with increasing complications if we consider that TE insertions generate somatic mosaicism in neuronal cells [3]. In addition, mobile elements have been heavily involved in tissue-specific promoter activity [4] that makes them good candidates for brain specific activation of genes related to schizophrenia. Eventually, TEs are thought to be important in regulation of methylation and DNA accessibility to transcription factors [5]. They are also an important source of small RNAs, which usually act to silence TEs [6]; a particular family of small RNAs, miRNAs, has already been shown to alter neural receptors' function [7].

#### Main aims of the study

Our aim is to investigate the functional relationship between TE-related sequences and genes identified as "best" candidates from genome wide association studies (GWAS) in psychiatric disorders, reconstructing their regulatory network with computational methods, and eventually validating them with exon-expression array. The same will be done independently with TE-related transcriptional start sites.

All this - together with methylation pattern analyses and target sequencing (without repeat masking) that we are planning - could be the start of a novel way of looking at the molecular bases of schizophrenia.

We found preliminary evidence of alternative regulation of tissue-specific gene expression in schizophrenic patients vs controls from genes that we identified as associated to schizophrenia from our GWAS and we present results for both tissue specific expression and alternative TSS in pivotal disease-related genes.

#### **Conclusion**

This *in silico* work is meant to predict and better understand the relationship between TEs and complex disease, paving the way for future experiments that will hopefully confirm and expand current knowledge of what was previously considered just "junk" DNA. This will allow us and others to extract many more data from GWAS by considering also results in previously discarded areas, and we also hope to give a general contribution to the understanding of genome regulation.

#### References

 Goodier JL, Kazazian HH: Retrotransposons Revisited: The Restraint and Rehabilitation of Parasites. Cell 2008, 135:23-35.

- Karlsson H, Bachmann S, Schröder J, McArthur J, Torrey EF, Yolken RH: Retroviral RNA identified in the cerebrospinal fluids and brains of individuals with schizophrenia. PNAS 2001, 98:4634-4639.
- Muotri AR, Chu VT, Marchetto MCN, Deng W, Moran JV, Gage FH: Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. Nature 2005, 435:903-910.
- Faulkner GJ, Kimura Y, Daub CO, Wani S, Plessy C, Irvine KM, Schroder K, Cloonan N, Steptoe AL, Lassmann T, Waki K, Hornig N, Arakawa T, Takahashi H, Kawai J, Forrest ARR, Suzuki H, Hayashizaki Y, Hume DA, Orlando V, Grimmond SM, Carninci P: The regulated retrotransposon transcriptome of mammalian cells. Nature genetics 2009, 41:563-571.
- Ślotkin RK, Martienssen R: Transposable elements and the epigenetic regulation of the genome. Nature Reviews Genetics 2007, 8:272-285.
- Kim VN, Han J, Siomi MC: Biogenesis of small RNAs in animals. Nat Rev Mol Cell Biol 2009, 10:126-139.
- Kocerha J, Faghihi MA, Lopez-Toledano MA, Huang J, Ramsey AJ, Caron MG, Sales N, Willoughby D, Elmen J, Hansen HF, Orum H, Kauppinen S, Kenny PJ, Wahlestedt C: MicroRNA-219 modulates NMDA receptor-mediated neurobehavioral dysfunction. PNAS 2009, 106:3507-3512.

Publish with **Bio Med Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- $\bullet$  yours you keep the copyright

Submit your manuscript here: http://www.biomedcentral.com/info/publishing\_adv.asp

